VOL. 27. Ser. A. Title-page and Index. [Issued November 1940.]

THE REVIEW OF APPLIED ENTOMOLOGY.

SERIES A: AGRICULTURAL.

ISSUED BY THE IMPERIAL INSTITUTE OF ENTOMOLOGY.

LONDON: THE IMPERIAL INSTITUTE OF ENTOMOLOGY. 41, QUEEN'S GATE, S.W. 7.

Price 6s. net.

All Rights Reserved.

IMPERIAL INSTITUTE OF ENTOMOLOGY.

Executive Council.

F. J. DU TOIT, Chairman, Union of South Africa.

Dr. WILLIAM ALLEN, Vice-Chairman, Canada.

Sir Donald Fergusson, K.C.B., United Kingdom.

F. L. McDougall, C.M.G., Australia.

NEVILL L. WRIGHT, F.I.C., D.I.C., New Zealand.

J. M. Adams, F.R.C.Sc. (I), Eire.

D. James Davies, C.B.E., Newfoundland.

Shamaldhari Lal, India and Burma.

B. F. Wright, Southern Rhodesia.

J. A. Calder, C.M.G., Colonies, Protectorates and Mandated Territories.

Sir David Chadwick, K.C.M.G., C.S.I., C.I.E., Secretary.

Director.

Sir Guy A. K. Marshall, C.M.G., F.R.S.

Assistant Director and Editor.
Dr. S. A. NEAVE, O.B.E.

Assistant Director.
Dr. W. R. THOMPSON, F.R.S.

Head Office—British Museum (Natural History), Cromwell Road, London, S.W.7.

Publication Office-41, Queen's Gate, London, S.W.7.

INDEX OF AUTHORS.

A reference in heavy type indicates that a paper by the author has been noticed.

Abbott, W. S., 531, 609. Abraham, R., 100, 113. Adamson, A. M., 55, 56. Aino, S., 442. Ainslie, C. N., 260. Ainsworth, G. C., 671. Akabori, J., 208. Akashi, M., 620. Akimova, A. S., 474. Akita, N., 338. Alden, C. H., 623. Aldrich, J. M., 294. Alibert, H., **347.** Allan, W., 222, 492. Allen, C. E., 541. Allen, M. W., 596. Allen, N., 254. Allen, T. C., 32, 33. Allman, S. L., 345. Almy, E. F., **421.** Ambros, W., 400, **564.** Amer, A., 508. Anagnostopoulos, P. T., 452, 453. Anderson, L. D., 530. Antongiovanni, E., 174. Aoyama, T., 618. App, B. A., 328. Appel, O., 290. Arant, F. S., 82. Araujo, R. L., 22, 23. Arbuthnot, K. D., 80. Arikado, H., 444. Arkhangel'skiĭ, P. P., 474. Arndt, W., 8. Arnim, von, 299. Asquith, D., 79. Austin, M. D., 1, 2, 332. Autuori, M., 34. Avens, A. W., 139. Ayyar, T. V. Ramakrishna, 129,

Babers, F. H., 81. Bach, P. H. De, 597. Back, E. A., 640, 648. Backhouse, J. R. C., 638.

392, 448.

Baerg, W. J., 178. Bailey, S. F., 410. Baker, E. W., 569. Balachowsky, A., 357, 392, 689. Balch, R. E., 368, 647, 662. Ballard, E., 127. Baranov, N., 263. Barber, G. W., 179, 430. Barber, T. C., 416. Barnes, H. F., 320, 353, 556, 635, 672. Barry, J. J., 180. Bartlett, K. A., 670. Basheer, M., 575. Batra, R. N., 486. Baudyš, E., 468. Bawden, F. C., 624. Beadle, L. D., 32. Beaumont, J. H., 188. Bebbington, A. G., 222, 492. Becker, G., 204, 268, 564. Becker, G. G., 53. Becker, W. B., 418. Becnel, I. J., 372. Bedard, W. D., 67. Bedford, H. W., 572. Beeson, C. F. C., 337, 611, 612. Beffa, G. Della, 5. Bégué, H., 224. Bei-Bienko, G. Ya., 315, 453. Bell, A. F., 185. Benton, C., 284. Beran, F., 65, 467. Berezhkov, R. P., 694. Berg, V. L., 543. Bergenstamm, J. E., 294. Bernhardt, E., 495. Berry, P. A., 132, 324. Bertand-Rossi, 224. Bertram, L., 398. Bess, H. A., 598. Besse, H., 478. Betrem, J. G., 522, 523. Bezrukova, T. I., 627. Bezrukova, V. F., 684. Bhatia, B. M., 612.

Bhatia, D., **559.** Bhatia, H. L., 228. Bhatia, M. L., **628**. Bieberdorf, G. A., **257**. Bienko, G. Ya. Bei-, **453**. Bigger, J. H., 67, 285. Billings, S. C., **140.** Bissell, T. L., **76.** Bisset, G. A., 456. Blanton, F. S., 652. Blattný, C., 468, 469, 470. Bleton, C. A., 124, 196. Bliss, C. I., 66, 243, 309, 531. Blockówna, H., 162. Blodgett, F. M., 53. Blunck, H., 100, 314, 334, 495, 608. Blyumberg, G. V., 310. Bobkov, S. D., 309. Boczkowska, M., 105. Bodenheimer, F. S., 336. Bodnár, J., 171, 458. Boelkel, H., 336. Bogunova, M. V., **303.** Böhme, H. Müller-, **158, 357.** Böhmel, W., 300. Bolens, G., 106. Bolwig, N., 518. Bondar, G., 22, 290, 566, 661, 662. Böning, K., 110. Bonnamour, S., 167. Borden, A. D., 432. Borkhsenius, N. S., **57**, **680**. Börner, C., 99, 392, 688. Bos, J., 578. Botkin, C. W., 273. Böttcher, F. K., 101, 231, 402. Bottel, A. E., 239. Bouet, G., 572. Bovey, P., 169, 336. Bovien, P., 117, 518. Bowmaker, P. A., 489, 491. Bowman, B. M., 475. Box, H. E., 182, 659. Boyce, A. M., 55, 261, 425, 432, 459, 649, 650. Bradley, W. G., **80.** Bramstedt, F., 100, 300. Brannon, L. W., 285, 595. Bratley, H. E., 654. Brauer, F., 294. Brédo, H. J., 476, 482, 483, 486. Brémond, P., 193, 359. Breny, R., 236. Brewer, E. G., 138.

Brierley, P., 369. Brightwell, S. T. P., 263. Brindley, T. A., 456. Brink, R. A., 33. Brittin, G., 224. Britton, W. E., 176. Brooks, J. W., **32**. Brown, G. T., 411. Brownlee, J. M., 258. Brubaker, R. W., 250. Bruneteau, J., 690. Brunson, A. M., 341. Brunson, M. H., 84. Bua, G., 98. Buchanan, L. L., 594. Buchanan, W. D., 644. Buchner, P., 401. Bue, I. G., 651. Bugdanov, G. B., 148. Buhl, C., 457, 511 Buikhovtzeva, 318. Burgess, A. F., 139, 615. Burnham, J. C., 104. Burr, M., 455. Busbey, R. L., 456. Busck, A., 188. Bushnell, R. J., 80. Busnel, R. G., 169. Butac, F. L., 183. Butler, C. G., **61**, **265**. Butovitsch, V., **122**, **219**. Buzacott, J. H., 88, 187. Buzin, N., 403. Bynum, E. K., 370.

Cairaschi, E. A., 622. Caldwell, N. E. H., **329.** Calinisan, M. R., 365. Callan, E. McM., **562, 563.** Callenbach, J. A., **32, 33.** Cameron, A. E., 122. Cameron, E., 38, 669. Campbell, R. E., 410. Candioli, P., **235, 515.** Candura, G. S., 172, 198. Canzanelli, A., **198, 199, 234,** 516. Capen, R. G., 280. Cardoso, J. G. A., **479.** Caresche, L., 49. Carnéri, A., 175. Carril, I. M. Del, **476.** Carter, R. H., 280, 456.

Carter, Walter, 624.

Cassab, A., 509. Cassidy, T. P., 416. Cassil, C. C., 188, 280. Castagne, E., **294.** Castets, G., 168. Caudle, H. B., 391. Chaboussou, F., 622. Chamberlain, E. E., 546. Chamberlain, T. R., 32. Chamberlin, F. S., 251, 254. Chamberlin, J. C., **325.** Chandler, S. C., **526.** Chang (Tsung-ping), 667. Chapman, P. J., 382. Charkovskii, M. P., 678. Charles, V. K., 392. Charter, C. F., 384. Chauvin, R., 48, 104, 604. Chen (Chung-liang), 613. Cheo (Ming-tsang), 614. Cherian, M. C., 127, 494, 561, 575. Chernuishev, P. K., 318. Chesnokov, P. G., 439. Chiaromonte, A., 175, 478. China, W. E., 126. Chisholm, R. D., 241. Chiu (Shin-foon), 531, 624. Christian, M. B., 248, 370. Chuginin, Ya. V., 683. Chuvakhin, V. S., 160. Clark, A. F., 329. Clark, J. C., 243. Clausen, C. P., 499. Cleare, L. D., 328, 505, 524. Cockerham, K. L., 249. Cohen, M., 96. Collins, D. L., 77. Compere, H., 39, 503, 506. Compton, C. C., 141. Comstock, J. A., **592.** Cook, M. T., 280. Corl, C. S., 310. Cosolo, S., 199. Costa Lima, A. da, 132, 183. Costantino, G., 5, 516. Cosway, C. A., 224. Cottier, W., 21, 56, 277, 336, 523, 550. Cotton, R. T., 341. Couch, J. N., 242. Couturier, A., 194, 390.

Creighton, J. T., 258.
Cristóbal, U. Lopez, 237, 601.
Crossman, S. S., 615.
Crosthwait, S. L., 532.
Cummins, J. E., 509.
Curl, L. F., 242.
Curran, C. H., 57, 567.
Currie, G. A., 282.
Currie, J. H., 383.
Cushman, R. A., 86.

da Costa Lima, A., 132, 183. da Fonseca, J. Pinto, 34, 86, 238. Dahms, R. G., 261, 456. Dashkevich, B. N., 308. Davidson, J., **345.** Davies, W. M., 550, **562**, 633. Davis, A. C., 79, 260. Davis, J. J., 407, 600. Dawson, J. C., **139.** Day, M. F., **282.** De Bach, P. H., 597. Debauche, H., 486. Decker, G. C., 138. Decoursey, J. D., 75. Deen, O. T., 249. de Fluiter, H. J., 161. Del Carril, I. M., 476. De Leon, D., **567.** Deleuze, J., 349. Della Beffa, G., 5. DeLong, D. M., 82, 421. de Meijere, J. C. H., 94. De Santis, L., 23, 24. de Stefani, T., 452. Diakonoff, A., 50, 205. Dibble, C. B., 408. Dietz, H. F., 532. Distant, W. L., 566. Ditman, L. P., 336, 497. Djou, Y. W., 129. Dmitriev, A. K., 150, 308. Dodonov, B. A., 310. Doeksen, J., 233, 234, 268. Dotti, F., 97, 104, 197. Doucette, C. F., **541**. Douglas, J. R., 412. Douglas, W. A., 537. Dozier, H. L., 539, 643. Drake, C. J., 138. Driggers, B. F., 536. Duarte, A. J., 264, 456, 479. Ducker, H. C., 222. Dugas, A. L., 373. Dumbleton, L. J., 159, 548, 550.

Cox, J. A., 141.

Crandell, H. A., 78.

Crawford, H. J., 538.

Dunnam, E. W., 243. du Plessis, C., 92, 93. Dussy, J., 349. Dyk, A., 400. Dykstra, T. P., 83.

Eastham, J. W., 66. Eckstein, K., 391, 563, 568. Eddy, C. O., **370, 372, 373,** 422. Edwards, W. D., 498. Edwards, W. H., 181, 238, 693. Efimenko, M. S., 407. Eggers, H., 522. Eichler, W., 109. Éidel'man, Z. M., 680. Ellisor, L. O., 258, 370, 371, 372. Emchuk, E. M., **405**, 406. Emel'yanova, N. A., **43**. Emerson, A. E., 384. Emerson, H. E., 55. Endo, T., 17. English, L. L., 259, 642. Enser, K., 171. Esaki, T., 362, 445, 446. Escherich, K., 110, 230. Essig, E. O., 34, 261, 431, 590. Evans, J. W., 20, 21, 184, 226, 510. Everly, R. T., 280, 455. Evlakhova, A. A., 308. Ewing, H. E., 34, 224. Ewing, K. P., 244, 245, 538. Ext, W., 171. Eyer, J. R., 31.

Faes, H., 438. Fahey, J. E., 273, 539, 643. Fahringer, J., 219, 615. Fallscheer, H., 383. Fantham, H. B., 459. Farrar, M. D., 67, 70, 260. Farstad, C. W., 176, 368. Fedorov, S. M., 43. Felt, E. P., 336, 499, 528. Fennah, R. G., 24. Fenton, F. A., 257, 340, 420, 427, 456. Ferla, A. La, 473. Fernando, M., 493, 638. Ferrant, V., 106. Ferrière, C., 506. Ferris, G. F., 290. Feytaud, J., 106, 107, 155, 193, 350, 583, 622.

Fiala, F., 468. Figueiredo jr., E. R., 23. Finck, E. von, **580**, 581. Fink, D. E., 142. Fintzescu, G., 357. Fischer, A., 201. Fischer, H., 353, 466. Fischer, W., 115. Fisher, R. A., 240. Fishkis, I. S., 686. Fitzgerald, D. Vesey-, 335. Flachs, K., **583.** Flanders, S. E., 144, 424, 595. Fleming, W. E., 104. Fletcher, R. K., 413. Flint, W. P., 67, 70, 284, 285, 408, 655. Floyd, E. H., 258. Fluiter, H. J. de, 161. Fluke, C. L., **32.** Folsom, J. W., **341.** Fonseca, J. Pinto da, **34, 86, 238.** Forsslund, K. H., **518.** Francke-Grosman, H, 401. Fransen, J. J., 104, 396. Franssen, C. J. H., 227, 280, 523. Franzke, A., 204. Frappa, C., 14, 62, 543, 637, 664. Frear, D. E. H., 52. Freeman, P., **360.** Freitag, J. H., **590.** Freudenstein, K., 101. Frey, W., 635. Frézal, J., **191.** Frickhinger, H. W., 606. Friend, R. B., 73, 342. Froggatt, J. L., 277, 544. Frost, S. W., 80. Fukaya, M., 619. Fukaya, S., 442. Fukuda, J., 620. Fukuda, S., 443. Fuller, M. E., 56, 57. Fulmek, L., 45. Fulton, R. A., 78. Furniss, R. L., 498. Furugori, N., 225. Fyfe, R. V., 89, 282.

Gäbler, H., 218, 334, 400, 579. Gahan, A. B., 434, 565. Gaines, J. C., 242, 245, 413, 417. Gaines, R. C., 243, 246. Gambrell, F. L., 529. Garman, P., 428, 429. Garthwaite, P. F., 692. Garus, J., 45. Gauze, G. F., 161. Gay, F. J., 89. Gayden, J. H., 258. Geijskes, D. C., 301, 567. Geisthardt, G., 521. Ghesquière, J., **572.** Ghidini, G. M., **174.** Gibson, A., 476, 477, 485. Gibson, K. E., 418. Gieysztor, M., 438. Gillett, J. A., 412. Gilmer, P. M., 247. Gilmore, J. U., 252, 253. Gilyarov, M. S., 44, 679. Giorgi, D., 198. Girault, A. A., 294. Girth, H. B., 165. Glaser, R. W., 165. Glasgow, H., **531.** Gnadinger, C. B., 310. Gobeil, A. R., 275, 463. Goetsch, W., 87. Goidanich, A., 173, 174, 472. Goncharova, A. A., 311. Gontarski, H., 630. Gooden, E. L., 541. Goodhue, L. D., 241, 528, 541. Goodliffe, F. D., 562. Goodwin, M. W., 534. Gorham, R. P., 601. Gösswald, K., 112, 231, 335, 688. Götz, B., **7, 496, 582.** Gould, E., **188.** Goux, L., 48, 103, 235. Goytia, A., **543.** Gradojević, M., **479.** Graham, A. B., 408. Graham, C. P., 259. Graham, J. J. T., 392. Graham, L. T., 370. Graham, S. A., 527. Grandi, G., 473. Grandori, L., 199. Grandori, R., 197, 200. Gray, K. W., 325. Gregor, F., 145. Grigor'eva, T. G., 317. Grison, P., 622. Griswold, C. L., 651. Grivanov, K. P., 554. Grosmann, H. Francke-, 401. Groves, K., 383. Grunberg, A., 512.

Gui, H. L., 342.

Guichard, F., **347.** Gunn, D. L., **224.** Gurney, W. B., **130.** Guy, H. G., **532.**

Haasis, F. A., 652. Hadersold, O., 113, 352. Haenseler, C. M., 536. Hafiz, H. A., 385. Hahmann, K., 298. Hall, W. J., 14. Haller, H. L., 135. Haller, M. H., 188. Hama, T., 619. Hamiel, G. R., 273. Hamilton, C. C., 74. Hammer, O. H., 533. Hanna, A. D., 512. Hanno, K., 400. Hansen, J. W., 649. Hansen, L. L., 31. Hanson, A. J., 178. Hanson, H. S., 504. Hardy, A. C., 167. Hardy, D. E., 596. Hardy, G. H., 294. Hardy, J. E., 262, 392, 670. Harel, 10. Harmston, F. C., 29, 427. Harrington, C. D., 33. Harris, C. S., 280. Harris, K. H., 342. Harris, W. V., **492**. Harrison, P. K., 250, **371**. Hartley, B. J., 222. Hartnack, H., 382. Hartzell, A., 272. Hartzell, F. Z., 529, 534. Harukawa, C., 624. Hase, A., **354.** Hatch, M. H., 78, 525, 592. Haug, G. W., 81. Haupt, W., 158. Hawkins, J., 650. Hawkins, L. A., 53. Hayes, W. P., 75. Headlee, T. J., 137. Hean, A. F., 360. Heikertinger, F., 160. Heinrich, C., 511. Heinze, K., 157, 267, 334, 636, 686, 687. Heiss, E. M., 82. Helson, G. A. H., 639. Hely, P. C., 278.

Hempel, A., 48. Hennig, W., 113. Henrard, P., 572. Hensill, G. S., 409. Hers, J., 486. Hespeler, O., 203. Hey, G. L., 236. Hicock, H. W., 73. Hildebrand, A. A., 276. Hille Ris Lambers, D., 224, 578. Hilmy, A. L., 506, 508. Hiraiwa, Y., 440. Hirayama, S., 444. Hixson, E., 79. Hockenyos, G. L., 422. Hodson, A. C., 597, 646. Hodson, W. E. H., 512, 519. Hoffmann, A., 225. Hofmann, C., 48, 231, 512. Hogan, T. W., 389. Holloway, J. K., 426. Hopkins, A. D., 325. Hopkins, J. C. F., 125. Hora, A. M., 585. Hori, H., 446. Hori, M., 104. Hornbostel, W., 334. Hoskins, W. M., 411. Hough, W. S., 533. Houser, J. S., 83. Howe, E. W., 500. Hsiao, Tsai Yu, 643. Huber, G. A., 280. Hudault, E., 691. Huggins, H. C., 116. Hughes, A. M., 456. Hughes, T. E., 456. Huitema, W. K., 624. Hukkinen, Y., 160. Hülsenberg, H., 513. Hundertmark, A., 223. Hunt, G. M., 180. Hunter, L., 237. Hunter, W. P., 258. Husain, M. A., 560. Husni, M., 508. Hussein, M., 481. Hutchinson, H., 488. Hutson, J. C., 638. Hyslop, J. A., 137.

Il'inskaya, L., 304. Imamura, S., 555. Immelman, M. N. S., 568. Ingram, J. W., 372, 656. Inouye, M., 91, 339.
Isaac, P. V., 575.
Isaakidès, C. A., 171, 478.
Isely, D., 165, 178, 327.
Ishibashi, R., 440.
Ishii, T., 91, 620.
Ishikura, S., 557.
Ishitani, F., 443.
Isitani, H., 208.
Ivanov, S. P., 406.
Iwasa, T., 616.
Iyriboz, Nihat, 452.

Jack, R. W., 124. Jacoby, M., 271, 567. Jacques, C., 131. Jancke, O., 111, 155, 458, 689. Janes, M. J., 651. Janisch, E., 98, 314, 564. Janjua, N. A., 575. Jannone, G., 570. Jary, S. G., 1, 2, 332, 519. Jaynes, H. A., 372, 656, 658. Jeffreys, F. J., 37, 548. Jepson, W. F., 611. Jeż, S., 350. Jobbins, D. M., 137. Johansson, E., 121. Johnpulle, A. L., 493. Johnson, A. C., 53. Johnson, G. V., 420. Jones, E. P., 14, 15, 16. Jones, E. T., 374, 375. Jones, G. D., 569. Jones, M. P., 408. Jones, S. C., 498. Jung, K., 104.

Kabakchi, B. P., 152. Kabos, W. J., **494.** Kaburagi, T., 441. Kac, 471. Kadow, K. J., 534. Kagan, M., 261. Kagy, J. F., 649, 653. Kalandra, A., 469. Kalashnikov, K. Ya., 684. Kalshoven, L. G. E., 48, 227, 493. Kamiński, E., 9. Kamiya, K., 209, 211. Kanda, S., 624. Kangas, E., 8. Kariya, S., 361, 616. Karlash, K., 403.

Karr, E. H., 648. Kaston, B. J., 68. Katz, Ya., 403. Kaufmann, O., 203, 204. Kaven, G., 232. Kawaai, I., 225, 555. Kawano, T., 339, 360. Kearns, C. W., 141. Kearns, H. G. H., 396. Keifer, H. H., 33, 288. Kéler, S., 162. Kelsall, A., 662. Kemp, H. K., 292. Kemper, H., 101, 201, 513. Kerkow, 203. Kerr jr., T. W., 425. Kharitonov, Ya., 683. King, K. M., 543. King, M. E., 395. Kirchner, H. A., 457. Kirkpatrick, A. F., 343. Klemm, M., 336. Klingstedt, H., 522. Klinkowski, M., 495. Knight, Harry H., 566. Knowlton, G. F., 29, 31, 48, 426, 427, 596. Knust, H. G., 88. Kobayashi, M., 210. Köhler, E., 156. Koidsumi, K., 615. Kojima, T., 441. Kolotov, D. G., 454. Kolubajiv, S., 517. Komárek, J., 170. Kôno, H., 19, 91, 211, 339, 443, 615. Koreishi, K., 206. Korsakov, M. N. Rimskii-, 214. Kostandyan, L. V., 320. Kovačević, Z., 10. Kowalczyk, S. A., 392. Koyama, T., 207, 361. Kozhanchikov, I. V. (Kozhantshikov, I. W.), 37, 147, 228. Kozikowski, A., 676. Kozlova, E. N., 41. Kreiter, E. A., 454. Krivin, B. G., 303. Kumazawa, T., 212. Kunike, G., 6, 47, 64, 399. Kurdyumov, N. V., 311. Kurentzov, A. I., 456. Kuroda, H., 620. Kuroda, M., 207, 443. Kuroda, S., 445.

Kurosawa, M., 50, 616. Kuryllo, A., 162. Kuwayama, S., 19, 340, 440, 618. Kuz'min, M., 322. Kuznetzov, 318. Kuznetzova, L. A., 313. Kvičala, B., 145.

Laan, P. A. van der, 17. Lacotte, 357. Lacroix, D. S., 177. La Ferla, A., **473.** Lafon, M., 624. LaHue, D. W., 427. Laing, J., 166. Lal, K. B., 385, 575. Lamas, C., J. M., 385. Lambers, D. Hille Ris, 224, 578. Lamerson, P. G., 165, 166. Lamiman, J. F., 409. Langford, G. S., **532.** Langston, J. M., **644.** Latta, R., 419. Laue, G., 102. Lazarevskii, M., 403. Lea, A., **62, 387.** Lean, O. B., 13. Lebedev, M. I., 623. Leece, C. W., 87. Leefmans, S., 514. Lehker, G. E., 408. Lehmann, H. C., 457. Leib, E., 204. Leon, D. De, 567. Leonard, M. D., 531. Le Pelley, R. H., 665. Lepesme, P., 196, 486, 584, 585. Lesnikovskii, G. M., 680. Leuzinger, H., 169. Lever, R. J. A. W., 212, 213, 365, 366, 641. Lima, A. da Costa, 132, 183. Lin (Ch'ang-Shan), 667. Lindblom, A., 355. Lindemuth, K., 511. Lindgren, D. L., 343. Lindinger, L., 223. Lindsay, E. A., 510. Linsley, E. G., 160. List, G. M., 72, 408. Little, E. C. S., 22. Liu (Chi-ying), 613, 614. Lizer y Trelles, C. A., 87. Lloyd, D. C., 302.

Lobdell, R. N., 656.
Lombardi, D., 471.
Lopez Cristóbal, U., 237, 601.
Lounsky, J., 605.
Lovibond, B., 95.
Lowman, M. S., 134.
Lucchese, E., 4, 629.
Ludwig, D., 336.
Lukash, I. I., 322.
Luk'yanovich, F. K., 313.
Lupo, V., 48, 511.

Ma (Tung-lwen), 614. McCay, C. M., 624. McClean, H. C., 642. McCoy, E. E., 165. McCrory, S. A., 133. McDaniel, E. I., 28, 461. McGarr, R. L., 244. MacGill, E. I., 672. McKay, M. B., 369. McKinney, K. B., 142. McKinstry, A. H., 222. McLane, L. S., 416. McNally, A. G., 601. McOmie, W. A., 597. Madden, A. H., 251. Madel, W., **7, 521.** Mader, E. O., **53.** Maeda, T., 210. Maercks, H., 519. Magie, R. O., 593. Mahdihassan, S., 113. Makino, M., 91. Makkus, W., 158. Malenotti, E., 115, 295. Malloch, J. R., 294. Mannheims, B., 398. Marcovitch, S., 104, 425, 426. Marcu, O., 355. Mariani, G., 197. Mariani, M., 630. Marloth, R. H., 330. Marshall, James, 383. Marshall, Joseph, 487. Martelli, M., 173, 174. Martin, C. H., **68, 190.** Martin, H., 396. Marushev, A. I., **554.** Masera, E., 224. Masi, L., 199. Mason, T. G., 220, 222. Massee, A. M., 160, 395, 636. Mathewson, A. A., 287. Mathlein, R., 118. Mattras, H., 357.

Maxwell, K. E., 55. Mayné, R., **106**, **107**. Meĭer, N. F., **588**. Meijere, J. C. H. de, **94**. Melis, A., 296. Melville, A. R., 665. Mendes, L. O. T., 34, 188, 289. Menozzi, C., 301. Mercet, R. G., 40. Merkel, L., 298. Mesnil, L., 167, 624. Metalnikov, S., 307. Metcalf, C. L., 655. Metzger, F. W., 104. Meyer, E., 511, 608. Miao, C. P., 360. Michelbacher, A. E., 411, 431, 569. Micklem, T., 94. Middelburg, H. A., 493. Middlekauff, W. W., 279. Middleton, W., 53, 138. Miestinger, K., 467. Milam, J., **652.** Miles, W. H., **96.** Milisavl'evic, D., **494.** Miller, D., 37, 449. Miller, F. W., 392. Miller, L. W., 293. Milne, P. S., 167. Minkiewicz, S., 163, 694. Misaka, K., 18, 206. Mistikawi, A. M., 481. Mitchell, B. L., 488. Miwa, Y., 205, 558. Miyake, T., 208, 617. Mizutani, Y., 209. Mo, Tjoa Tjien, 347. Modena, M. P., 199. Monro, H. A. U., 416, 541. Moore, M. H., 456. Morae, C., 238. Moreland, R. W., 244, 417. Moretti, G. P., 199. Morgan, W. L., 57. Morgenthaler, O., 169. Morimoto, S., 361. Morita, T., 557. Moriyama, C., 557. Moriyama, T., 618. Moroshkina, O. S., 474. Morrill, A. W., 177. Morris, H. M., 270, 317. Morrison, H., 392. Morrison, L., 547. Moss, E. G., 254.

Mossop, M. C., 125, 637. Mote, D. C., **643**. Motzkus, E., 203. Muesebeck, C. F. W., 133, 324, **461,** 614. Mühle, E., 466. Mukerji, D., **280.** Mukerji, S., **486.** Müller, H., **520.** Muller, H. R. A., **280.** Muller, Klothilde, 230. Müller-Böhme, H., 158, 357. Munger, F., **419**. Mungomery, R. W., 88. Munro, J. A., **33, 70, 71, 651, 655.** Munro, J. W., 265, **336.** Murachev, M., 153. Murat, M., 487. Murphy, A. M., 286. Murray, C. W., 188. Murray, F. V., 104.

Nägeli, W., 518. Nakayama, S., 225, 362. Narayanan, E. S., 558. Nash, R. W., **540**. Nawa, U., **207**, **361**. Nefedov, N. I., **386**. Negrul', A., 403. Nelson, O. A., 643. Nepveu, P., **621.** Nettles, W. C., **408.** Neu, W., 608. Nevskii, V. P., 552, 587, 610. Nichols, R. F. W., 153. Nicholson, A. J., 476. Nikitina, T. F., 316. Niklas, O. F., 581, 632. Nikolskiĭ, V. L., 152. Nitsche, G., 688. Nixon, G. E. J., 19, 263, 505, 604, 632, 669. Noble, N. S., **293, 448, 564.** Noble, R. J., 448. Noguchi, T., 440. Nolte, H. W., 170, 267, 298, 335, 401, 466. Norris, K. R., 131. Notini, G., 117. Notley, F. B., 127. Novienko, A. I., 318. Nowell, W., 290. Nowicki, S., 232, 403. Nozu, R., 555, 556. Nunberg, M., 676.

Obarski, J., **162.** Odagaki, S., 617. Odera, S., **617.** Ogijewicz, B., 350. Ogilvie, L., **671.** Ohta, Y., 210, 339. Oishi, T., 440. Okamoto, D., 619. Oliver, R. W., 25. O'Neill, W. J., 536. Onoe, T., 208, 620. Oosterling, H., **577.** Orelli, O. Schneider-, **160.** Osborn, H., **527**. Osborn, H. T., 370. Osborn, W. M., 274. Osburn, M. R., **256, 257.** Osima, K., 618. Osterberger, B. A., **248**, **370**, **371**. Otero, J. I., 280. Oudemans, A. C., 301. Ovchinnikova, M. I., 151. Ovinge, A., 233, 465. Ozaki, S., 620. Ozols, E., 47.

Paĭkin, D. M., 149. Painter, R. H., **341.** Palmer, M. A., 104, 542. Paniagua, S., 477. Pape, H., 437. Pappenheim, E., 49. Parfent'ev, V. Ya., **321.** Parham, B. E. V., **213.** Parisi, E., **199.** Park, M., 493. Park, T., 27. Parker, J. H., **274.** Parker, J. R., 240. Parker, R. L., 165, 166. Parks, T. H., 83, 407. Parnell, F. R., 221. Parsons, F. S., 222, 223, 487, 488. Pasquier, R., 480. Patay, R., 584. Patch, L. H., **645**. Paterson, N. F., 60. Paul, L. C., 543. Pawłowicz, J., 438. Peacock, A. D., 176. Pearce, G. W., 139. Pearson, E. O., 489. Peat, J. E., **491.**

Pechuman, L. L., 77. Peirson, H. B., 275. Pemberton, C. E., 90, 366. Pengelly, R. C., 27. Pepper, B. B., 179, 536. Persing, C. O., 425, 649, 653. Pescott, R. T. M., 90, 187, 389. Petch, C. E., 664. Peters, G., 11, 201. Petherbridge, F. R., 673. Peus, F., 102. Pfeffer, A., 469. Phillips, W. J., 280. Phillips, W. R., 541. Phillis, E., 220, 222. Pickles, A., 24, 543. Pierce, W. C., 255. Pierpoint, R. L., 532. Pilat, M. V., 307. Pinto da Fonseca, J., 34, 86, 238. Pitcher, R. S., 2. Plank, H. K., **423.** Plessis, C. du, **92**, **93**. Plumb, G. H., 73. Podkanowicz, J., 9. Poeteren, N. van, 106, 269. Polezhaev, V. G., 551, 585. Ponomarenko, D. A., 108. Ponomareva, L. M., 304. Poole, R. F., 254. Poos, F. W., 280. Popova, A. I., 315. Popova, E. I., 303. Porra, J., 392. Pospelov, V. P., 306, 455. Pospelova, V. M., 678. Potter, C., 154, 623. Potts, S. F., 143, 645. Poutiers, R., 194. Powell, H. R., 130. Prashad, B., **391.** Pravdin, F. N., **44.** Prell, H., 101. Prendergast, D. T., 459. Prentice, A. N., 222, 491. Priesner, H., 507. Prikhodkina, T. D., 677. Pringle, J. A., **60.** Printz, Ya., 403. Profft, J., 267, 301, 633, 636. Prüffer, J., 9, 162. Prutenskii, D. I., 153. Pruthi, H. S., 50, 228, 558. Pussard, R., 266, 458, 621. Pyenson, L., 86, 136, 413.

Quanjer, H. M., **624.** Quayle, H. J., **342.**

Rademacher, B., 334. Rahman, K. W., **503.** Rainey, R. C., **489.** Rainwater, C. F., **246.** Ramakrishna Ayyar, T. V., 129, 392, 448. Ramchandra Rao, Y., 484, 485, 559, 624. Rau, G. J., 48. Raucourt, M., 168, 675. Ray, W. W., 427. Reed, L. B., 240, 249, 287. Reed, R. H., 260. Regueral, F. G., 6. Reichmuth, W., 102. Rekach, V. N., 215. Reppert, R. R., 408. Richardson, C. H., 134, 537. Riggs, D. S., 68. Rimskii-Korsakov, M. N., 214. Ris Lambers, D. Hille, **224**, **578**. Ritcher, P. O., **32, 592.** Roark, R. C., **78, 224, 285, 392,** 538, 593, 694. Roberti, D., 512. Robertson, W. H., 66. Rodd, A. E., 609. Rodd, V. E., 312. Rodendorf, B. B., 439. Rodionov, Z. S., 551, 585. Roebuck, A., 674. Roesler, R., 689. Rogers, W. S., 395. Rolfe, S. W., 3. Romanova, V. P., 304. Romney, V. E., **422**, **593**. Roos, K., **270**. Rosenberg, H. T., 38. Ross, A. A., 186. Ross, E., 569. Rossi, Bertand-, 224. Rossi, L., 104. Rubtzov, I. A., 145, 311, 313, 554. Ruffo, S., **471, 473.** Ruhmann, M. H., 67. Rungs, C., 193, 392, 691. Rusk, H. W., 539. Russo, G., 568.

Ruszkowski, J., 162.

Saalas, U., 267, 465. Saito, H., 617. Saito, S., 209, 616. Sakai, K., 207, 445. Sakharov, N. L., **109**, 406, **680**. Salaman, R. N., **236**. Sales, E. M., **434**. Salman, K. A., **140.** Salt, G., **35, 390.** Sameshima, T., **362.** Sanders, F. R., **126.** Santis, L. De, **23, 24.** Saraiva, A. C., **479, 484.** Sattler, H., **398.** Sauer, F. G., **22.** Saugstead, S., 33. Saunders, L. G., **27.** Savchenko, E. N., 406. Savitz'ka, Z., 404, 406. Sawa, R., 209. Sawamoto, T., **443.** Sazama, R. F., 273. Sazonov, P. V., 308. Scales, A. L., **246.** Schaefer, C. A., **323.** Schaefer, E. E., **91.** Schaerfenberg, B., 398. Schaper, P., 107, 158. Schechter, M. S., 135. Schedl, K. E., 7, 105, 219, 397, 512. Schifino, L. A., 70, 71. Schilder, F. A., **688.** Schimitschek, E., 218, 522. Schiuma, R., **213.** Schmidt, E., **224.** Schmidt, G., **437.** Schmidt, H., 114. Schneider, F., 385. Schneider-Orelli, O., 160. Schrock, M., 240. Schuch, K., 112, 202, 204, 520, 520.Schultz, E. F., 238. Schulze, B., 203. Schulze, K., **521.** Schumacher, F., 218. Schwartz, M., 158, 193. Schwartze, C. D., 280. Schwarz, C. L., 203. Schwarz, M., 106, 107. Schwerdtfeger, F., 160, 299, 300, 579, 606. Scleros, N., **453.** Scott, L. B., 652.

Seamans, H. I., 368.

Searls, E. M., 33, 415. Seaton, L., 32, 287, 423. Séguy, E., 113, 114. Seiferle, E. J., 134, 537. Selivanova, S. N., **305.** Sellke, K., **7, 634.** Sen, H. K., **128.** Sen, P., **627.** Serrano, L. A., 436. Servadei, A., 3. Severin, H. H. P., **590.** Seydel, C., 294. Shabliovskiĭ, V. V., **42.** Shadle, A. R., **541**. Shafik, M., **506**, **508**. Shands, W. A., **254**. Sharp, S. S., **372, 645.** Shaw, H., **330, 456.** Shcherbinovskii, N. S., 681. Shepherd, B. L., **374.** Shevchenko, M. I., 147, 440. Shibuya, T., 616. Shil'onok, A. A., 610. Shinji, O., 104, 224. Shiraiwa, H., 616. Shmal'ko, V. S., 685. Shotwell, R. L., 76, 139, 240. Sidel'nik, I., **321.** Sidorovnina, E. P., **305.** Siebenmann, 517. Siegler, E. H., 419, 528. Silayan, H. S., 451. Silvestri, F., 34, 48, 478, 568, 623. Simmonds, H. W., 22. Sinel'nikova, Z. S., **552, 623.** Siraiwa, H., 103, 456. Sison, P., 364. Skoblo, I. S., 42, 682. Skryabina, E. A., 308. Slate, W. L., **176.** Sloan, W. J. S., **187, 281.** Smallman, B. N., 52. Smee, C., **125.** Smirnov, E. S., 551. Smirnova, O. N., **683.** Smit, C. J. B., **393**. Smith, A. J., **360.** Smith, C. E., **250, 370.** Smith, C. F., 29, 455. Smith, F. F., 79. Smith, G. L., 246. Smith, J. H., 89, 365, 602. Smith, K. M., 456, 671. Smith, L. E., 142, 419. Smith, L. M., 139. Smith, R. C., 138, 374, 375.

Smith, R. H., 143. Smith, T. C., 140. Smolák, J., 512. Snapp, O. I., 255. Snelling, R. O., **274.** Snyder, T. E., **180.** Sokolov, A. M., **684.** Soliman, L. B., 160. Sonan, J., 206, 339, 340, 446, 447, 618. Soraci, F. A., 29, 279. Sorauer, P., 113, 290. Sorenson, C. J., 29, 48, 462. Sorokin, S., **319.** Spencer, H., 256, 257. Spessivtseff, P., 123. Speyer, W., **99, 112, 514.** Spoon, W., **336, 465.** Squire, F. A., 222, 223. Stahel, G., **567.** Stahler, N., **423.** Stanley, J., **52, 175.** Stanley, W. W., **104, 426.** Stearns, L. A., **534.** Steer, W., **331, 456.** Stefani, T. de, 452. Steinegger, P., **46.** Steiner, H. M., 535. Steiner, L. F., 272, 643, 653. Steiner, P., 520, 583. Stellwaag, F., **102**, **465**, 689. Stepantzev, I. N., **147**. Stepniewska, K., 675. Stewart, R., 96. Stirrett, G. M., 25. Stofberg, F. J., 330. Stone, M. W., 410. Storey, H. H., **94, 153.** Storey, I. F., 672. Störmer, 157. Stracener, C. L., 248, 370. Straňák, F., **468.** Strawiński, K., **163.** Strickland, A. G., 164, 292. Strickland, E. H., **175, 539.** Stritt, W., 6. Strom, L. G., 336. Strong, L. A., **261**, **376**. Subklew, W., **224**, **694**. Subramaniam, T. V., 291, 625. Sugak, A., **150.** Sugihara, Y., **211.** Sugimura, R., 556. Sullivan, W. N., **134.** Summerland, S. A., 374. Sundaram, C. V., **561.**

Suster, P. M., 608. Sviridenko, P. A., 568. Swain, A. F., 343. Swain, R. B., 528. Sy, M., 114, 159. Szymański, W., 162.

Tadasa, Y., **340.** Takahashi, H., 206, 339, 446, 447. Takahashi, R., 103, 205, 224, 336, 557, 624. Takahashi, T., 210. Takana, S., 363. Tamanuki, K., 615. Tammes, F. M. L., 478. Tamura, I., 210, 338, 445, 556. Tanaka, K., 617. Tanko, B., 171. Taranukha, M. D., 456. Tarwid, K., 161. Tattersfield, F., 694. Taylor, G. G., **546.** Taylor, T. H. C., 568. Teissier, G., 624. Telenga, N. A., 303. Tempel, W., 354. Tenhet, J. N., 500. Terui, M., 225. Thalenhorst, W., 65, 354. Thielmann, K., 216. Thiem, H., 46, 112, 114, 155, **297, 298, 299,** 507, **513.** Thomas, C. A., 511. Thompson, W. R., 436. Thornley, H. F., 29. Thorpe, W. H., 264, 391. Thung, T. H., 493. Tiburec, 470. Tiegs, O. W., 104. Tihenko, V. J., 409. Tilemans, E., 605. Timokhina, N. I., 626. Tischler, W., 63, 352, 631. Tjoa, Tjien Mo, 347. Todd, J. N., **69.** Tokunaga, M., 624. Tompkins, C. M., 273. Tonnoir, A. L., **57.** Toshevikova, A. G., 309. Townsend, C. H. T., **22, 143.** Townsend, J. F., 429. Trägårdh, I., 219, 514. Trappmann, W., 100, 202. Trehan, K. N., 560.

Treiman, F. S., 404.
Trelles, C. A. Lizer y, 87.
Treml', 318.
Trinchieri, G., 481.
Trouvelot, B., 107, 168, 193, 357.
Tsai Yu Hsiao, 643.
Tsuchiya, K., 338.
Tsumagri, H., 210.
Tucker, R. W. E., 660.
Turaev, N. S., 454.
Turner, N., 71, 456.
Turner, P. E., 384.
Tziopkalo, V. L., 406.

Uichanco, L. B., 478, 485.
Ullyett, G. C., 323.
Underhill, G. W., 141.
Uspenskiř, F. M., 625.
Uvarov, B. P., 138, 336, 474, 475, 522, 604.
Uyttenboogaart, D. L., 223.

Vampler, E. L., **411.** Vance, A. M., 414. van der Laan, P. A., 17. Vanderwalle, R., 605. van Poeteren, N., 106, 269. van Zwaluwenburg, R. H., 90. Vappula, N. A., 694. Vasser, R. É., 148, 313. Vayssière, P., 358, 486. Veitch, R., 87. Vel'tishchev, P. A., 321. Venables, E. P., **511**. Venturi, F., 3. Verma, P. M., 560. Vesey-Fitzgerald, D., 335. Vidal, J. P., 570. Vielwerth, V., 468. Viktorov, V. F., 152. Vinokurov, G. M., 153, 320. Vinson, C. G., 133. Vogel, R., 46. Volgin, V. I., 319. Volkov, V. F., 307. Volterra, V., 161. Voskresenskaya, A. K., 310. Vukasović, P., 515.

Wahlen, F. T., **350.** Wahlgren, E., **160, 512.** Wakeland, C., **412.**

Wakely, S., 295. Walker, G. L., 71. Walker, H. G., **530.** Wallace, G. B., **388**, **492**. Wallace, J. M., **286.** Walters, E. V., 427. Walton, E. V., 287. Walton, R. R., 427. Walton, W. R., 240. Ward, K. M., 89, 186. Wardlaw, C. W., 288. Wardziński, K., 232. Wason, E. J., 57. Watanabe, C., 338, 614, 615, 620, 624. Waters, H. A., **421.** Waterston, J. M., **85.** Watson, E. B., **25.** Watson, J. R., 654. Watzl, O., 635. Webb, D. van V., 387. Weber, H., 170. Webster, R. L., 178, 525. Weddell, J. A., 186. Weidner, H., 201. Weiland, G. S., 336. Wellington, J. S., 271. Welsch, J., 113, 507. Wever, A. L., 642. Whelan, D. E., 600. Whitaker, W. C., 83. Whitcomb, W. D., **496.** Whitehead, T., 633. Whittington, F. B., 532. Wickens, G. M., 226. Wiesmann, R., 232, 280. Wigglesworth, V. B., 436. Wilcoxon, F., 272. Wilkinson, D. S., **37, 57, 505.** Wilkinson, H., 666. Wille, J. E., 164, 385, 511. Williams, C. B., **608.** Williams, F. X., 90. Williams, J. L., 456. Willis, H. R., 646. Wilson, C. C., 569. Wilson, F., 59, 262. Wilson, G. F., 115, 295. Wilson, H. F., 31, 32. Winning, E. von, 218. Wisecup, C. B., 249, 250, 287. Witman, E. D., 421. Woke, P. A., 241, 341, 524. Wolcott, G. N., 84, 85, 424, 434, 464.

Wolfenbarger, D. O., 644. Woodworth, C. E., 28. Worth, C. B., 655. Worthley, H. N., 52, 535. Wright, D. W., 116, 674. Wright, E., 284. Wünn, H., 223, 402.

Yagi, N., 442. Yago, M., 225, 441. Yamasaki, T., 446. Yamazaki, M., 444. Yanagihara, M., 363. Yashiro, H., 444. Yasumatsu, K., 446. Yokoo, R., 557. Yokoo, T., 337. Yuasa, H., 17, 556. Yu Hsiao, Tsai, 643.

Zaaijer, J. W., 624.
Zacher, F., 8, 219, 220, 355, 513.
Zaitzeva, A. Ya., 307.
Zakharov, L. Z., 109, 553.
Zakhvatkin, A. A., 1.
Zelensky, V., 691.
Zeller, S. M., 498.
Zenyakin, L. A., 40.
Zetterstedt, J. W., 512.
Ziegler, L. W., 653.
Zolotarev, E. Kh., 586.
Zolotarevsky, B. N., 11, 280, 481, 484, 570, 571.
Zvoruikina, N. A., 587.
Zwaluwenburg, R. H. van, 90.
Zweigbaumówna, Z., 162.

GENERAL INDEX.

In the case of scientific names, the page reference is cited only under the heading of the generic name.

When a generic name is printed in brackets, it signifies that the name is not the one adopted.

Abagrotis vetusta (see Agrotis). abboti, Oiceticus; Sphecodina.

abbreviatus, Diaprepes.

abdominalis, Eutrichopoda; Macrocentrus.

Abies, restrictions on importation of, into Britain, 356; pests of, in Germany, 48, 401; Chermes nordmannianae on, in Switzerland, 517; Acantholyda erythrocephala not observed U.S.A., 651. on,

Abies alba, pests of, in Czecho-

slovakia, 469.

Abies concolor, pests of, in U.S.A., 284, 324, 567.

Abies grandis, Chermes piceae on, in U.S.A., 499.

Abies sachalinensis, pests of, in Japan and Sakhalin, 19, 211, 339, 615.

abietina, Pristiphora (Nematus). abietinus, Dryocoetes.

abietis, Chermes (Adelges); Diprion (Neodiprion); Megastigmus.

Ablerus clisiocampae, parasite of Malacosoma disstria in Minnesota, 597.

abrupta, Empoasca.

abruptorius, Exenterus.

abscissor, Microbracon (Bracon).

abyssinialis, Hypena. abyssinicus, Anagyrus.

Acacia, Ochrogaster contraria on, in 545; Anomala Australia, expansa on, in Formosa, 363; Icerya purchasi on, in Morocco, Aphis laburni on, Russia, 216; Coleoptera in timber of, in S. Africa, 60.

Xystrocera mollissima, globosa on, in Formosa, 558.

Acacia, White (see Robinia). Acacia, Yellow (see Caragana).

varinervis Antholcus introduced into New Zealand against, 449.

Acalypha senensis, Helopeltis bergrothi on, in Sudan, 574.

Acanthocoris sordidus, bionomics and control of, in Japan, 446.

acanthodactyla, Platyptilia.

Acantholyda erythrocephala, bionomics and control of, on Pinus spp. in U.S.A., 279, 651.

Acantholyda nipponica, on pine in Japan, **339.**

Acanthoscelides (see Bruchus).

Acarapis woodi, attacking honeybees in Switzerland, 169.

Acaulona peruviana, parasite of Dysdercus ruficollis in Peru, 164. Acer (see Maple).

Acer macrophyllum, pests of, in U.S.A., 498, 499.

Acer negundo, Leptocoris trivittatus on, in Kansas, 374.

Acer saccharum, Malacosoma disstria on, in U.S.A., 377.

Aceratoneuromyia australia, type of (See Syntomosphygenus, 434. rum indicum.)

aceris, Cryptococcus; Phenacoccus; Rhinocola.

Acetylene, no reaction of Melolontha to, 200.

Achaea janata, banding against, on trees in Burma, 693.

achemon, Pholus.

Acherontia atropos, on potato in Britain, 332; on olive in Egypt,

Achroia grisella, induced response in Nemeritis canescens to odour of, 264, 265; penetration of fluids into tracheae of, 580.

Achrysocharella ruforum, parasite of Diprion pini in Danzig and E.

Prussia, 105.

Achrysopophagus aegyptiacus, reared from Coccinellids in Egypt, 40; possibly reared from Pseudococcus spp. in S. Africa and Tanganyika, 40.

Acidol, 627.

acidusa, Anastrepha. acinaciformis, Coptotermes. Acolastus amyntas, on elliptica in Porto Rico, 502.

acontioides, Lyncestis.

acraea, Estigmene.

acridiorum, Hexamermis.

Acritocera negligens, on coconut in Fiji, 212.

Acrobasis caryae, thrips predacious on, in Louisiana, 372.

Acrocercops cramerella, on cacao in Philippines, 452.

Acromyrmex, suggested measures against, in Argentina, 87.

aculeatus, Haplothrips. acuminata, Aelia.

acuta, Cantheconidea.

Acyrthosiphon (see Macrosiphum).
Adalia bipunctata, predacious on
Anuraphis roseus in Connecticut,
428.

Adansonia digitata (Baobab), relation of *Dysdercus* spp. to, in Africa, **222**, **490**, **573**, **574**.

Adelges (see Chermes).

Adelphocoris lineolatus, bionomics and control of, on lucerne in Russia, 475.

Adesmus borgmeieri, sp. n., on kapok in Brazil, **290.**

Adhesives, banding with, 28, 170, 172, 277, 354, 398, 499, 517, 611, 628, 693; spraying with, against *Tetranychus telarius*, 583; other methods of catching insects with, 125, 245, 249, 466, 633, 684.

Adia genitalis (see Hylemyia).

Adina cordifolia, Xylocopa in timber of, in India, 337.

Adonia variegata, in wheat fields in Central Asia, 453.

adonidis, Entomoscelis. adonidum, Pseudococcus.

Adoretus versutus, on beans in Fiji,

Adrama apicalis, on tea in Formosa, 339.

adreptella, Carposina.

adspersus, Exenterus.

adustus, Spilocryptus; Xylion. advena, Ahasverus (Cathartus).

Aegeria exitiosa, measures against, on peach in U.S.A., 255, 526; bionomics of, 526.

Aegeria pictipes, measures against, on peach in Illinois, 527.

Aegeria typhiaeformis (tiphiaeformis), parasites of, on apple, etc., in Italy, 4, 629, 630; bionomics of, 630.

aegopodii, Cavariella.

aegyptiaca, Icerya.

aegyptiacus, Achrysopophagus.

Aelia, type of injury to cereals by, in Germany, 352; measures against, 353.

Aelia acuminata, bionomics of, on cereals in Germany, 63, 64, 632.

Aelia klugi, ecology of, on cereals in Germany, 631.

Aelia rostrata, ecology of, on cereals in Germany, 631.

Aelia sibirica, on wheat in Central Asia, 453.

aeneus, Chalcodermus; Corymbites (Selatosomus); Meligethes.

Aeolarchis sphenotoma, parasites and control of, on Pandanus in Fiji, 213.

Aeolesthes induta, measures against, on tea in Formosa, 340.

aeolopa, Lobesia.

Aeolothrips fasciatus, predacious on insects in wheat fields in Central Asia, 453.

aequalis, Euaresta; Heterobostrychus.

aequatus, Rhynchites.

aeripennis, Corymbites (Ludius).

Aeroplanes, for distributing insecticidal dusts, 299, 514, 610, 645; power dusters compared with, 299; autogiro (q.v.) compared with, for spraying against forest pests, 645; Psallus seriatus trapped in, at high altitudes, 245; insects intercepted in quarantine in, in Hawaii, 368.

Aeropus sibiricus, in Russian Union, 146, 147, 320, 680; causing white ear in wheat, 680; relation of outbreaks of, to humidity, 146, 147.

aeruginosus, Dermatodes.

Aeschynomene, relation of Nomadacris septemfasciata to, in Tanganyika, 388.

Aesiotes leucurus, bionomics and control of, on conifers in New South Wales, 602.

Aesiotes notabilis, bionomics of, on Araucaria cunninghami in Queensland, 603.

aestivum, Apion. aestuans, Xylocopa.

Aethemenes chloris, bionomics of, on rice in Sierra Leone, 271.

affinis, Agria; Dibrachys; Pempheres; Phytomyza.

Afghanistan, Cydia pomonella in, 575.

Africa, French West, locusts in, 11-14, 280, 477, 478, 480, 481, 482, 487, 570, 571, 572; pests of oil palms in, 347-349; beneficial insects in, 347, 349.

Africa, Italian East, new Encyrtid parasites of Coccids in, 40 503;

INDEX. 711

locusts in, 478, 570; termites of, 174; absence of Stephanoderes hampei from, 175.

Africa, Portuguese East, locusts in,

62, 479, 484.

Africa, South, Citrus pests in, 330; cotton pests in, 223, 487, 488, **489, 490, 491**; pests and diseases of tobacco in, 360; Coleoptera in timber in, 60; locusts in, 91, 92, 93, 393; 'termites in, 180, 384; beneficial insects in, 40, 323, 394, 489, 503, 506; ants hindering biological control of certain pests in, 323; new Microsporidian parasites of Cactoblastis spp. in, 459; beneficial insects introduced into Mauritius from, 611.

Africa, South-West, Locustana par-

dalina in, 394.

africana, Gryllotalpa.

africanus, Ceroplastes; Rhaconotus menippus.

agamemnon, Oryctes.

Agar, in preparations for feeding parasites, 426; as an activator for nicotine, 645.

agricola, on vines Agarista Queensland, 365.

agaristae, Euplectrus.

Agelastica alni, action of HCN on,

Ageniaspis, possibly parasitic on Leucoptera in Tanganyika, 127. agilis, Phorocera.

Aglia tau, Pentatomid predacious on, in Germany, 354.

Agonoderus lecontei (pallipes, auct.), identity of, 595.

pallipes (limbatum), Agonum synonymy of, 595.

agonus, Limonius Pheletes (see ectypus). Agria affinis, parasite of spruce

sawflies in Czechoslovakia, 518. agricola, Agarista.

Agricultural Institutions, value of entomological collections in, 568.

Agrilus hyperici, introduced into against Hypericum Australia perforatum, 546.

Agrilus ruficollis, susceptibility of blackberry varieties Oklahoma, **79.**

Agriotes, feeding habits of larvae of, in Germany, 398; measures against, on lupins in Russia, 683. Agriotes lineatus, possibly

Britain, 96.

Agriotes obscurus, in Britain, 96; in Siberia, 679; bionomics of and (1875)

soil conditions in relation to, 96,

Agriotes sericeus, on tobacco in Japan, 210.

Agriotes sputator, distribution of, in Russia, **146**.

Agromyza, bionomics of, on barley in Japan, **209.**

Agromyza lantanae, attacking Lantana in India, 612.

Agromyza phaseoli, measures against, on beans in New South Wales, 58. agropyri, Phenacoccus.

Agropyrum, pests of, in Russia,

322, 680.

repens, Contarinia Agropyrum tritici on, in Italy and Holland, 3, 268; pests of, in Siberia, 312,

Agropyrum smithi, liable to favour Cephus cinctus, 368.

graminis Agrostis, Dasyneura probably on, in Britain, 635; Pediculopsis graminum on, in Russian Union, 151.

Agrostis vulgaris, Dasyneura graminis on, in New York, 635. Agrotis, on beet in Britain, 674.

Agrotis malefida (see Feltia).

Agrotis pronuba, suitability of, as host for Trichogramma evanescens,

Agrotis segetum, bait for, on tobacco in S. Rhodesia, 125; on potato in Czechoslovakia, 469; in Russian Union, 42, 146, 306, 307, 455, 589, 627, 677, 680, 682; bionomics and physiology of, 42, 146, 455, 677, 682; race of Trichogramma evanescens parasitic on, 589; beet protecting Scorzonera against, 680; experiments with fungi against, 306, 307; tests of insecticides on, 309, 310, 627.

Agrotis vetusta, measures against, on lucerne and maize in Wisconsin, 32; fungus infesting, 32.

Agrotis ypsilon, on tobacco in Fiji, 213; on cabbage and sugar-cane in U.S.A., 250, 657; seasonal abundance of, 250; parasites of,

Ahasverus advena, measures against, in oil-palm kernels in Dahomey, 349.

aino, Diachasma (Opius).

Air-tight Storage, against cereal pests, **667**.

Aira flexuosa, Dasyneura airae on, in France, 635.

airae, Dasyneura.

Alabama, miscellaneous pests in, 82, 379, 599; Naupactus spp. and legislation against them in, 259, 275, 378; beneficial insects in, 82.

Alabama argillacea (on cotton), in U.S.A., 243, 246, 258, 260; in W. Indies, 271, 661; flight of, physiological factors 260: affecting abundance of, 258; measures against, 271, 661; tests of insecticides on, 243, 246.

alaskensis, Pikonema.

Albania, summary of plant quarantines in, 592.

Albasol, as an emulsifier, 141.

Alberta, cereal pests in, 176, 539; Psyllids of, 175.

alberti, Eberhardia.

albifrons, Decticus; Spilochalcis. Technomyrmex • (see albipes,

detorquens).

albisignata, Idatiella. albitarsis, Anastatus.

Albizzia, Tineid on, in Formosa, 447.

alboannulatus, Dirhicnus (Pteromalus).

albohirtum, Lepidoderma (Lepidiota). albolineatus, Dendrolimus.

albomarginatus, Chorthippus.

albobunctatus, Crossotus.

Albumen, in baits for ants, 87.

Alcohol, for extracting insecticidal principles of plants, 135, 450. (See Methylated Spirit.)

Alder, pests of, in Britain, 58, 563; pometaria Alsophila Canada, 663; Linoclostis gonatias on, in Formosa, 447; Cicadella viridis on, in Germany, 398; locust damaging, in Japan, 618; effect of feeding Lymantria monacha on, 399.

aldrichi, Palpostoma; Sarcophaga. alecto, Telenomus (Prophanurus).

Aleurites, Buzura suppressaria on, in China, 614; Coccid on, in Nyasaland, 126.

Aleurites montana, Achaea janata on, in Burma, 693; as a shadetree for coffee against Xylotrechus quadripes in Indo-China, 49; pests of, in Java, 523.

Aleurodes brassicae (on cabbage), bionomics of, in Britain, 61, 265: in Italy, 265.

Aleurodiosis, 637.

Aleurolobus barodensis, new parasite of, on sugar-cane in India, 385.

Algeria, Cydia pomonella in, 191-193; Phloeosinus spp. on conifers in, 226; locusts in, 480; restrictions on importation of vines against Tunisia from, into Phylloxera, 11.

aliberti, Eloeidiphilos. alienus, Lasius niger.

Alissonotum, on sugar-cane Formosa, 446.

Alkylphenylbenzenesulphonic Acid, addition of, to derris spray, 249.

Allium, Aphid on ornamental, in Washington, 541. (See Onion and Leek.)

allyni, Eupelmus.

Almond, pests of, in Australia, 21, 57; pests of, in California, 650; Mimastra cyanura on, in India, 503; Monosteira unicostata on, in Morocco, 359; Coccids on, in Uzbekistan, 474; nutritive value of kernels of, for Ephestia kuehniella, 64.

alni, Agelastica.

alopecuri, Dasyneura.

Alopecurus, Phytomyza nigra on, in Japan, 208.

Alopecurus pratensis, thrips on, in Finland, 160; Dasyneura alopecuri on, 635.

Alophoropsis brasiliensis (see Hyalomyia).

alpestris, Dasyneura.

Alphus canescens, on Anonaceae in Brazil, **566.**

Alpine Currant (see Ribes alpinum). Alsike Clover (see Trifolium hybridum).

Alsophila pometaria (on elm, etc.), bionomics of, in Canada, 662-664; in U.S.A., 28, 74; measures against, 74, 663.

alternatus, Nabis.

Altingia excelsa, pests of, in Java,

Aluminium Arsenate, 283; spraying with, against Leptinotarsa decemlineata, 583, 584.

Aluminium Hydroxide, as a corrective for calcium arsenate, 373.

Aluminium Nitrate, affecting wetting power of sprays, 412. Aluminium Silicate, as a

a carrier, 431.

Aluminium Sulphate, as an adhesive for sprays, 256, 257, 532, 533.

Alurnus corallina (see Mecistomela). Alyssum, Dociostaurus maroccanus associated with, in Transcaucasia 321.

INDEX. 713

amabilis, Eublemma: Prepodes similis.

Amarantus, Hymenia recurvalis on, in New South Wales and Fiji, 639, 641.

Amazon Fly (see Metagonistylum

minense). ambigua, Myrmelachista.

ambiguella, Clysia.

Amblynotus longitarsis, parasite of Leucopis obscura in Britain, 263.

Amblyteles camelinus, parasite of Vanessa cardui in Transcaucasia,

Amercedoides nitida (see Revena). America, cactus-feeding Phycitinae of. 511.

America, North, Tarsoneminae of, 224.

americana, Larra; Malacosoma; Periplaneta.

americanum, Mezium.

americanus, Polistes crinitus.

Amerrhinus ynca, on Cocos coronata in Brazil, 661.

amicroploides, Macrocentrus.

Amitermes, in Australia, 545.

Ammonia, in bait for Dacus psidii, 131; and oleic acid, in nicotine spray, 514.

Ammonium Bifluoride, in baits for grasshoppers, 134.

Ammonium Fluoride, in baits for grasshoppers, 134.

Ammonium Nitrate, in baits for

Dacus oleae, 98. Ammonium Polysulphide, against mites, 461.

Ammonium Sulphate, in baits for Trypetids, 98, 270.

amoenus, Anagyrus.

Amoora wallichii, Xylocopa latipes in timber of, in India, 337.

Amorpha fruticosa, rotenone in, 502. Amorphoidea lata, measures against, on cotton in Philippines, 184.

Amorphophallus Ptinus koniac, japonicus in flour of, in Japan, 445.

spp., parasitised by Ampeloeca Apanteles congregatus in U.S.A.,

ampelophaga, Theresimima (Procris, Theresia).

Amphimallus solstitialis, bionomics of, damaging lawns in Holland, 269; beet protecting Scorzonera against, in Russian Union, 680. Amphipyra tragopoginis, on poplar

and willow in Morocco, 196. Amphorophora avenae (see Macro-

siphum).

Amphorophora cosmopolitana (see A. lactucae).

Amphorophora dirhodum (see Macrosiphum).

Amphorophora lactucae, on currant: and gooseberry in Britain, 333.

Amphorophora rubi, on raspberry in Britain, 333; variety of raspberry resistant to, in Washington, 280.

amputator, Oncideres.

amyntas, Acolastus; Pleurotropis. Amyosoma chilonis (see Microbracon chinensis).

Anabasine, as an insecticide, 593. Anabasine Sulphate, spraying with, against Aphids, 151, 216; and soap, 151.

Anabrus simplex, outbreaks of, in U.S.A., 29, 599.

Anacardium occidentale, piercing moths damaging, Sierra Leone, 271.

Anacentrinus subnudus, bionomics of, on sugar-cane, etc., Louisiana, 248.

Anacridium moestum melanorhodon, reactions of, to light, 603.

Anagrus armatus, suggested introduction of, into Florida against Saccharosydne saccharivora, 657.

Anagrus atomus, parasite of Erythroneura pallidifrons in Britain,

Anagyrus, African species of, 503; parasite of mealybugs in Natal,

Anagyrus abyssinicus, sp. n., parasite of *Pseudococcus* in Eritrea, 503.

Anagyrus amoenus, sp. n., parasite of Pseudococcus in Transvaal, 503.

Anagyrus bugandaensis, sp. parasite of Pseudococcus kenyae in Uganda, 503.

Anagyrus coccidivorus, introduced into Porto Rico from Hawaii against Pseudococcus brevipes, 502.

Anagyrus kivuensis, sp. n., parasite of Pseudococcus in Belgian Congo, 503.

Anagyrus niger, sp. n., parasite of Pseudococcus in Uganda, 503.

Anagyrus nigrescens, sp. n., parasite of Pseudococcus in Transvaal,

Anagyrus pullus, sp. n., parasite of Pseudococcus in Eritrea, 503.

Anagyrus subproximus, characters of group of, 504.

Anaitis spp., ants preventing establishment of, against Hypericum perforatum in Australia, 282.

Analeurops cuthbertsoni, on tobacco

in S. Rhodesia, 125.

analis, Bruchus (Callosobruchus).
Anamastax, identity of type of, 294.
(See Tritaxys.)

Anaphes anomocerus, parasite of Psallus seriatus in U.S.A., 381,

538.

Anaphothrips obscurus, bionomics of, on wheat in Sweden, 121.

Anarsia lineatella, on plum in Cyprus, 270.

Anastatus spp., parasites of Dendrolimus spectabilis in Japan, 211.

Anastrepha, in Brazil, 34; Dirhinus giffardi introduced into Porto Rico against, 502; in Trinidad, 24.

Anastrepha acidusa, bait-spray for, 256.

Anastrepha fraterculus, parasite of, in Brazil, 34.

Anastrepha fraterculus var. mombinpraeoptans, parasites of, in Porto Rico, **502**.

Anastrepha suspensa, parasites of, in Porto Rico, 502; bait-spray for, 256.

anastrephae, Opius.

Anatis quindecimpunctata, predacious on Anuraphis roseus in Connecticut, 428.

anceps, Hudsona.

Ancylis comptana (on strawberry), parasites of, in U.S.A., 374, 376, 377; measures against, 376.

ancylivorus, Macrocentrus. Andres-Maire Traps, for Spodoptera

mauritia, 577.

Andropogon, relation of locusts to, in Fr. W. Africa, 571.

Aneristus ceroplastae, parasite of Pulvinaria polygonata in Formosa, 558.

Anethum graveolens, new Aphid on, in California, 590; celery virus transmitted to, 590.

Anetia nigripes (see Lydella).

Angitia, parasite of Cydia nigricana in Britain, 38, 39; parasite of C. molesta in Victoria, 640.

Angitia cerophaga (parasite of Plutella maculipennis), in Britain, 262; introduced into New Zealand, 262.

Angitia fenestralis (parasite of Plutella maculipennis), in Britain, 262; introduced into New

Zealand and New South Wales, 262, 545; reared from *Pissodes* notatus in Finland, 8.

Angitia molestae (parasite of Cydia molesta), in U.S.A., 377, 544; introduced into Australia, 544.

Angitia rufipes, 38. angusticeps, Thrips.

angusticollis, Psylliodes; Zootermopsis.

angustifrons, Cheiloneurus.

Aniline, in oil emulsions against vine moth pupae, 103.

Anisodactylus binotatus, measures against, on strawberry in Germany, 457.

Anisoplia austriaca, on wheat in Czechoslovakia, 468; fungus infesting, in Russia, 306; experiments with fungi against, 307.

anitus, Microphanurus. annexa. Feltia.

annulata, Campsomeris.

annulipes, Caliroa (Eriocampoides); Chelonus.

anobiidivorus, Heterospilus.

Anobiids, frass and débris due to boring by, 568.

Anobium punctatum (striatum), in building timber in U.S.A. and Russia, 78, 322; suitable for testing wood preservatives, 204.

Anogeissus acuminata, Heterobostrychus aequalis in timber of, in Burma, 693.

Anomala, Lacon murinus experimentally feeding on, 398.

Anomala cuprea, bionomics of, in Japan, 17, 338; morphology of, 17.

Anomala dubia var. frischi, damaging lawns, etc., in Holland, 269.

Anomala expansa, bionomics of, on sugar-cane, etc., in Formosa, 363, 446.

Anomala horticola, bionomics of, damaging lawns, etc., in Holland, 269.

Anomala orientalis, on sugar-cane and pineapple in Hawaii, 366, 367; parasites of, 367.

Anomala rufocuprea, on tobacco and soybeans in Japan, 210, 338; effect of light on, 338.

Anomala viridis, on Altingia excelsa in Java, 227.

Anomis flava (see Cosmophila).

Anomis luridula (texana), measures against, on cotton in Peru, 164. anomocerus, Anaphes.

Anona, Coccus longulus on, in Egypt, **509**.

Anona squamosa, pests of, in Brazil, 566, 567.

anonella, Cerconota.

anonicola, Codiohispa; Meroprion. Anophia leucomelas (on sweet potato), in Japan, 444, 445, 620; in Loochoo Islands, 444; bionomics of, 444, 445; measures against, 445.

antancarus, Enicospilus. antennata, Graptolitha.

antennatum, Callidium.

Antestia, measures against, coffee in Kenya, 667; vector of Nematospora, 389.

Antestia lineaticollis, on coffee in

Tanganyika, 127.

Antheraea pernyi (Chinese Oak Silkworm), experiments on possibility of establishment of, in Russian Union, 403, 586.

Antholcus varinervis, introduced New Zealand into against Acaena, 449.

Anthomyia, species of Hylemyia referred to, 113.

Anthomyiids, nomenclature injurious, 113, 114.

Anthonomus grandis (Cotton Boll Weevil), legislation against introduction of, into Barbados, 661; in Haiti, 661; in U.S.A., 244, 246, 247, 380, 381; susceptibility of varieties of cotton to, **247**, **381**; experiments with insecticides against, 244, 245, 246, 247.

Anthonomus piri, bionomics of, in Germany, 632.

pomorum Anthonomus (Apple Blossom Weevil), in Britain, 333; Czechoslovakia, 469: Germany, 298; on pear Manchuria and China, 340, 450; hibernating in pine bark, 298; measures against, 298, 450.

Anthonomus pyri (see A. piri).

Anthonomus rectivostris, measures against, on cherries in Germany, 156.

Anthonomus rubi, measures against, on strawberry in Germany, 155.

Anthonomus signatus, measures against, on strawberry in U.S.A., 380.

Anthonomus spilotus, 632.

Anthonomus varians, bionomics of, on pine in Rumania, 355.

Anthonomus vestitus, bionomics of. on cotton and Hibiscus in Peru. 164.

Anthores leuconotus, on coffee in Nyasaland, 126.

Anthracene Oil Emulsion, (See Tar Distillates.)

Anthrenus, characters and habits of species of, in Germany, 47.

Anthrenus fuscus, 47.

Anthrenus museorum, 47.

Anthrenus pimpinellae, 47.

Anthrenus scrophulariae, bionomics of, damaging woollen goods in Germany, 47.

Anthrenus verbasci (varians). bionomics of, in Germany and Victoria, 7, 47, 187, measures against, 188.

Anthrenus vorax, infesting house insulation in U.S.A., 640.

Anthriscus cerefolium, celery virus transmitted to, 590.

Anthrocephalus pomonellae (see Stomatoceras).

Anticarsia gemmatalis, measures against, on soy beans in U.S.A., 258, 371, 373, 422; bionomics of,

Antigua, attempted introduction of Coccinellid against Aspidiotus destructor on coconut in, 25: Diatraea saccharalis and its biological control on sugar-cane in, 25, 183, 658, 659, 660; Lixophaga into diatraeae introduced Lucia from, 182.

Antimony Potassium Tartrate (see Tartar Emetic).

antipodialis, Chromocryptus.

antiqua, Hylemyia (Delia); Orgyia. antonii, Helopeltis.

Ants, list of Formosan species of, 446; associated with Aphids, 44, 276, 323, 363, 376; associated with Coccids, 323, 335, 363, 611, 665; other Arthropods in nests of, 288; destroying other insects, 82, 100, 231, 239, 282, 323, 335, 363, 364, 489, 564, 611, 642; natural enemies of, 129, 288; experimentally infested Beauveria bassiana, 688; measures against, 25, 44, 87, 287, 288, 323, 435, 541, 567, 611, 665; differences in sensitivity of species of, to arsenical baits, 112.

Ants, Carpenter (see Camponotus). Ants, Leaf-cutting (see Acromyrmex and Atta).

Ants, Red Forest (see Formica

Anuraphis padi (helichrysi), sprays against, on plum in Britain, 332, 333.

Anuraphis persicae-niger, measures against, on peach in Tasmania,

Anuraphis roseus (on apple), in Britain, 333; in U.S.A., 428, 535, 536; bionomics and biological control of, 428; other measures against, 333, 428, 535, 536: characters distinguishing Aphis pomi and, 428.

Anuraphis schwartzi, on peach in

Germany, 687.

Anuraphis tulipae, not transmitting breaking disease of tulips in

Oregon, 369.

Aonidiella aurantii (on Citrus), in Argentina, 23; in Australia, 292; in California, 342, 343, 460, 649; natural enemies of, in Cyprus, 270; in Egypt, 509; in India, 503; in Japan, 207; in New Zealand, 523, 524; in S. Rhodesia, 14; bionomics of, 207, 524; measures against, 15, 23, 292, 460, 509, 524; toxicity of dinitroorthocyclohexylphenol to, 649; resistance of strains of, to HCN and other fumigants, 342, 343.

Aonidiella citrina, bionomics of, on Citrus in Transcaucasia, 148, 149.

Aonidiella orientalis. measures against, on lac host-trees in India,

Aonidiella perniciosa (see Aspidiotus).

aonidum, auct., Chrysomphalus (see C. ficus).

Apamea secalis (see Trachea). Apamea testacea (see Luperina).

Apanteles, parasite of Phthorimaea operculella in Brazil, 289; parasite of Pissodes validirostris in Finland, 8; parasite of Proto-parce sexta in Florida, 251; probably parasitic on Malacosoma neustria in Holland, 205; parasite of Pyrausta phoenicealis in Japan, 210; parasite of P. aurata meridionalis in Morocco, 193; emendations to key to, 505.

Apanteles bedelliae, parasite of Bedellia somulentella in Kansas,

Apanteles carposinae, sp. n., parasite of Carposina adreptella in New Zealand, 37.

Apanteles coleophorae, sp. n., parasite of Coleophora fuscedinella in Britain, 58.

Apanteles compressiventris, hosts and distribution of, 505.

Apanteles congregatus (parasite of Sphingids), in U.S.A., 178, 253, 655; bionomics and distribution of, 253, 254.

Apanteles diatraeae, introduced into U.S.A. against Diatraea sacchara-

lis, 656.

Apanteles difficilis, parasite Erannis in Crete, 172.

Apanteles dignus, sp. n., parasite of Phthorimaea spp. in California and Virgin Islands, 133.

Apanteles dioryctriae, sp. n., parasite of Dioryctria peverimhoffi in

Morocco, 58.

Apanteles flavipes, parasite Diatraea sticticraspis in Mysore,

Apanteles insignis, sp. n., parasite of Platyptilia acanthodactyla in California, 133.

Apanteles langei, sp. n., parasite of Pyrausta insequalis in California,

Apanteles liparidis, bionomics of, in Japan, 91, 209, 211, 442.

Apanteles machaeralis, parasite of Hapalia machaeralis in Burma, 693.

Apanteles plutellae, hosts and distribution of. 505.

Apanteles scutellaris, establishment of, against Phthorimaea operculella in Hawaii, 367.

Apanteles sicarius, parasite Plutella maculipennis in Britain,

Apanteles solitarius, establishment of, against Stilpnotia salicis in U.S.A., 499.

Abanteles spurius, parasite Vanessa cardui in Transcaucasia, 587.

Apanteles tachardiae, unsuitable for use against Holcocera pulverea in India, 129.

Apate monacha, on coconut in Jamaica, 239.

apatelae, Arthrolytus (see Dibrachys cavus).

Aphaereta, key to European species of, 604.

Aphaereta difficilis, sp. n., parasite of Hylemyia spp. in Morocco, 604.

Aphalara calthae, on potato in Germany, 636.

Aphanistes ruficornis, parasite of Sphinx pinastri in Poland, 677. Aphelinus diaspidis (see Aphytis).

Aphelinus mali (parasite of Eriosoma lanigerum), bionomics of, 10, 549, 589; utilisation of, 20, 117, 127, 162, 271, 548-550, 588; factors affecting efficiency of, in New Zealand, 548-550.

Aphelinus mytilaspidis (see Aphytis).
Aphidencyrtus aphidivorus, parasite of Leucopis obscura in

Britain, 263.

aphidis, Pachyneuron.

Aphidius, Coelonotus considered a synonym of, 615.

Aphidius japonicus (see Coelonotus). Aphidius rosae, parasite of Macrosiphum onobrychis in Utah, 31.

Aphidius salignae, sp. n., parasite of Lachnus salignus in Japan, 615.

aphidivorus, Aphidencyrtus.

Aphids, and plant diseases, 83, 84, 87, 125, 126, 157, 158, 213, 226, 236, 267, 274, 280, 289, 301, 302, 303, 333, 365, 369, 370, 448, 472, 226, 228, 262, 263, 271, 305, 334, 371, 382, 428, 578, 588, 615, 628, 629, 674, 687; technique for rearing Hippodamia on, 81; factors affecting wing development in, 323; trapped by leaves of tomato and bean, 113, 142; tests of insecticides on, 236, 259, 272, 332, 458; of California, 261; of Canada, 104; of Central Europe, 392; of Italy, 224, 512; of Lapland, 512; of Sweden, 160; classification and new species of, 100, 104, 160, 224, 261, 336, 339, 392, 552, 590, 596; check-list of genera of, 392.

Aphis, transmitting disease of Passiflora in Australia, 448; key to species of, on Ribes, 597. Aphis apigraveolens, sp. n., transmitting disease of celery in

California, 590.

Aphis apii, transmitting disease of celery in California, 590.

Aphis avenae (see Macrosiphum). Aphis brassicae (see Brevicoryne).

Aphis chloris, bionomics of, on Hypericum perforatum in Britain, 59.

Aphis coprosmae, in New Zealand, 336.

*Aphis fabae, in Britain, 673, 674; in Germany, 334, 686, 688; bionomics of, 674, 686; measures against, on beans and beet, 334, 674; relation of, to virus diseases of plants, 686, 688; considered distinct from A. rumicis, 674.

Aphis ferruginea-striata, sp. n., vector of food-plants and relation to celery virus of, in California, 590, 591.

Aphis forbesi (see Cerosipha).

†Aphis frangulae, bionomics and relation to virus diseases of plants of, in Germany, 633, 634, 686, 688; on cotton in Russia, 215, 216.

†Aphis gossypii, in Czechoslovakia, 471; in India, 228; in Manchuria, 361; in Peru, 164; in U.S.A., 243, 245, 256, 590; vector of virus disease of celery, 590; on Citrus, 256; on cotton, 164, 228, 243, 245; hairiness of cotton leaves in relation to, 243; Leucopis griseola predacious on, 228; dust of derris with chloropicrin ineffective against, 471; effect of calcium arsenate dusts on infestation by, 164, 243.

Aphis grossulariae, on currant and gooseberry in Britain, 333, 596. Aphis heraclella, on celery in

California, 590.

Aphis laburni (leguminosae), transmitting rosette disease of groundnuts in Nyasaland, 126; bionomics of, on cotton, etc., in Russia, 215, 216; measures against, 216.

Aphis maidis, in Fiji, 641; in Florida, 658; in Italy, 472; in Philippines, 667; in Queensland, 87; on maize, 88, 641; transmitting sugar-cane mosaic, 87, 658, 667; possible vector of virus disease of sugar sorghum, 472; measures against, 88.

*Aphis medicaginis, on certain lupins in Russia, 151; measures

against, 151.

Aphis middletonii, vector of disease of celery in California, 590.

Aphis neomexicanus, on Ribes in U.S.A., 596.

* In view of the divergent opinions on the identity of the Aphids of this group, no attempt is here made to show the synonymy.

† These references refer to the same species: some authors consider gossypii

a synonym of frangulae.

*Aphis papaveris, on beet in Czechoslovakia, 468.

Aphis pawneepae, ant associated with, on Cercis canadensis in

Kansas, 376.

Aphis pomi (on apple), in Britain, 332, 333; resistance of, to abnormal cold in France, 622; in U.S.A., 428, 534; measures against, 333, 534; studies on ovicides against, 236, 332; characters distinguishing Anuraphis roseus and, 428.

Aphis rhamni, in Britain, 563; in Germany, 157, 633, 686, 688; in Holland, 563, 578; in U.S.A., 563; relation of, to virus diseases of plants, 157, 563, 578, 633, 686, 688; Rhamnus primary food-

plant of, **578**, 686.

Aphis ribiensis, on Ribes in Utah, 596.

Aphis ribi-gillettei, sp. n., on currant and gooseberry in Utah, 596.

Aphis ribis, on Ribes gracile in

Missouri, 597.

*Aphis rumicis, 448; in Britain, 673; relation of, to virus diseases of plants in Germany and U.S.A., 267, 370, 590, 652; test of greenhouse fumigant on, 272; identity of, 673.

Aphis sacchari (on sugar-cane), in Jamaica, 238; in Queensland, 186; relation of, to diseases of

sugar-cane, 186.

Aphis sanborni, on gooseberry in U.S.A., 596.

Aphis spiraecola, on Citrus in U.S.A., 256, 259, 460, 461; insecticides against, 259, 460, 461.

Aphis tavaresi (on Citrus), measures against, in New Zealand, 277; in S. Rhodesia, 15.

Aphis varians, on Ribes in U.S.A., 596.

Aphis, Barley (see Brachycolus noxius).

Aphis, Black Bean (see Aphis

fabae).

Aphis, Citrus (see Aphis spiraecola). Aphis, Cotton (see Aphis gossypii). Aphis, Pea (see Macrosiphum

onobrychis).

Aphis, Peach (see Myzus persicae). Aphis, Potato (see Macrosiphum solanifolii).

Aphis, Rosy Apple (see Anuraphis roseus).

Aphis, Strawberry Root (see Cero-sipha forbesi).

Aphis, Woolly Apple (see Eriosoma lanigerum).

Aphis, Yellow Sugar-cane (see Sipha flava).

Aphomia gularis (in stored products), in China, 451; in Germany, 355, 520, 521; imported into Germany from Manchuria, 520; in Japan, 360; bionomics of, 520, 521; measures against, 451, 521.

aphyopterus, Microcryptus brachy-

pterus.

Aphytis chrysomphali (parasite of Coccids), in Argentina, 24; in Cyprus, 270.

Aphytis diaspidis, parasite of

Coccids in Brazil, 87.

Aphytis mytilaspidis (parasite of Coccids), in Argentina, 24; in Transcaucasia, 149.

Aphytis proclia, parasite of Aspidiotus perniciosus in Argentina,

24.

apicalis, Adrama. apicipennis, Exora. apigraveolens, Aphis.

apii, Aphis.

Apiomerus crassipes, predacious on *Horistonotus uhleri* in S. Carolina, **500.**

Apiomyia bergenstammi, in Corfu, 5; bionomics and control of, on pear in Italy, 5.

Apion, studies of migrations of, in Russia, 681.

Apion aestivum (on clover), bionomics of, in Russia, 682; distribution of, in Sweden, 117; susceptibility of varieties of clover to, 118; fungi infesting, and their utilisation, 118.

Apion apricans (on clover), in Czechoslovakia, 470; bionomics of, in Russia, 568, 681, 682; distribution of, in Sweden, 117; susceptibility of varieties of clover to, 118; fungi infesting, and their utilisation, 118; net for collecting adults of, 470.

Apion carduorum, on artichokes in France, 357.

Apion carduorum var. galactitis, bionomics and control of, on artichokes in France, 357.

Apion flavipes, bionomics of, in Russia, 568, 682.

^{*} In view of the divergent opinions on the identity of the Aphids of this group, no attempt is here made to show the synonymy.

Apion seniculum, bionomics and control of, on Trifolium hybridum in Latvia, 47.

Apion ugandum, on Voandzeia in

Nyasaland, 126.

Aplastomorpha calandrae, parasite of Pempheres affinis in India, 506. Apleurotropis lalori, sp. n., parasite of Promecotheca in New Guinea

Territory, 294.

Aporia crataegi (in orchards), in Czechoslovakia, 145, 468; proposed measures against, in Poland, 163; parasite of, 145.

Appelia (see Anuraphis).

Apple, Cydia pomonella on, in Afghanistan, 575; C. pomonella on, in Algeria, 192; pests of, in Argentina, 24, 188; pests of, in Australia, 20, 57, 89, 130, 159, 186, 279, 292, 345, 346, 390; pests of, in Baluchistan, 50-52, 577; pests of, in Britain, 236, 333, 396, 676; spray programme for, in Britain, 396; pests of, in Canada, 66, 382, 542, 662, 663, 664; Illiberis pruni on, in China, 450, 663; Eriosoma lanigerum on, in Cyprus, 271; pests of, in Czechoslovakia, 468, 469; pests of, in Denmark, 117; Melolontha hippocastani on, in Finland, 267; pests of, in France, 168, 390, 357, 622, 690; pests of, in Germany, 114, 155, 156, 298, 301, 398, 513; pests of, in Holland, 301, 578; Aspidiotus perniciosus on, in Hungary, 690; pests of, in India, 503, 577; pests of, in Italy, 4, 104, 197, 629, 630; pests of, in Japan, 207, 209, 361, **556, 557, 616;** pests of, in Jugoslavia, **10, 690;** pests of, in Manchuria, 340; pests of, in Morocco, 283, 359; pests of, in New Zealand, 159, 548, 550; E. lanigerum on, in Palestine, 127; pests of, in Poland, 162, 163, 675, 694; E. lanigerum on, in Palestine, 163, 675, 694; E. lanigerum on, in Palestine, 185; and permissions. S. Rhodesia, 125; A. perniciosus on, in Rumania, 690; pests of, in Russian Union, 303, 305, 316, 455, 474, 552, 587, 588, 589, 610, 623, 681, 683; pests of, in Switzerland, 232, 438; pests of, in U.S.A., 31, 33, 52, 77, 133, 134, 141, 166, 178, 272, 273, 343, 382, 384, 419, 420, 428, 429, 430, 432, 525, 532, 533, 534, 535, 536, 542, 642, 643, 648, 652; list of Cecidomyiids attacking, 557; susceptibility of varieties of, to

E. lanigerum, 100, 141; effect of feeding Lymantria spp. on, 405, 399, 406; insecticides causing injury to, 6, 134, 309, 420, 425, 430, 534, 535, 536, 680.

Apple (Fruit), restrictions on importation of, into Britain, 356; Rhagoletis pomonella intercepted in Russian Union, 439; effect of methyl bromide against pests in, 541, 542; problems of spray residues on, 133, 139, 168, 188-190, 192, 193, 273, 525, 539, 642, 648.

Apple Aphis, Rosy (see Anuraphis

roseus).

Apple Aphis, Woolly (see Eriosoma lanigerum).

Apple Blossom Midge (see Contarinia mali).

Apple Blossom Weevil (see Anthono-

mus pomorum). Apple Curculio (see Tachypterellus quadrigibbus).

Apple Leaf Jassid (see Typhlocyba

froggatti). (see Hyponomeuta Apple Moth

padellus malinellus). Light-brown Apple Moth,

Tortrix postvittana). Apple Worm, Lesser (see Enarmonia

prunivora).

Apple-tree Borer, Flat-headed (see Chrysobothris femorata).

apricans, Apion.

apricarius, Chorthippus (Stenobothrus).

Apricot, pests of, in Baluchistan, 51, 577; Myzus persicae on, in Britain, 236; Saissetia oleae on, in California, 649; M. persicae on, in Germany, 157, 687; samsonovi on, Eurytoma India, 503; Cydia pomonella on, Coccids on, in in Italy, 4; 474, 552; Uzbekistan, not attacked by Aegeria typhiaeformis, 630; penetration leaves of, by insecticides, 309. penetration of

Apricot (Fruit), Ceratitis capitata intercepted in, in Russian Union, 439; (dried), seldom attacked by Carpophilus hemipterus, 201.

Apriona cinerea, on apple in India, 503.

Aptinothrips rufus, on wheat and grasses in Sweden, 121.

aquifolii, Phytomyza (see P. ilicis). Arabia, locusts in, 480, 481, 482. Arabis, Dasyneura alpestris causing

leaf galls on, 673.

arabis, Dasyneura (see D. alpestris).

Arachis hypogaea (see Ground-nuts). Aradus cinnamomeus, trapped by leaves of tomato in Germany, 113. Araecerus fasciculatus, in stored cacao in Ceylon, 638.

Araucaria cunninghami, Aesiotes notabilis on, in Queensland, 603.

arboreus, Microcerotermes. archangelskyae, Leucodiaspis.

Arctia caja, on tobacco in Japan, 210.

Arctia caja phaeosoma, on cereals and peas in Japan, 556.

Arctia villica, parasitised by Apanteles compressiventris, 505.

Arctium lappa, Acridid damaging, in Japan, 618.

Arctonis chrysorrhoea, parasite of, in Japan, 338.

Arcyptera fusca, in Jugoslavia, 480. Arcyptera microptera (see Pararcyptera).

Areginal, fumigation of flour mills with, against Ephestia kuehniella, 400.

Aresket, as a wetter and spreader for sprays, 71, 72.

Argentina, Citrus pests in, 23, 238; orchard pests in, 24, 188, 237, 601, 602; miscellaneous pests in, 87, 238, 413, 623; Coccids in, 23, 24, 87, 238; locusts and grasshoppers in, 213, 214, 476, 543; beneficial insects and biological control in, 23, 24, 214, 237, 238, 601, 602; amendments to plant quarantines in, 344; insects introduced into other countries from, against Opuntia, 345, 459.

Ephialtes (Calliephiargentinus, altes).

argillacea, Alabama,

Argyresthia ephippella, measures against, on cherry in Austria, 232. Argyria sticticraspis (see Diatraea). Argyroploce, new parasite of, on Inocarpus in Fiji, 19.

Argyroploce illepida, parasites of, on beans in Fiji, 213, 641.

Argyroploce leucotreta, measures against, on orange in S. Africa,

Argyrotaenia citrana (see Eulia). aridula, Chaetocnema.

Ariri Palm (see Cocos vagans).

aristella, Lonchaea.

Aristotelia, on Iris in Japan, 441. Arizona, Rhynchota on cotton in,

381, 416, 538; miscellaneous pests in, 142, 377, 378; Paratrioza cockerelli in, 409, 422; parasites of Psallus seriatus in, 538.

Arkansas, Diceroprocta vitripennis on cotton in, 165; Listroderes obliquus in, 327; miscellaneous pests in, 178, 537, 599; parasite of Psallus seriatus in, 538.

Arma custos (see Auriga).

armatus, Anagrus; Phloeosinus.

armigera, Heliothis.

Armillaria mellea, 504, 562.

armoraciae, Phaedon; Phyllotreta. Armyworm, Nutgrass (see Laphygma exempta).

Armyworm, Southern (see Xylomyges eridania).

Arrhenodes, type and synonymy of,

Arrhenodes minutus (septentrionis), synonymy of, 595.

arrogans, Trigonophymus.

Arrowroot, Calpodes ethlius on, in

St. Vincent, 271.

Arsenic, in baits, 32, 67, 112, 134, 250, 291, 537; reactions of wireworms to, 28, 29; in dust against Laphygma exempta, 368; uses of, against termites, 181, 465; danger to bees and birds of insecticides containing, 4, 101, 156, 159; problem of residues of, on fruit, 139, 168, 191, 192, 193,

273, 293, 331, 432, 495, 539, 642. Arsenic Pentoxide, in bait for locusts, 602; use of, against Rhizopertha in pith helmets, 612.

arteagai, Opsophyto.

Artemisia, Pyrausta nubilalis on, in Britain, 116, 295; Geometrid on, in Sicily, 630.

Arthrochlamys spilota, bionomics of, on rhododendron in Japan, 212.

Arthrolytus apatelae (see Dibrachys cavus).

Artichoke (Cynara), Phytomyza atricornis on, in California, 600; weevils on, in France and Italy, 357, 473; liable to attack by Dectes spinosus, 546.

Artona catoxantha, on coconut in Philippines, 451.

Arundinaria, new Coccid on, in Formosa, 103.

arundinariae, Kuwanaspis. arundinis, Hyalopterus.

Asanilic Acid, toxicity of, to grasshoppers, 134.

Ascia rapae (see Pieris).

Ascogaster canifrons, parasite of Cydia pomonella in Central Asia, 588.

Ascogaster carpocapsae (parasite of Cydia molesta), introduced into Australia from U.S.A., 544; utilisation of, against C. nigricana in Canada, **39.**

Ascogaster quadridentata, parasite of Cydia nigricana in Britain, 38: parasite of C. pomonella in Italy, 4; not experimentally parasitising Enarmonia caryana in Transcaucasia, 377; bionomics of, 38.

Aserica formosae, soil characters in relation to, in sugar-cane fields in Formosa, 206.

Aserica japonica (on soy beans), effect of light on adults of, in

Japan, 338.

Ash (Fraxinus), cankerworms on, in N. America, 74, 663; Prociphilus oriens migrating to Abies

from, in Japan, 211.

Asia, Central, Ulobaris loricata on beet in, 152; cotton pests in, 147, 609, 625; pests of forest and shade trees in, 153, 474; Aphids on leguminous plants in, 552; orchard pests in, 315, 474, **552, 587, 610, 623;** pests of rubber-producing plants in, 44; wheat pests in, 453; Coccids in, 315, 336, 474, 552; beneficial insects in, 453, 588, 589, 609, 626.

asiatica, Chionaspis. asozanus, Gelis (Pezomachus).

asparagi, Crioceris.

Asparagus, Crioceris asparagi on, in Utah and Italy, 426, 471.

Asparagus Beetle (see Crioceris asparagi).

Aspens (see Populus tremula and P. tremuloides).

Aspergillus, Necrobia rufipes feeding on, 584.

asphaltinus, Otiorrhynchus.

Asphondylia, on wild serradella in Portugal, 495; food-plant range of species of, 673.

Aspidiotiphagus citrinus, parasite of Aonidiella citrina in Trans-

caucasia, 149.

Aspidiotiphagus lounsburyi, parasite of Aspidiotus hederae Nyasaland, 126.

Aspidiotus chinensis, bionomics of, on imported orchids in Japan,

Aspidiotus destructor (on coconut), in Fiji, 212; in Mauritius, 611; in W. Indies, 25, 502; Coccinellids and biological control of, 25, 212, 502, 611; ants fostering, 611.

Aspidiotus hederae, parasite of, on Aleurites in Nyasaland, 126.

Aspidiotus jourdani, sp. n., Morocco, 392.

Aspidiotus juglans-regiae, measures against, on walnut in California, 649.

Aspidiotus lataniae, measures against, on vines in Brazil, 34.

A spidiotus orientalis (see A onidiella). Aspidiotus ostreaeformis, on fruit trees and bush-fruits in Czechoslovakia, 468.

Aspidiotus perniciosus (on fruit trees), in Austria, 690; current in Germany, 467; Hungary, 690; legislative measures against, in Jugoslavia, 690; in Manchuria, 340; in Punjab, 503; in Rumania, 690; in Russian Union, 303, 315, 681; in U.S.A., 343, 534; bionomics of, 315, 316; Coccinellids predacious on, and their utilisation, 303, 304, 316; other measures against, 467, 534, 681, 690; emulsions dissolving wax coating of, 467; resistance of strains of, to lime-sulphur, 343.

Aspidiotus personatus (see Chrysom-

phalus).

Aspidiotus prunorum, on fruit trees in Uzbekistan, 474.

Aspidiotus slavonicus, on poplar and willow in Uzbekistan, 474.

assimilis, Gryllulus (Gryllus). assmuthi, Termes (Odontotermes).

assulta, Heliothis (Chloridea). Asterolecanium bambusae, Cladosporium infesting, on bamboo in

Porto Rico, 501. Asterolecanium miliaris, on bamboo in Porto Rico, 501.

Asterolecanium palmae, on coconut in Jamaica, 239.

Asterolecanium variolosum (on oak), in Australia, 21, 546; in New utilisation Zealand, 546; Habrolepis dalmanni against, 21,

546: tar distillate against, 21. Astragalus aureus, relation Dociostaurus maroccanus to, Transcaucasia, 321.

Astragalus sinicus, identity Phytomyza on, in Japan, 444.

asynamorus, Tachycines. atalantae, Theronia.

Ataxia hubbardi, unsuitable for use against Xanthium pungens Australia, 546. ater, Hylastes.

Athalia rosae (colibri, spinarum) (on crucifers), in Denmark, 117; in Germany, 110; in Russian Union, 313, 439; tests of potential insecticides against, 313.

athasaria, Ellopia.

Athesapeuta cyperi (attacking nutgrass), introduced into Australia and Fiji from Hawaii, 367, 546.

Athous obscurus (haemorrhoidalis),

in Britain, 96.

atlanticus, Smerinthus ocellatus.

Atomaria linearis (on beet and mangel), in Britain, 332, 674; in Denmark, 117.

atomus, Anagrus. atra, Phyllotreta.

Atreides plebeja, parasitised by Apanteles congregatus in U.S.A., 253.

atricapillus, Chrysocoris (see C. javanus).

atricornis, Phytomyza.

Atriplex, beet pests on: in New South Wales, 639; in Poland, 351, 352; in Russian Union, 152, 322, 677.

atropos, Acherontia.

Atta, in Argentina, 87; in Brazil, 271; methods for studying structure of nests of, 271, 567; measures against, 87, 287, 567.

Atta cephalotes, in Dutch Guiana, 567.

Atta sexdens, in Dutch Guiana, 567.

Atta texana, importance of, in Texas, 287.

Attagenus japonicus, in stored tobacco in Japan, 210.

Attagenus pellio, habits of, in Germany, 7, 224, 513; measures against, 513.

Attagenus piceus, infesting house insulation in U.S.A., 640.

attenuata, Estola; Psylliodes.

aucubae, Macrosiphum (see M.

solani).

Aulacaspis pentagona, food-plants and biological control of, in Brazil, 34, 86, 87; on papaya in Porto Rico, 435; oil emulsions against, 34, 435.

Aulacophora (see Ceratia).

Aulacorthum (see Macrosiphum).

aurantiaria, Erannis.

aurantii, Aonidiella (Chrysomphalus); Prospaltella; Pulvinaria; Scirtothrips.

aurata, Pyrausta.

auratus, Carabus; Dirhinus; Rhynchites.

aurea, Leskia.

auricularia, Forficula.

Auriga custos, predacious on Leptinotarsa decembineata in France, 107.

auripennis, Xylocopa. auronitens, Entedon.

aurulenta, Buprestis.
austauti, Smerinthus populi.

Austracris guttulosa, in Queensland, 602.

australe, Dolichocolon.

Australia, forest pests in, 262, 545, 546; Lyctus in timber in, 509; miscellaneous pests in, 21, 544, 545; Ctenolepisma longicaudata in, 510; grasshoppers in, 56, 476, 545; Helicobia australis associated with grasshoppers in, 57; Megastigmus spp. in, 565; Pentalonia nigronervosa transmitting disease of banana in, 289; termites in, 90, 180, 282, 545, 546; beneficial insects and biological control of insects and plants in, 56, 262, 282, 294, 367, 459, 504, 544, 545, 546; criticism and summaries of plant quarantines in, 184, 274, 344, 592; Gambrus stokesi introduced into U.S.A. from, 382. (See under the various States.)

Australia, South, Aonidiella aurantii on Citrus in, 292; orchard pests in, 159, 292; vine pests in, 164; Chortoicetes terminifera in, 345.

Australia, Western, Austroicetes cruciata in, 56, 291; Lepidosaphes ulmi on apple in, 130; pests of pastures in, 131, 292; legislation dealing with control of fruit-flies in, 130; beneficial insects and biological control in, 56, 131, 292, 294; Biscirus lapidarius introduced into New Zealand from, 551; Microphanurus basalis introduced into Queensland from, 603.

australia, Aceratoneuromyia (see Syntomosphyrum indicum); Pseudomicromelus (Dibrachys).

australicum, Trichogramma.

australis, Harpobittacus; Helicobia; Leptoglossus; Oedaleus; Typhlocyba (see T. froggatti).

Austria, Brotolomia meticulosa on Cyclamen in, 171; orchard pests in, 232, 690; Lygus spinolae on vines in, 45; new Cecidomyiid on Trisetum flavescens in, 635; toads destroying insects in, 160;

INDEX. 723

parasites of Galerucella luteola introduced into U.S.A. from, 324.

austriaca, Anisoplia.

Austroicetes cruciata (jungi), in Australia, 56, 130, 291, 545; phases of, 545; Cyrtomorpha flaviscutellaris destroying eggs of, 56; baits for, 130, 291.

Autogiro, for applying dusts and sprays against forest pests, 74,

279, 645.

Autoserica (see Aserica).

avellanae, Eriophyes.

avenae, Macrosiphum (Amphoro-

phora, Aphis).

Avocado, Papilio scamander grayi on, in Brazil, 23; new Eriophyid on, in California, 288; pests of, in Fiji, 641; Xyleborus morstatti on, in Mauritius, 611; Eutermes costalis attacking, in Porto Rico, 464.

Azalea, effects of insecticide treatments on, 605.

Azotus, hosts of species of, 385.

Azotus delhiensis, sp. n., parasite of Aleurolobus barodensis in India, 385.

Azya trinitatis, establishment of, against Aspidiotus destructor in Porto Rico, 502; introduced into Santo Domingo, 502.

B.

baccarum, Dolycoris.

bacchus, Rhynchites.

Bacillus cereus, measures against, causing disease in *Xylomyges* in breeding cages, 81; toxicity of, to cockroaches, 81.

Bacillus sphingidis, causing disease in Protoparce spp. in U.S.A., 251,

253.

Bacteria, Beneficial, 100, 251, 253, 269, 300, 302, 307, 455, 496, 587, 597, 602, 666; methods of culturing and utilising, 307, 455.

Bacteria, Injurious, 302, 463; relation of insects to, 167, 563.

Bactra truculenta (attacking nutgrass), introduced into Australia and Fiji from Hawaii, 367, 546.

badius, Termes; Xyleborus. Bagrada hilaris, on crucifers in

S. Rhodesia, 125.

bahiense, Monalonion. baileyi, Calepitrimerus.

Bait-sprays, for Anastrepha, 256; doubtful value of, for Dacus

oleae, 270, 453; for Lepidoptera, 232, 300, 301; formulae for, 300. Baits, for ants, 44, 87, 112, 323, 435, 665; for Coleoptera, 28, 97, 127, 185, 249, 250, 321, 322, 323, 346, 373, 466, 501, 656, 679, 683; for crickets, 124, 652; for Ctenolepisma, 510; for Dysdercus, 164; for Forficula, 510; for locusts and grasshoppers, 67, 92, 93, 130, 134, 240, 260, 291, 320, 476, 477, 478, 479, 480, 537, 570, 602, 637; for moths, 20, 31, 32, 89, 89, 192, 194, 273, 293, 300, 494, 496, 582, 682, 684; for Lepidopterous larvae, 32, 67, 125, 178, 213, 252, 287, 368, 379, 380, 390, 595, 609; for Trypetids, 98, 131, 270; for other Diptera, 116, 200, 520; formulae for, 32, 67, 89, 92, 98, 102, 124, 127, 131, 249, 250, 260, 270, 320, 321, 322, 323, 379, 390, 476, 478, 494, 496, 510, 520, 595, 602, 609, 652; types of containers for, 98, 435; mixer for, 478.

bajaria, Erannis (Hybernia).

bajulus, Hylotrupes.

Balaninus (see Curculio).

ballestrerii, Megastigmus (Trogocarpus).

balteatus, Syrphus.

Baluchistan, locusts in, 485, 559, 560; orchard pests in, 50-52, 577.

Bambarra Ground-nuts (see Voand-zeia subterranea).

Bamboo, Ceracris kiangsu damaging, in China, 451; pests of, in Formosa, 340, 363; carpenter bees in, in India, 337; new Coccid on, in Japan, 624; pests of, in W. Indies, 181, 423, 501; resistance of introduced species of, to Dinoderus minutus in Porto Rico, 501; soaked in water against Dinoderus spp., 181.

Bamboo Locust (see Ceracris

kiangsu).

Bamboo Powder-post Beetle (see Dinoderus minutus).

bambusae, Asterolecanium; Chaetococcus; Idiococcus.

Banana, pests of, in Australia, 184, 289, 329, 330, 346; Temnoschoita quadripustulata on, in Dahomey, 348; pests of, in Fiji, 213, 366; Papuana huebneri on, in Gilbert Islands, 366; Colaspis hypochlora on, in Br. Guiana, 524; Cosmopolites sordidus on, in Mysore, 291; Dacus psidii on, in New

Caledonia, 131; Erionota thrax on, in Philippines, 451; C. sordidus on, in Tahiti, 365; pests of, in W. Indies, 239, 288, 289; Pentalonia nigronervosa and virus diseases of, 213, 288, 289; cold water treatment of rhizomes of, against C. sordidus, 291.

Banana Rust Thrips (see Scirto-

thrips signipennis).

Banana Scab Moth (see Nacoleia octasema).

Banana Weevil (see Cosmopolites sordidus).

bancrofti, Tarsonemus.

Banding, against *Cydia* spp., **5**, **31**, **114**, **197**, **273**, **293**, **383**, **579**, **588**, **623**, **639**; against other Lepidoptera, **7**, **28**, **458**, **663**, **689**; treatment of materials for, **197**, **293**, **623**, **639**. (See Adhesives.)

Banyan, Chinese (see Ficus retusa). Baobab (see Adansonia digitata).

Barathra brassicae, measures against, in China, 451; on tobacco in Japan, 210; on crucifers in Russian Union, 109, 439, 589; Trichogramma spp. reared on, 35, 36, 109, 391, 589.

Barbados, cotton pests in, 223, 660, 661; sugar-cane pests in, 660; Coptotermes in building timber in, 56, 384; beneficial insects and biological control in, 660; legislation against introduction of Anthonomus grandis into, 661.

Barbarea vulgaris, relation of wireworms to, in Siberia, 679.

barbata, Platygenia; Sarcophaga.

barbatum, Stromatium.

Barbitistes constrictus, attacking conifers in Germany and Poland, 401; not predacious on Lymantria monacha, 401.

Baris interstitialis, introduction of, into Australia against Xanthium

pungens, 546.

Barium Chloride, uses of, as an insecticide, 232, 323, 470, 473.

Barium Chromate, spraying with, against *Plusia orichalcea*, **577**.

Barium Fluoride, tests of toxicity of, to insects, 134, 627.

Barium Fluosilicate, in baits, 124, 125, 478, 510, 537; dusting with, 139, 177, 199, 244, 247, 286, 373, 425, 431; carriers for, 139, 177, 247, 286, 431; spraying with, 70, 71; tests of toxicity of, to insects, 286, 373, 537, 627; formulae containing, 70, 124, 177, 247, 286, 425, 431; and Bordeaux

mixture, 70; and cubé, 177; magnesium oxide as a corrective for foliage injury by, 425; penetration of leaves by, 309; problem of residues of, 177, 432; Dutox form of, 70, 425; other insecticides compared with, 177, 247, 627.

Bark-beetles, of Russian Far East, 456; methods for calculating infestation by, 219; natural enemies of, 324, 339, 397, 473, 522; associated with Ophiostoma ulmi on elm, 65, 68, 77, 378, 396, 397, 529; associated with fungi on Abies, 284; feeding of, 391. (See Scolytids.)

Bark-beetles, Elm (see Hylastes rufipes and Scolytus multistriatus).

Barley, pests of, in Britain, 122, 562; pests of, in Czechoslovakia, 468, 469; pests of, in Denmark, 117; pests of, in Japan, 208, 209, 556; Mayetiola destructor on, in Morocco, 691; pests of, in New Zealand, 547, 548; Syringopais temperatella on, in Palestine, 127; pests of, in Russian Union, 312, 454, 554; thrips on, in Sweden, 121; pests of, in U.S.A., 32, 284.

Barley (Stored), pests in, and their control, 119, 120, 196, 318, 319. Barley Aphis (see *Brachycolus noxius*).

barodensis, Aleurolobus.

Barriers, against grasshoppers and locusts, 214, 476, 478.

Barringtonia speciosa, Dacus passiflorae on, in Fiji, 212.

basalis, Microphanurus; Milionia.

basilinea, Trachea.

basizonus, Microcryptus.

Bassus [Braconid] (see Microdus).
Bassus [Ichneumonid], parasite of
Cydia molesta in Victoria, 640.

Basswood (see *Tilia glabra*).

batatae, Euscepes (see E. post-fasciatus).

Baumspritzmittel, 45, 65, 298. (See Tar Distillates.)

Bean Aphis, Black (see Aphis fabae).

Bean Beetle, Mexican (see Epi-

lachna varivestis).
Bean Bruchid (see Bruchus obtectus).

Bean Fly (see Agromyza phaseoli). Bean Thrips (see Hercothrips fasciatus).

Beans, Heliothis armigera on, in S. Africa, 489; Aphis fabae on, in Britain, 674; pests of, in Fiji, 641; Cantharis obscura on, in

INDEX. 725

France, 623; pests of, Germany, 334, 686; pests of, in Japan, 208, 618; Maruca testulalis on, in Mauritius, 611; pests of, in New South Wales, 58, 449; Piesma quadratum not surviving on, in Poland, 351; pests of, in U.S.A., 32, 69, 77, 82, 142, 240, 340, 370, 421, 432, 599, 600; time of sowing and varieties of, in relation to A. fabae, 334; Ceresa bubalus reared on, 390; leaves of, trapping small insects, 142; Aphids and virus diseases in, 370, 686; Rhynchota associated with Nematospora in, 389; translocation of derris in, 234; tests of toxicity of insecticides to, 79, 422, 426.

Beans (Stored), pests and their control in, 173, 199, 208, 521; nutritive value of groats of, for

Ephestia kuehniella, 64.

Beans, Broad or Horse (see Beans). Beans, Lima (see Lima Beans). Beans, Soy (see Soy Beans).

Beans, Velvet (see Stizolobium).

Beauveria spp., infesting insects, 39, 63, 91, 92, 306, 307, 334, 608, 688; experiments with, and attempted utilisation of, 92, 105, 106, 306, 307, 688.

beckii, Lepidosaphes.

Bedellia somnulentella, parasites of, on bindweed in Kansas, 375.

bedelliae, Apanteles.

Beech, pests of, in Germany, 299, 353, 354; new mite on, in Holland, 301; Barbitistes constrictus feeding on, 401; effect of feeding Lymantria monacha on, 399.

Bees, Carpenter (see Xylocopa).

Bees, Honey, Acarapis woodi attacking, 169; insect enemies of, 238, 515, 631; experimentally infested with Beauveria bassiana, 688; question of risk of insecticides to, 4, 98, 101, 156, 231, 301, 402.

Beet, pests of, in Britain, 96, 673, 674; legislation restricting importation of, into Britain, 356; Cicadella viridis on, in China, 451; pests of, in Czechoslovakia, 468, 469, 470; pests of, in Denmark, 117; Hymenia recurvalis on, in Fiji, 641; pests of, in Germany, 686, 687, 688; Cleonus mendicus on, in Italy, 301; pests of, in Morocco, 283; H. recurvalis a potential pest of, in

New South Wales, 639; Piesma spp. on, in Poland, 105, 163, 350-352; pests of, in Russian Union, 152, 321, 322, 407, 677, 678, 679, 680, 684; book on pests and diseases of, in Russian Union, 439; Loxostege sticticalis on, in Saskatchewan, 27; pests of, in U.S.A., 286, 381, 412, 593; insects and virus diseases of, 163, 286, 351, 356, 412, 439, 686, 688; barium chloride causing injury to, 470; trap-strips of, as a protection against pests of Scorzonera tau-saghyz, 679, 680; mites in stored seed of, 686.

Beet Bug (see *Piesma quadratum*). Beet Leafhopper (see *Eutettix tenellus*).

verienus)

Beet Webworm (see Loxostege sticticalis).

Beet Weevils (see *Cleonus mendicus* and *C. punctiventris*).

Begonia, Monalonion flavisignatum on, in Brazil, **566.**

behri, Etiella.

Belgium, Leptinotarsa decemlineata in, 106, 583; legislation against introduction of L. decemlineata into Britain from, 356.

bella, Strumia.

Bemisia, transmission of cassava mosaic by, in Africa, 94, 153; transmitting virus disease in tobacco in Africa and Netherlands Indies, 124, 125, 360, 493, 637; review of food-plants and distribution of species of, 637.

Bemisia gossypiperda, 94; on cotton in India, 560, 576; possibly associated with "crazy top" in cotton in Sudan, 574; measures against, 560.

Bemisia longispina, characters of, 637.

Bemisia manihotis, characters of, 637.

Bemisia nigeriensis, **94.**

Bemisia rhodesiaensis, measures against, transmitting leaf-curl in tobacco in S. Rhodesia, 124, 125.

Bemisia vayssierei, sp. n., possibly transmitting disease of tobacco in Madagascar, 637.

beneficiens, Telenomus (Phanurus).

Bent Grass (see Agrostis).

Bentonite, toxicity of dusts of, to Bruchus obtectus, 531; and nicotine sulphate, dusting with, against Pyrausta nubilalis, 179; uses of, in sprays, 133, 179, 272, 273, 525, 532, 545; problem of

residues of, on apples, 133, 525; formulae containing, 133, 179, 532; preventing detonation of sulphur nitride, 79; comparison of types of, 545.

Bentonite Sulphur, in mixed sprays,

664.

Benzene, soil treated with, against Leptinotarsa decemlineata, 583; fumigation of stored grain with, against mites, 319; as a solvent, 332, 493; toxicity of derivatives of, to Cydia pomonella, 419.

Ber (see Zizyphus jujuba).

bergenstammi, Apiomyia (Oligotrophus).

bergi, Neomaskellia. bergrothi, Helopeltis.

berlesei, Caloglyphus; Prospaltella. Bermuda, risk of introduction of Popillia japonica from U.S.A. into, 85, 86.

Bessa selecta (see Ptychomyia). Besserioides sexualis, gen. et sp. n., parasite of Dysdercus sidae in Queensland, 57.

Beta-naphthol, bands treated with, against Cydia spp., 197, 293, 639; and oils, formula containing, 197 unsatisfactory in sprays against Cydia pomonella, 134.

Betula (see Birch).

Betula japonica, Lamiids on, in Russian Far East, 42.

Betula lenta, resistant to Phyllotoma nemorata in Quebec, 275.

betulae, Byctiscus; Kuwania; Pulvinaria; Rhynchites.

betuleti, Byctiscus (see B. betulae). Bhima idiota (see Pyrosis).

biannulipes, Peregrinator.

Bibio hortulanus, on barley in Denmark, 117.

bicolor, Leefmansia; Metriona; Olibrus; Phloeosinus; Rhynchites (Merhynchites); Spinaria.

bidens, Picromerus.

bifasciatus, Anastatus; Leptomastix. bifida, Cerura.

biformis, Platypus.

Big Bud Mite (see Eriophyes ribis). Bigonicheta setipennis, liberation of, against Forficula auricularia in U.S.A., 379.

biguttatus, Coptocercus.

biguttula, Empoasca (Chlorita). biguttulus, Chorthippus; Cremastus

(see C. flavo-orbitalis).

bilineata, Cidaria (Hydriomena); Lema.

bilineatus, Prosoestus (Derelomus).

Bill-posters' Paste, as an adhesive for cryolite dips, 380, 652.

bimaculata, Virachola (Deudorix).

binotalis, Crocidolomia.

binotatus, Anisodactylus; Stenotus. Bioclimatics, treatise on, 325–327. bioculatus, Paratetranychus (Tetranychus).

Biological Control, general discussions and papers on, 161, 302, 392, 436, 514.

Biosteres, parasite of Dacus spp. in Fiji, 212.

Biosteres (Opius) fijiensis, parasite of Dacus spp. in Fiji, 212.

Biosteres longicaudatus, parasite of Dacus ferrugineus in Madras, 577. Biosteres persulcatus, parasite of Dacus ferrugineus in Madras, 577. Biosteres tryoni, parasite of Dacus

in Fiji, 213. Lachnosterna (Phyllo-

bipartita,

phaga). bipunctata, Adalia. bipunctatus, Nephotettix. bipunctifer, Schoenobius. bipustulatus, Chilocorus.

Birch, Melolontha hippocastani on, in Finland, 267; Phyllotoma nemorata on, in Quebec, 275; Polyphylla fullo damaging roots of, in Germany, 522; food-plant of Cryptorrhynchus lapathi, 563; Pyrausta nubilalis experimentally fed on, 229; Asiatic Coleoptera in timber of, 465.

Birds, destroying noxious insects, 78, 97, 100, 105, 114, 122, 159, 217, 251, 253, 387, 394, 397, 449, 465, 470, 496, 504, 588, 608, 632, 633; destroying beneficial insects, 159, 282, 550, 564; question of danger of arsenical insecticides to, 159; grain mites spread by, 552; relation of household pests to nests of, 101, 513.

bisaltide, Doleschallia.

Biscirus lapidarius (predacious on Smynthurus viridis), occurrence and utilisation of, in Australia and New Zealand, 131, 292, 551. biselliella, Tineola.

Cassava Bitter (see Manihot utilissima).

Bittersweet (see Solanum dulcamara).

bituberculatus, Dolichoderus; Tetranychus.

Emulsion, Bitumen against Hylemyia brassicae, 116.

bivittata, Metriona. bivittatus, Melanoplus. INDEX. 727

Bixa orellana, Calotermes papua on, in New Guinea Territory, 544.

bizanensis, Microterys.

Black Cyanide, Russian form of calcium cyanide (q.v.), 149, 683.

Black Leaf, 155, 534.

Blackberry, Aphid on, in Britain, 333; Carposina adreptella on, in New Zealand, 37; risk of introducing insects into New Zealand to destroy, 449; pests of, in 79, 178; U.S.A., Dasyneura plicatrix on, 673.

blackburni, Lamprosema.

Blackthorn (see Prunus spinosa).

blanda, Systena taeniata.

blandiatrix, Eutelia.

Blastophagus (see Myelophilus). Blatta orientalis, temperature and

humidity relations of, 224. goniaeformis

Blepharipeza Tritaxys).

Blissus hirtus, damaging lawns in Canada, 601: recorded as B.

leucopterus, 601.

Blissus leucopterus, B. hirtus recorded as, in Canada, 601; measures against, damaging lawns in Trinidad, 25; on cereals in U.S.A., 261, 274, 284, 374; attractiveness of different cereals for, 284; resistance of varieties of Sorghum and wheat to, 274, Collops quadrimaculatus 374; predacious on, 261.

Blood Albumen, as a wetting agent for sulphur, 257, 460; as an emulsifier for wool grease, 426; basic copper arsenate compatible

with, **421.**

Boarmia selenaria dianaria, adaptation of, to Citrus in Sicily, 630.

boas, Oryctes.

Boehmeria nivea, pests of, in Transcaucasia, 587.

boerneri, Chermes (Pineus).

boetica, Cosmolyce (Lampides). measures flavolineatus, against, on vines in Brazil, 34.

Bolivia, summary of plant quarantines in, 592.

bombina, Cratosomus.

Bombyx mori, 586; experimentally infested with Beauveria bassiana, 688; Dermestids attacking, 196, 198; review of literature_ on Arthropods injurious to, 224; parasitised by Sturmia sericariae in Japan, 442, 443; used in tests of insecticides, 298, 688.

Conotrabondari, Chalcodermus;

chelus; Monalonion.

boninsis, Pseudococcus.

Books, pests of, and their control in Fiji, 366.

Borassus, destruction of, against Oryctes spp. in Dahomey, 347.

Borax, in bait for Trypetids, 98, 270; effect of using, to remove spray residues from apples, 648; and boric acid, treatment of wood with, against Lyctus brun-

neus, 509, 510.

Bordeaux Mixture, use of, against Aesiotes, 602; against Empoasca and Epitrix, 53, 70, 342, 501; repellent to Leptinotarsa, 584; Monalonion, against 566: apparently reducing virus infection of potato, 157; formulae for, 70, 278, 342, 425, 536, 602; formulae containing, 70, 125, 127, 192, 278, 342, 422, 425, 536; and arsenicals, 34, 70, 125, 139, 188, 192, 207, 235, 342, 421, 534, 583, **643, 653, 662**; interactions of, with calcium arsenate, 171, 373; and barium fluosilicate, 70; and cryolite, 425; and derrisol, 127; and nicotine, 125, 501, 534, 536; and oil emulsions, 188, 192, 278, 497, 536; thiuram sulphides compatible with, 532; causing foliage injury, 643; magnesium oxide as a corrective for injury by, 425; interactions of, with other sprays on Citrus, 278.

borgmeieri, Adesmus.

Boric Acid, treatment of timber with, against Lyctus, 509; use of, in removing spray residues from apples, 648, 649.

Borkhausenia pseudospretella, damaging wine corks in Scotland, 122; characters of eggs of, 49.

bormansi, Dolichopoda.

Borreria verticillata, Larra americana associated with, 85.

Bostrychoplites cornutus, on Vitex timber in Mauritius, 611.

Botanical Collections, damaged by Trogoderma tarsale, 196.

Bothriophryne, key to species of, 503; characters of, 504.

Bothriophryne dispar, sp. n., parasite of Ceroplastes in Tanganyika, 503.

Bothriophryne fuscicornis, sp. n., parasite of Ceroplastes in S. Africa, 503.

Bothriophryne purpurascens, sp. n., parasite of Ceroplastes in S. Africa, 503.

Bothriophryne (Trichomasthus) tenuicornis, 504.

Bothynoderes punctiventris (see Cleonus).

botrana, Polychrosis.

Botrytis bassiana (see Beauveria). boucheanus, Dibrachys (see D cavus).

bowringi, Colaphellus.

Box-elder (see Acer negundo).

Box-elder Bug (see Leptocoris trivit-tatus).

brachialis, Bruchus.

Brachmia macroscopa, effect of temperature on, on sweet potato in Korea, 362.

Brachonyx pineti, bionomics and control of, in Rumania, 355.

Brachonyx pineti var. obscurella, on pine in Rumania, 355.

Brachyacma palpigera, on Tephrosia toxicaria in Porto Rico, 502.

Brachycolus noxius, measures against, on wheat in Central Asia, 453, 454.

Brachylacon (see Lacon).

Brachymeria, parasite of Earias and Platyedra in Philippines, 184.

Brachymeria euploeae, associated with Virachola isocrates in India, 385; bionomics of, parasitising Oreta carnea in Sumatra, 385, 386.

Brachymeria excarinata (parasite of Plutella maculipennis), bionomics of, in India, 575, 576.

Brachymeria obscurata, hosts of, in Japan, 211.

Brachymeria vitripennis, parasite of Cassida nebulosa in Morocco, 284.

brachypterus, Microcryptus.

Brachystegia, 62.

Brachytrypes membranaceus, measures against, on tobacco in S. Rhodesia, 124.

Brachytrypes portentosus, soil characters in relation to, in sugar-cane fields in Formosa, 206.

Bracon abscissor (see Microbracon). Bracon nicevillei (see Glyptomorpha). Bracon triangularis (see Microbracon).

Braconids, list of, in Germany, 113.
Bran (including pollard), insects infesting, 64, 224; in baits, 32, 67, 68, 92, 130, 134, 178, 240, 249, 260, 291, 321, 390, 476, 478, 511, 520, 537, 602; baits for Trypetids prepared from, 98, 270; formulae containing, 32, 92, 260, 321, 390, 476, 511, 520, 602.

brasiliana, Acaulona.

brasiliensis, Clistomorpha (Hyalomyodes); Eurhizococcus; Gasterocercodes; Hyalomyia (Alophoropsis); Pseudiastata.

brassicae, Aleurodes; Barathra; Brevicoryne (Aphis); Dasyneura; Hylemyia (Chortophila, Delia); Phaedon; Pieris; Plusia (Autographa); Tyloderma.

Brassolis sophorae, on coconut in Br. Guiana, 524.

braueri, Tritaxys.

brauneri, Otiorrhynchus.

Brazil, pests of Anonaceae in, 566, 567; new weevil on cabbage in, 132; cacao pests in, 566, 662; pests of Cocos spp. in, 22, 661; coffee pests in, 34, 188, 238; cotton pests in, 22, 34, 86, 136, 413; pests of grape-vines in, 34; Phthorimaea operculella in, 289; miscellaneous pests in, 23, 86, 290; new Coccids in, 48, 623; Nosema infesting Lepidoptera in, 459; beneficial insects and biological control in, 22, 23, 34, 85, 86, 87, 137, 183, 188, 238, 253, 289, 414, 505; Larra americana introduced into Porto Rico from, 85, 435.

brevicollis, Dociostaurus crucigerus.

brevicomis, Dendroctonus.

brevicornis, Wesm., Microbracon. brevicornis, auct., Microbracon (see

M. hebetor).

Brevicoryne brassicae (on crucifers), in Britain, 332; measures against, in China, 450; resistance of, to abnormal cold in France, 622; relation of, to mosaic disease of turnip, etc., in New York, 274; in S. Rhodesia, 125; in Russian Union, 439; in Tasmania, 21.

brevipes, Pseudococcus.

brevis, Calotermes (Cryptotermes); Dinoderus; Hoplocampa; Stylocryptus.

brevistigma, Tetrastichus.

brevistylus, Dacus (see D. ciliatus).

brevitarsis, Liocola.

brevivalvus, Megastigmus (Epime-gastigmus).

Brithys pancratii, on amaryllidaceous plants in S. Rhodesia, 125. British Columbia, Coccids of, 511; miscellaneous pests in, 66, 568; proposed introduction of parasites from Britain into, against Phytomyza ilicis, 669.

British Isles, cereal pests in, 96, 122, 332, 562; forest pests in, 58, 262, 391, 504; bark-beetles associated with Dutch elm disease in, 65; pests and diseases of

willows in, 562, 563, 627; pests of greenhouse and ornamental plants in, 115, 295, 333, 356, 437, 512, 519, 669, 673; pests of hops in, 1, 333; mushroom mites in, 1; orchard pests in, 236, 332, 333, 676; pests of small fruits in. 2, 236, 333, 395, 596; list of Coleoptera associated with fruits in, 160; spray programmes for fruits in, 396; pests of stored products in, 122, 154, 265, 623, 638, 666; pests of turf and grasses in, 95, 122, 635; pests of vegetables and root crops in, 38, 61, 96, 115, 116, 122, 236, 262, 265, 332, 333, 562, 563, 578, 671, 672, 673, 674, 675; problem of wireworms in, 96; Pyrausta nubilalis in, 116, 295; Stenomalus muscarum in houses in, 201; new Thomasiniana on hawthorn in, 673; Aphids and virus diseases in, 236, 395, 563, 671, 672; beneficial insection, 38, 58, 95, 115, 122, 262, 265, 390, 391, 504, 562, 628, 632, 669; utilisation of *Encarsia formosa* in, 333, 437; new mite predacious on Thrips tabaci in, 672; plant pest legislation in, 355, 356; introduction of beneficial insects into other countries from, 38, 39, 162, 262, 282, 504, 669.

bromeliae, Diaspis.

Bromus, relation of Dociostaurus maroccanus to, in Transcaucasia, 321; new Anthomyiid on, in Siberia, 312.

Bromus secalinus, Ochsenheimeria taurella and white ear in, in Russian Union, 680.

Brontispa limbata, on palms in Mauritius, 611.

Brood, meaning of term, 137.

Brotolomia meticulosa, bionomics of, on Cyclamen in greenhouses in Austria, 171.

Brown-tail Moth (see Nygmia phaeorrhoea).

Bruchidius (see Bruchus).

Bruchobius mayri (see Sphaerakis). Bruchocida orientalis, parasite of Pempheres affinis in India, 506.

Bruchophagus gibbus, on lucerne in Russia, 108, 475.

Bruchus analis, bionomics of, on lima beans in Japan, 617; intercepted in Japan, 208.

Bruchus brachialis (Vetch Bruchid), parasites and biological control of, in U.S.A., 381, 599.

Bruchus chinensis, fumigation against, in China, 451; bionomics of, in Japan, 208, 361, 557, 616, 617.

Bruchus cisti, on wild lupins in Italy and Sicily, 201, 495; not found in Iberian Peninsula, 495.

Bruchus maculatus, intercepted in beans in Japan, 208; larval anatomy and hatching of, 280.

Bruchus obtectus (Bean Bruchid), in Italy, 199, 200; intercepted in Japan, 208; nutritional studies on, 624; experiment with Trichogramma evanescens and, 36; tests of insecticides against, 199, 200, 531.

Bruchus phaseoli, bionomics of, on lima beans in Japan, 617.

Bruchus pisorum (Pea Bruchid), in Czechoslovakia, 468; on beans in Japan, 208; in Russia, 147; in U.S.A., 325, 456; Triaspis thoracicus introduced into U.S.A. against, 381; bionomics of, 208, 325, 456; X-rays for detecting, in stored peas, 147; measures against, 325, 456.

Bruchus quadrimaculatus (see B. maculatus).

Bruchus rufimanus, bionomics of, on beans in Japan, 208; not occurring in Formosa, 208.

Bruchus seminarius, on wild serradella in Portugal, 495.

Bruchus terrenus, bionomics of, on

beans in Japan, 208. brumata, Operophtera (Cheimatobia). brunneus, Heterobostrychus; Lyctus. brunsvicensis, Chrysomela (Chryso-

lina).

Bryobia praetiosa, measures against, in orchards in New South Wales and California, 57, 261, 432.

Bryobia sarothamni, sp. n., on Sarothamnus in Holland, 301.

bubalis, Ceresa.

Bucculatrix gossypii, on cotton in Queensland, 603.

Bucentes geniculata, parasite of Tipula spp. in Britain, 95.

bucyrus, Cactoblastis.

Bufo marinus (destroying insects), in Porto Rico, 656; establishment of, in Fiji, 641; utilisation of, in Florida, 656; probable establishment of, in Philippines, 452; establishment of, in Queensland, 185.

Bufo viridis, observations on, in Russia, 407.

C2

Bufo vulgaris, destroying insects in Austria, 160.

bugandaensis, Anagyrus.

Building Timbers, Coleoptera in, 6, 7, 78, 110, 203, 204, 268, 321, 322, 335, 469, 499, 520, 584, 600; termites in, 55, 56, 125, 180, 239, 464, 465, 498, 499, 545, 641, 693; other pests of, 337, 498, 499, 693.

Bulgaria, new Coccid on roses in, 402.

bulgariense, Lecanium.

Bumelia lycioides, host-fruit of Ceratitis capitata in France, 195. buoliana, Rhyacionia (Evetria).

Bupalus piniarius (Pine Geometrid), in Czechoslovakia, 469; measures against, in Germany, 231, 512.

Buprestis aurulenta, on pines in California, 329; in New Zealand, 329.

Buprestis langi, in dock timber in Û.S.A., 499. buqueti, Thaneroclerus.

Burma, forest and timber pests in, 612, 692, 693; catalogue of Longicorns in, 612; beneficial insects and biological control in, 693; parasites introduced into India from, 612, 693.

Busseola fusca, legislative measures against, on maize in Kenya, 666; new Tachinid parasite of, in S.

Rhodesia, 567.

β butoxy β' thiocyanodiethylether, 155, 236, 258, 332. (See Thiocyanates.)

Butternut (see Juglans cinerea).

Butyl Phenyl-phenol Sodium Sulphonate, 179.

buxi, Pinnaspis.

buxtoni, Calotermes; Margarodes. Buzura suppressaria, bionomics and control of, on tung-oil tree in China, **614.**

Byctiscus betulae (betuleti), on vines in Russian Union, 403.

Bysocrypta fusiformis, susceptibility of varieties of elm to, in Japan, 443.

Bysocrypta gallarum, susceptibility of varieties of elms to, in Japan, 443.

Bythoscopus mundus, on grapevines and willow in Japan, 440.

Byturus unicolor, measures against, on raspberry in Washington, 380.

C.P. 100 (see Cyclohexylamine Derivatives).

Cabbage, pests of, in Australia, 21. 545; new weevil on, in Brazil, 132; pests of, in Britain, 61, 96, 97, 116, 265, 578, 608; Colaphellus bowringi on, in China, 450: Crocidolomia binotalis on, in Belgian Congo, 572; Athalia rosae on, in Denmark, 117; pests of, in Fiji, 641; pests of, in France, 459, 622; pests of, in Germany, 110, 511, 636, 686, 687; Contarinia nasturtii on, in Holland, 514; Plutella maculipennis on, in Jamaica, 239; Forficula auricularia feeding on, in Ontario, 601; pests of, in Russian Union, 109, 153, 306, 307, 310, 316, 320, 455, 589, 679; Pieris brassicae on, in S.A., 32, 33, 249, 250, 370, **372, 379, 541;** susceptibility of varieties of, to pests, 61, 317; Aphid transmitting virus disease of, 686; problem of arsenical residues on, 370.

Cabbage Bug, Harlequin (see Mur-

gantia histrionica).

Cabbage Butterflies (see Pieris brassicae and P. rapae).

Cabbage Fly (see Hylemyia brassi-

Cabbage Whitefly (see Aleurodes brassicae).

Cacao, pests of, in Brazil, 566, 662: Monalonion on, in Colombia, 566; Helopeltis antonii on, in Java, 523; pests of, in New Guinea Territory, 277, 544; Acrocercops cramerella on, in Philippines, 452; thrips on, in W. Indies, 24, 25.

Cacao (Stored), pests in, and their control, 64, 154, 521, 584, 638.

Cacao Thrips (see Selenothrips rubrocinctus).

Cacoecia (see Tortrix).

Cactoblastis bucyrus, C doddi previously identified as, 459.

Cactoblastis cactorum (destroying Opuntia), new Nosema infesting, in S. Africa, 459; utilisation of, in Queensland, 345.

oddi, attempted of, against S. Africa and Cactoblastis doddi, establishment Opuntia in S. Africa and Australia, 459; new Nosema infesting, 459.

cactorum, Cactoblastis.

cadaverum, Glycyphagus.

Cadmium, compounds effective in baits for grasshoppers, 134.

Cadurcia vanderwulpi, sp. n., parasite of Hapalia machaeralis in India, 263.

caementarium, Sceliphron.

caesareus, Staphylinus.

Cages, for rearing and studying insects, 28, 30, 78, 81, 345.

caja, Arctia.

Cajanus cajan (indicus), Corythuca gossypii on, in Jamaica, 239; Etiella behri on, in Seychelles, 336; Inglisia conchiformis on, in Uganda, 40.

calacte, Macrocentrus.

Calamagrostis epigeios, relation of Pentatomid cereal pests to, in

Germany, 631.

Calandra granaria (Grain Weevil), in Germany, 582; in Poland, 162; in Russian Union, 304, 551, 585, 685; in Sweden, 118-120; in U.S.A., 341; movement of, in stored grain, 585; bionomics of, 118-120, 305; parasite of, 304, 551; measures and fumigation tests against, 120, 162, 309, 341, 344, 585, 685; tests of Naaki

dust against, 582.

Calandra oryzae (in stored cereals), in China, 451; in Kenya, 667; parasitised by Lariophagus in Italy, 199; in Japan, 339, 360; mass rearing of Lariophagus on, in Russia, 304; in Sweden, 118-120; in U.S.A., 248, 341; bionomics of, 104, 118-120, 305; feeding and oviposition preferences of, 64, 65, 119, 339; water content of adults of, killed by desiccation, 585; uses and tests of fumigants against, 120, 341, 451; other measures against, 120, 248, 360, 667.

Calandra sasakii, measures against, in stored rice in Japan, 360.

calandrae, Aplastomorpha. calcarator, Zele (Phylacter).

Calcid, 165.

calcitrator, Collyria.

Calcium Arsenate, against beet pests, 283, 451, 470; against Lepidoptera on cabbage, 370, 380; against cotton pests, 164, 243, 244, 245, 246, 247, 380, 417, 418; effects of, on infestation by Aphis gossypii, 164, 243; against forest pests, 74, 428, 461, 645; against pests of leguminous plants, 82, 108, 258, 373; against orchard pests, 70, 273, 383, 528, 610, 662, 664; against pests of solanaceous plants, 67, 70, 158,

342, 372, 431, 525, 583, 634; against pests of sweet potato, 286, 373, 445; against pests of other crops, 191, 364, 378, 426, 441; in bait against Heliothis, 609; soil treatment with, against Lachnosterna, 425, 447; tests of toxicity of, to insects, 243, 246, 249, 277, 286, 528; application of, from aircraft, 74, 610, 645; formulae for, in dusts, 191, 243, 244, 245, 246, 247, 277, 283, 342, 372, 380, 417, 418, 426, 461; carriers for, 191, 246, 247, 277, 283, 372, 380, 418, 426, 461; spraying with, 70, 158, 159, 247, 273, 283, 342, 373, 383, 441, 445, 451, 470, 528, 529, 583, 610, 634, 661, 664; adhesives and spreaders for, 70, 373, 445; mopping mixture containing, 247; and cubé, 417; and bentonite sulphur, 664; and Bordeaux mixture, 70, 342, 373, 584, 662; effects of, on suspension capacity of Bordeaux mixture, 171; and lime, 283, 451, 664; effect of addition of magnesium sulphate and lime to, 158, 634; and limesulphur, 664; and nicotine sulphate, 529; and oil emulsions, 529; and Paris green, 243, 244, 245, 417, 418; and sodium fluosilicate, 283; and sulphur, 244, 245, 246, 247, 380, 417; particle size and other factors affecting toxicity of, 243, 246, 528; preparation and forms of, 372, 373, 643; standards for, 115: and injury to plants, 82, 247, 248, 380, 529; correctives for foliage injury by, 373; residues of, on grapes, 191; question of danger of, to birds, 159; other insecticides compared with, 79, 243, 244, 245, 246, 247, 249, 250, 373, 417, 441, 470, 525, 627, 634.

Calcium Arsenite, dusting with, against lucerne weevils, 108; toxicity of, to insects and plants,

627.

Calcium Caseinate, as a spreader for sprays, 207, 208, 279, 330, 445; basic copper arsenate compatible with, 421; dusting sulphur made wettable by, 460.

Calcium Chloroacetate, dusting with,

239

Calcium Cyanamide, tests of action of, on insects, 200; experiments with, against Leptinotarsa decemlineata, 107, 158, 159, 349; soil

treatment with, against other

pests, 358, 520, 617, 619.

Calcium Cyanide, against ants, 24, 287; dusting with, 463, 561; fumigation of Citrus with, against Coccids, 149, 150, 165; as a greenhouse fumigant, 357, 526; fumigation of mushroom houses with, 260; against wireworms, 683; causing injury to carnations, 526.

Calcium Fluoride, in baits for

grasshoppers, 134.

Calendula officinalis, not attacked by Micromyzus oliveri in Washington, 541.

Calepitrimerus baileyi, gen. et sp. n., on apple in California, 33.

Calidea, key to species of, in Ethiopian Region, 360; associated with Nematospora and boll rots, 360, 389.

calidella, Ephestia.

California, Citrus pests in, 55, 261, 342, 343, 377, 424, 425, 432-434, 459-461, 649, 650; pests of forest and shade trees in, 140, 144, 284, 329, 569, 599, 649; miscellaneous pests in, 239, 569, 592; orchard pests in, 33, 34, 261, 383, 409, 432, 434, 649, 650; Stromatium fulvum in imported timber in, 160; vegetable pests in, 83, 240, 379, 431, 590, 599, 600; Aphids of, 261, 590; records and new species of Eriophyids in, 33, 55, 288, 409; grasshopper surveys in, 569; Symphyla in, 411; wireworms in, 379, 410; Aphids and virus diseases of celery, etc., in, 590, 591; beneficial insects and biological control in, 30, 133, 144, 410, 424, 434, 437, 569; Lasioderma serricorne imported into Germany in beans from, 521. californica, Plusia (Autographa).

californicus, Pheletes (Limonius); Prionus.

caliginosellus, Crambus.

Caliroa annulipes, on lime in Germany, 204.

Caliroa limacina (cerasi, auct.) (on fruit and nut trees), in Australia, 21, 279; new Tritneptis probably parasitic on, in California, 434; in Germany, 114; bionomics of, 21, 114; measures against, 21, 279.

Callabismodes tabaci, parasite of, in Brazil, 183.

Calliceras kamiyae, sp. n., hyperparasite of Dendrolimus spectabilis in Japan, 91.

Calliceras manilae (see Ceraphron). Callidium antennatum hesperum, in building timber in U.S.A., 499.

Calliephialtes (see Ephialtes).

Callimome druparum, on apple in New York, 533.

Calliptamus italicus, in Central Asia, 627; in Cyprus, 270; financial loss due to, in Greece, 478; in Jugoslavia, 479; in Libya, 478; in Poland, 9; breeding areas of, in Spain, 477; crops attacked by, 9, 478; measures against, 477, 478; tests of arsenical and fluorine compounds against, 627.

Callopistria floridensis, measures against, on greenhouse ferns in

N. America, 592.

callosa, Sitona.

Calocoris norvegicus, on potato in Britain, **332.**

Caloglyphus, bionomics and identity of species of, on mushrooms in Britain, 1.

Caloglyphus berlesei, 1.

Calomel (see Mercurous Chloride).

Calomycterus setarius, distribution of, in U.S.A., **600.**

Calopogonium coeruleum, rotenone in, **502.**

Calopogonium orthocarpum, rotenone in, 502.

Caloptenus italicus (see Calliptamus). Calosoma, Pinthaeus sanguinipes

predacious on, in Germany, **354.** Calosoma inquisitor, predacious on forest pests in Germany and Poland, **354, 438.**

Calosoma sycophanta (predacious on Lymantria spp.), bionomics and suggested utilisation of, in Germany, 170, 354; utilisation of, in U.S.A., 170.

Calotermes, in Australia, 545.

Calotermes brevis, damaging furniture and building timbers in W. Indies, 55, 56, 239, 434, 464, 465; factors increasing damage by, in Jamaica, 239; susceptibility of varieties of timber to, 434, 465; characters of, 55; measures against, 55, 465.

Calotermes buxtoni, in building

timbers in Fiji, 641.

Calotermes papua, measures against, damaging cacao and Bixa in New Guinea territory, 544.

Calotermes repandus, in building timbers in Fiji, 641.

Calotermes taveuniensis, in Fiji, 641. Calotropis gigantea, Poecilocerus pictus on, in India, 577.

Calpe emarginata, bionomics of, in

Japan, 209.

Calpodes ethlius, measures against, on arrowroot in St. Vincent, 271. calthae, Aphalara.

Calymnia trapezina, possibly predacious on Evannis spp. in Poland, 438.

Calyptus strigator, parasite Pissodes validirostris in Finland,

Calystegia japonica, Anophia leucomelas on, in Japan, 445.

camelinus, Amblyteles.

Camellia oleifera, weevil on, in Formosa, 340.

cameroni, Spalangia.

Camnula pellucida, in Canada, 476, 485; in U.S.A., 485; migrations of, 485; organisation of measures against, 476.

Camphor Sassafras Base Oil, as an intermediary solvent for derris in

oil sprays, 530.

camphorae, Gammaphytoptus.

Camponotus herculeanus pennsylvanicus, apparatus for protecting insect specimens from, in New York, 541.

Camponotus laevigatus, possibly in building timber in U.S.A., 499. Camponotus maculatus vicinus, in

building timber in U.S.A., 499.

Campoplex convexus, parasite of Pristiphora wesmaeli in Germany,

camposi, Chalcodermus.

Campsomeris annulata, establishment of, against Lachnosterna smithi in Mauritius, 611.

Campsomeris erythrogaster, establishment of, against Lachnosterna smithi in Mauritius, 611.

Campsomeris marginella modesta, parasite of Anomala orientalis in Hawaii, 367.

Campsomeris rodriguezensis, establishment of, against Lachnosterna smithi in Mauritius, 611.

Campylochaeta, parasite of Sphinx pinastri in Poland, 677.

Campylomma nicolasi, possibly causing "crazy top" of cotton possibly in Sudan, 574.

Canada, cereal pests in, 176, 229, 295, 368; danger of introduction of Agropyrum smithi into, on

account of Cephus cinctus, 368; forest pests in, 25, 53, 67, 176, 368, 369, 540, 647; orchard pests in, 39, 382, 542; pests of ornamental plants in, 592, 673; Cydia nigricana on peas in, 38, 39; Leptocoris trivittatus in, 374; grasshoppers and expenditure on their control in, 476, 477, 485; Aphids of maritime Provinces of, 104; annual insect population records in, 141; beneficial insects and biological control in, 38, 39, 67, 253, 369; regula concerning transportation regulations nursery stock in, 176; beneficial insects introduced into U.S.A. from, 381; legislation restricting importation of potatoes Britain from, 356; Hercothrips fasciatus on oranges imported into Japan from, 616. (See also under the various Provinces.)

canaliculatus, Lyctus (see L. linearis). Canary Islands, Tephritis in, 333. cancellatus, Carabus.

canella, Diatraea.

canescens, Alphus; Nemeritis.

canifrons, Ascogaster.

Cankerworm, Fall (see Alsophila pometaria).

Cannabis sativa (see Hemp).

Cantharis livida, in orchards in France, 623.

Cantharis obscura, food-plants and control of, in France, 622, 623. Cantharis rustica, in orchards in France, 623.

Cantheconidea acuta, predacious on Oreta carnea in Sumatra, 386.

Cantheconidea robusta, parasites of, in Ceylon, 19.

canus, Pheletes (Limonius).

Cape Goosebery, False (see Nicandra physaloides).

capensis, Enneadesmus forficula.

capitata, Ceratitis. capitatum, Piesma.

capito, Locusta migratoria.

Capitophorus, characters and scope of, **99, 100.**

Capitophorus fragariae, and virus diseases of strawberry in Britain, 395: measures against, 395.

Capitophorus korschelti, sp. n., on Alpine and black currant in Germany, 100.

Capitophorus ribis, on currant and gooseberry in Britain, 333; C. korschelti recorded as, in Germany, 100.

capreae, Cavariella.

Capsicum, insects causing leaf-curl of, in Ceylon and India, 493, 494; Myzus persicae transmitting virus disease to tobacco from, in Netherlands Indies, 493; Paratrioza cockerelli on, in Texas, 651; not attacked by Phthorimaea operculella, 289.

capulifera, Pimpla.

Carabus auratus, measures against, on strawberry in Germany, 457. cancellatus, measures Carabus

against, on strawberry

Germany, 457.

Carabus pallipes (see Agonum). Caragana, Etiella zinckenella on, in Russia, 152.

Caraway, Aphids and virus disease of, in California, 590.

Carbolic Acid, tests of repellents containing, against Chrysobothris femorata, 420.

Carbolineum (see Tar Distillates).

Carbon Bisulphide, against borers in oil palms, 347, 348; a Coleoptera in furniture, against against pests of stored products, plant material, etc., 184, 230, 319, 341, 349, 358, 539, 667; not affecting germination of coffee seed, 667; against ants and termites, 25, 56, 287, 567, 693; against other pests in soil, 11, 88, 177, 259, 358, 522, 583, 617, 621, 693; in spray against Parlatoria oleae, 553; emulsions of, 177, 259, 539, 553; fire hazard of, 341, 358; and carbon tetrachloride, 341; and dichlorobenzenes, 88; and naphthalene, 177.

Carbon Dioxide, fumigants used

with, 359, 542.

Carbon Tetrachloride, and ethylene dichloride, as a fumigant, 120, 184, 341, 378, 539; and carbon bisulphide, 341; and toluene, as a solvent for dinitro-o-cyclohexylphenol, 261.

Carcelia kockiana, parasite of Argyroploce illepida in Fiji, 213.

Carcelia lucorum (see Zenilia).

carcharias, Saperda.

Cardamom (see Elettaria cardamomum).

Dysdercus; cardinalis, Rodolia(Novius).

Cardiophorus formosanus, soil characters in relation to, sugar-cane fields in Formosa, 206. cardui, Vanessa (Pyrameis). carduorum, Apion (Ceratapion).

Carica papaya (See Papaya).

carinatus, Cheiloneurus; Eriophyes. Carnation, Tortrix pronubana imported into Germany with cuttings of, 218; pests of, in U.S.A., 79, 525; Tetranychus telarius dis-seminating rust fungus on, 79; insecticides causing injury to, 272, 526.

carnea, Oreta.

Carolina, North, miscellaneous pests in, 377, 380, 599; tobacco pests in, 254, 380; Typophorus viridicyaneus on sweet potato in, 285; beneficial insects and biological control in, 599.

Carolina, South, Epilachna varivestis on beans in, 69; Heliothis armigera in, 179, 180; Horistonotus uhleri in, 500; miscellaneous pests in, 379, 380; beneficial insects, etc., in, 500,

501, 538.

carolina, Dissosteira.

Carolinaia setariae, vector of sugarcane mosaic in Florida, 658.

Carpenter Ants (see Camponotus). Carpenter Bees (see Xylocopa). carpini, Trypopitys (Priobium).

Carpocapsa (see Cydia).

carpocapsae, Ascogaster; Hemiteles. Carpocoris, bionomics and ecology of, on cereals in Germany, 353, **631, 632**; measures against, **353.** Carpocoris pudicus, parasites of,

on cereals in Germany, 632. Carpocoris pudicus fuscispinus, parasites of, on cereals in Germany,

64, 632.

Carpomyia incompleta, intercepted in fruits of Zizyphus in Russian Union, 439.

Carpomyia pardalina (see Myiopardalis).

Carpomyia schineri, intercepted in fruits of Zizyphus in Russian Union, 439.

Carpomyia vesuviana, on Zizyphus jujuba in India, 503; intercepted in fruits of Zizyphus in Russian Union, 439.

Carpophilus, measures against, in oil-palm kernels in Dahomey, 349. Carpophilus hemipterus (in dried

fruit), habits of, in France and

Germany, 167, 201.

Carpophilus humeralis, attacking seed sugar-cane in Hawaii, 367. Carposina adreptella, new parasite of, on raspberry and blackberry in New Zealand, 37.

carposinae, Apanteles.

Carrot, pests of, in Britain, 96;

Psila rosae on, in Czechoslovakia,
468; Forficula auricularia feeding on, in Ontario, 601; Loxostege sticticalis on, in Siberia, 313;
Aphids on, in Tasmania, 21;
pests of, in U.S.A., 376, 496, 590;
Aphids and virus disease of, 590.
Carrot Beetle (see Ligyrus gibbosus).

Carrot Rust Fly (see Psila rosae).

Carum carvi (see Caraway).

caryae, Acrobasis.

caryana, Enarmonia.

casei, Piophila.

Cashew (see Anacardium occidentale).

Casmara patrona, measures against,

on tea in Formosa, 340.

Cassava, new mite on, in Brazil, 290; Coscineuta virens on, in Trinidad, 25; locusts damaging, 571; insects and virus diseases of, 94, 95, 153.

Cassava, Sweet (see Manihot dulcis

var. aipi).

Cassia, Míctis profana on, in Fiji, 641; Xyleutes persona on, in Burma, 693.

Cassida vittata, bionomics and control of, on beet in Morocco, 283, 284.

Castanea sativa (see Chestnut).

castanea, Diparopsis.

castaneum, Tribolium.

Castor, Tetranychus telarius on, in India, 127; susceptibility of, to dichloroethyl ether, 272.

Casuarina, Coccids on, in Porto

Rico, 434.

Catabomba pyrastri (see Lasioph-thicus).

Cataglyphis bicolor, Microbracon greeni reared on, in India, 129.
Catalpa, Sphingids on, in U.S.A.,

252, 655.

catalpae, Ceratomia.

Catascopus facialis, introduced into Samoa against Oryctes rhinoceros, 641.

Cathartus, measures against, in oil-palm kernels in Dahomey, 349.

Cathartus advena (see Ahasverus).
Catocala oberthüri, on poplars and willows in Morocco, 196.

Catorama herbarium, parasite of, infesting books in Fiji, **366**; previously recorded as Sitodrepa panicea, **366**.

Catoxantha (see Chrysochora).

catoxantha, Artona.

caucasicus, Otiorrhynchus; Phenacoccus.

caudata, Lepidiota.

Cauliflower, pests of, in Britain, 96, 116; Contarinia nasturtii on, in Czechoslovakia, 468; Cidaria fluctuata on, in Denmark, 117; pests of, in France, 167, 458, 459, 621; Contarinia nasturtii on, in Holland, 514; Hylemyia brassicae on, in Russia, 317; pests of, in U.S.A., 72, 541; types of injury to, by C. nasturtii, 167; susceptibility of varieties of, to Rhytidoderes plicatus, 459; danger of injury to, by mercury bichloride, 116.

cautella, Ephestia. Cavariella aegopodii, on carrot and

parsnip in Tasmania, 21. .Cavariella capreae, transmitting

virus disease of celery in California, 590; on carrot and parsnip in Tasmania, 21.

cavus, Dibrachys.

Cecidomyiids, list of, attacking apple and pear, 557; studies on food-plant range of, 672.

Cecropia adenopus, Monalonion knighti on, in Brazil, 566.

Cedar (*Cedrus*), pests of, in N. Africa, **58**, **226**.

Cedrela toona, Hypsipyla robusta on, in India, 612; Xylocopa latipes in timber of, 337.

cedri, Phloeosinus.

Cedria paradoxa, liberation of, against Margaronia pyloalis and Hapalia machaeralis in India and Burma, 612, 693.

Cedrus (see Cedar).

Celastrus angulatus, preparation of insecticide from, 450, 451.

Celeriac, Aphids and virus disease of, in California, 590.

Celerio lineata, outbreak of, in California, 239; trap-ditches against, 239.

celerio, Hippotion.
Celery, Aphids on, in U.S.A., 541,

590; list of other Aphids on, 590; virus diseases of, 590, 591. Celite, dust mixture containing, 75.

Cemiostoma scitella (see Leucoptera). Centaurea, relation of Dociostaurus maroccanus to, in Trans-

caucasia, **321.**Centeterus (parasite of *Pyrausta* nubilalis), bionomics of, in Italy,

174. Cephaleia sachalinensis, on pine in Japan, 339.

Cephalobus, parasite of Diplosis mori in Korea, 338.

Cephalobus persegnis, parasite of Horistonotus uhleri in S. Carolina, 500.

cephalonica, Corcyra.

Cephalosporium lecanii, methods of utilising, against Ceroplastes sinensis in Russian Union, 308.

cephalotes, Atta.

Cephus cinctus, on wheat in Canada, 176, 368, 600; thelyotokous parthenogenesis in, 176; danger of introduction of Agropyrum smithi into wheat areas on account of, 368.

Cephus pygmaeus (on cereals), in Czechoslovakia, 468; in Denmark, 117; in Russian Union, 680; cytology of strains of, in Europe

and Canada, 176.

Ceracris kiangsu, outbreaks and distribution of, in China, 451, 477; measures against, 477.

ceramicus, Xyleutes.

Ceraphron manilae, parasite of Fornicia in Java, 347.

cerasi, Myzus; Rhagoletis.

cerasi, auct., Caliroa (see C. lima-cina).

Ceratation (see Ation).

Ceratia coffeae, on cucurbits in Fiji, 641.

Ceratia quadrimaculata, on cucurbits in Fiji, 641.

Cevatia similis, repellent against, on cucurbits in Formosa, 618.

Ceratitis capitata, in S. Africa, 330; utilisation of Tetrastichus giffardi against, in Brazil, 34; in Cyprus, 270; bionomics of, in France, 195; in Italy, 98; in Queensland, 346; intercepted Russian Union, 439; not found in Trinidad, 24; legislation against importation of, U.S.A. from Hawaii, 344; climatic conditions unfavourable to, in Florida, 327; reported in Venezuela, 24; on Citrus, 270, 330, 439; on coffee, 34; other host-fruits of, 195, 439; experiment on oviposition of, 346; reproductive organs of, 512; measures against, 98, 270, 330, 344.

Ceratitis rosa, measures against, on orange in S. Africa, 330.

Ceratomegilla fuscilabris, predacious on Anuraphis roseus in Connecticut, 428.

Ceratomia catalpae, in U.S.A., 253, 421, 655; parasite and hyperparasite of, 253, 655, 656; toxicity of basic copper arsenate to, 421.

ceratoniae, Myelois.

Ceratostomella ulmi (see Ophiostoma).

Ceratovacuna lanigera (see Oregma). cercerisella, Gelechia.

Cercis canadensis, pests of, in Kansas, **376**.

Cerconota anonella, on Anona squamosa, etc., in Brazil, **566.** cerealella, Sitotroga.

cerealium, Limothrips.

Cereals, grasshopper damaging, in Argentina, 214; pests of, in Britain, 96, 332; pests of, in Germany, 352, 353, 519, 631, 632; locust damaging, in India, 291; grasshopper damaging, in Korea, 557; Laphygma exempta on, in Nyasaland, 126; pests of, in Poland, 403, 676; pests of, in Russian Union, 679, 680; Thysanoptera on, in Sweden, 121; Oscinella spp. on, in Switzerland, 270; pests of, in U.S.A., 32, 67, 284, 287; cutworms on, in Victoria, 389; type of injury to, by bugs, 352, 353, 548, 554; insects and other factors associated with white ear disease in, 121, 680; question of survival of grain mites in fields of, 312, 318. (See Maize, Wheat, etc.)

Cereals (Stored), pests of, and their control, 8, 24, 118–120, 137, 162, 172, 173, 196, 199, 220, 248, 265, 305, 312, 318, 319, 339, 341, 350, 355, 360, 378, 551, 552, 582, 585, 600, 667, 685; chloropicrin reducing germination of, 319.

Ceresa bubalus (on fruit trees), in France, 390, 689; in Jugoslavia, 474; in Switzerland, 169, 438; in U.S.A., 689; bionomics of, 169, 390, 689; measures against, 690.

Cerodontha denticornis (on cereals), in Japan, 208; in New Zealand, 547, 548.

Ceromasia sphenophori, parasite of Rhabdocnemis obscura in Hawaii, 90, 367.

cerophaga, Angitia. ceroplastae, Aneristus.

Ceroplastes, new Encyrtid parasites of, in Africa, 503; on coffee in Kenya, 503.

Ceroplastes africanus, new Encyrtid hyperparasite of, in Sudan, 40.

Ceroplastes itatiayensis, sp. n., in

Brazil, 48.

Ceroplastes sinensis, on Citrus in Russian Union, 150, 308; experiments with Cephalosporium lecanii against, 308; other measures against, 150.

Cerosipha forbesi, bionomics of, on strawberry in Ontario, 276.

Cerura bifida, on poplar and willow in Morocco, 196.

cerurae, Microplitis.

cervicalis, Zelus.

cervinus, Dascillus.

Ceuthorrhynchus macula-alba (Poppy Weevil), in Czechoslovakia, 468, 469, 470; bionomics of, 469; measures against, 470.

Ceuthorrhynchus pleurostigma, on cauliflower in France, 459.

Ceuthorrhynchus punctiger, bionomics and control of, on Taraxacum hok-saghyz in Kazakstan, 45.

Ceylon, leaf-curl and pests of Capsicum frutescens in, 493, 494; Nephantis serinopa on coconut in, 638; catalogue of Longicorns in forests in, 612; pests of stored cacao in, 638; Homona coffearia on tea in, 19; beneficial insects and biological control in, 19, 638; summary of plant quarantines in, 592; natural enemies of Pseudococcus introduced into Kenya from, 665.

Chaetexorista javana, parasite of Setora nitens in Java, 347.

Chaetocnema, on ramie in Transcaucasia, 587.

Chaetocnema aridula, on wheat in Central Asia, 453.

Chaetocnema concinna, on beet in Britain, 674.

Chaetocnema cylindrica, on rice in Japan, 556.

Chaetococcus bambusae, on bamboo in Porto Rico, 501.

Chaetodacus (see Dacus).

Chaetophorus hickelianae, in Italy, 512.

Chaetoptelius vestitus (see Hylesinus). chalcedonia, Elaphria.

Chalcidoids, list of, in Germany, 352.

chalcites, Plusia (Phytometra); Pterostichus (Poecilus).

chalcodermi, Urosigalphus.

Chalcodermus aeneus (Cowpea Curculio), bionomics of, in U.S.A., **76, 82, 644;** measures against, **82.**

Chalcodermus bondari, bionomics and control of, on cotton, etc., in Brazil and Argentina, 413, 414.

Chalcodermus camposi, sp. n., on

cacao in Brazil, **566.**

Chalcodermus marshalli, on cacao in Brazil, **566.**

Chalk, as carrier for dusts, 313, 367. Charaeas graminis, natural enemies of, on grasses in Scotland, 122.

Cheese Skipper (see *Piophila casei*).

Cheiloneurus African species of 40

Cheiloneurus, African species of, 40, 503.

Cheiloneurus angustifrons, sp. n., probably hyperparasite of Ceroplastes africanus in Sudan, 40.

Cheiloneurus carinatus, sp. n., hosts and distribution of, in Africa, **40.** Cheiloneurus chiaromontei, doubt-

fully distinct from C. cyanonotus, 40.

Cheiloneurus cyanonotus, distribution and hosts of, in Africa, 40; C. chiaromontei doubtfully distinct from, 40.

Cheiloneurus latiscapus, sp. n., reared from mealybugs parasitised by Anagyrus in Natal, 40. Cheiloneurus orbitalis, sp. n., hosts

of, in Natal, 40.

Cheimatobia brumata (see Operophtera).

Cheiranthus, Aleurodes brassicae on, in Britain, 61.

Cheiropachus, key to N. American species of, 434.

Chelonus, parasite of Pyrausta aurata meridionalis in Morocco, 193.

Chelonus annulipes, in Canada, 381; introduced into Porto Rico against Diatraea saccharalis, 382; hosts and utilisation of, in U.S.A., 80, 381, 382, 656; relation of host physiology to development of, 80.

Chelonus pectinophorae, introduced into U.S.A. from Korea against Platyedra gossypiella, 381.

Chermes abietis, measures against, on spruce in New York, 529; characters of, 91.

Chermes boerneri, recorded as C. pini on pines in Australia, 262, 546; restricted to Pinus radiata in U.S.A., 546; question of biological control of, 262, 546.

Chermes ishiharai, on Picea spp. in Japan, 91; characters of, 91.

Chermes japonicus, on Picea spp. in Japan and Sakhalin, 91; characters of, 91.

Chermes karafutonis, sp. n., on Picea jezoensis in Sakhalin, 91.

Chermes nordmannianae (on Abies), in Czechoslovakia, 469; in Germany, 48; increase of, in Switzerland, 517; measures against, 48, 517.

Chermes nüsslini (see C. nordmannianae).

Chermes piceae, on Abies grandis in U.S.A., 499.

Chermes pini (on pines), C. boerneri recorded as, in Australia, 262, 546; predacious enemies of, in Britain, 262, 263.

Chermes strobi, predacious enemies of, on pines in Britain, 262, 263.

Chermes strobilobius, on Larix decidua in Czechoslovakia, 469.

Chermes tsugae, on Tsuga heterophylla in U.S.A., 499.

Chermes viridis, on Larix decidua in Czechoslovakia, 469.

Cherry, pests of, in Australia, 20, 21, 279; Argyresthia ephippella on, in Austria, 232; Myzus cerasi on, in Britain, 333; Illiberis pruni on, in China, 668; Phalera flavescens on, in Formosa, 206; pests of, in France, 622, 689; pests of, in Germany, 114, 155, 156, 297, 513; Aphids on, in Hungary, 458; Malacosoma indica on, in India, 577; pests of, in Japan, 211, 338, 361, 620; Monosteira unicostata on, in Morocco, 359; pests of, in U.S.A., 429, 498, 535, 643; Coccids on, in Uzbekistan, 474; Rhagoletis cerasi not attracted by extracts of, 297; R. cerasi intercepted in fruit of, in Russian Union, 439.

Cherry Case-bearer (see Coleophora pruniella).

Cherry Fruit-fly (see Rhagoletis cerasi).

chersis, Sphinx.

Chervil, Curled (see Anthriscus cerefolium).

Chestnut, pests of, in France, 358; pests of, in Greece, 452; pests of, in Italy, 358; Lachnus tropicalis on, in Japan, 615; Dryocoetes minor on, in Turkey, 522; treatment of nuts of, against pests, 358, 359.

Cheyletus eruditus, predacious on Tyroglyphus farinae in Russian Union, 319.

chiaromontei, Cheiloneurus.

Chickweed (see Stellaria media).
Chile, sawfly introduced into New
Zealand from, against Acaena,

449.

Chillies (see Capsicum).

Chilo infuscatellus, on sugar-cane in Formosa, 206, 339, 363, 447; dispersal of larvae of, 339; relation of growth of sugar-cane to oviposition by, 206; natural enemies and biological control of,

363, 364, 447.

Chilo simplex (on rice), in China, 613; in Hawaii, 367; in Japan, 209, 338, 441, 442, 619, 620; in Manchuria, 340; other foodplants of, 340; parasites and biological control of, 367, 442, 619, 620; flooding reducing infestation of rice by, 209; measures against, 209, 441, 613; tests of insecticides on, 18, 338, 620.

Chilo trypetes, sp. n., on sugar-cane

in Punjab, 456.

Chilo zonellus, effect of, on sugarcane varieties in India, 558.

Chilocorus, predacious on Coccids in

Dahomey, 349.

Chilocorus bipustulatus (predacious on Coccids), in Cyprus, 270; in Russian Union, 304, 316; bionomics and utilisation of, 304.

Chilocorus distigma, establishment of, against Coccids in Seychelles,

335.

Chilocorus politus, utilisation of, against Coccids in Mauritius, 611.

Chilocorus renipustulatus, predacious on Aspidiotus perniciosus in Russian Union, 316.

Chilocorus renipustulatus ab. inornatus, introduced into Caucasus from Russian Far East against Aspidiotus perniciosus, 303; bionomics of, 303, 304.

Chilocorus schioedtei, predacious on Coccids in Dahomey, 349.

Chilocorus wahlbergi, establishment of, against Coccids in Seychelles, 335.

chilonis, Amysoma (see Microbracon chinensis).

China, Buzura suppressaria on Aleurites in, 614; forest pests in, 360, 449, 451; Diacrisia obliqua on lima beans in, 129; miscellaneous pests in, 450, 451;

orchard pests in, 450, 667-669; rice pests in, 451, 613; new Aphids in, 336; new Coccid in, 48; locusts and grasshoppers in, 450, 451, 477, 486; beneficial insects in, 669; Eupristina verticillata introduced into Hawaii from, to pollinate Ficus retusa, 367; Coccid on orchids imported into Japan from, 617.

Chinch Bug (see Blissus leucopterus).

chinensis, Aspidiotus; Bruchus
(Callosobruchus); Melaphis;

Microbracon; Oxya.

Chinese Banyan (see Ficus retusa). Chinese Oak Silkworm (see Antheraea pernyi).

Chionaspis asiatica, on fruit trees

in Uzbekistan, 474.

Chionaspis pinifoliae, measures against, on pines in New York, 529, 530.

Chionaspis tegalensis, Coccinellid predacious on, on sugar-cane in Mauritius, 611.

Chirida signifera, measures against, on sweet potato in Jamaica, 239.

Chirothrips hamatus, on Alopecurus in Finland, 160; on wheat in Sweden, 121.

Chirothrips manicatus, on wheat in

Sweden, 121.

chittendeni, Dialeurodes.

Chlorida festiva, on egg-plant in Jamaica, 239.

Chloridea (see Heliothis).

Chloris, relation of Nomadacris septemfasciata to, in Tanganyika, 387, 388.

chloris, Aethemenes (Nezara); Aphis.

Chlorita (see Empoasca).

Chloroacetamide, tests of, in baits

for grasshoppers, 537.

Chlorochroa sayi, and boll rots in cotton in Arizona, 416, 417; measures against, 417.

Chlorocresylic Acid, as repellent for *Hylemyia brassicae*, **116**.

Chlorocryptus, parasite of Setora

nitens in Java, 347.
Chloropicrin, ineffective against ants, 44; as a soil fumigant, 44, 617, 622; against pests of stored products, 225, 319, 338, 349, 378, 451; causing injury to Citrus, 617; reducing germination of grain, 319; azaleas killed by emulsion of, 605; effect of addition of, to insecticidal dusts, 470, 471.

Chloropisca notata, bionomics and control of, on cereals in Japan, 556.

Chlorops circumdata (see Chloro-

pisca notata).

Chlorops oryzae, on rice in Japan, 207, 555, 620; bionomics of, 207; measures against, 555.

Chlorops taeniopus (on cereals), in Britain, 332, 562; in Czechoslovakia, 468; parasites of, 562.

choaspes, Microphanurus.

Chondacris rosea, measures against, in China, 477.

Chopardina importata (see Dolichopoda bormansi).

Chorthippus albomarginatus, in Russian Union, 320, 386; ecology of, 386.

Chorthippus apricarius, in Poland, 9; ecology of, in Russia, 386.
Chorthippus biguttulus, ecology of,

in Russia, 386.

Chortoicetes terminifera, in Australia, 56, 130, 345, 476, 545, 602; bionomics of, 345, 545; Trichopsidea oestracea parasitic on, 56; measures against, 130, 476, 602.

Chortophila, doubtful identity of, 113, 114. (See Hylemyia.)

chosensis, Hemiteles.

Chromaphis nirecola, on elms in Japan, 443.

chromataria, Earias.

Chromatomyia ilicis (see Phytomyza). Chromic Oxide, tests of, in baits for grasshoppers, 134.

Chromocryptus antipodialis, parasite of Cydia molesta in Australia, 640; parasitised by Dibrachys

cavus, 640.

Chrysanthemum, pests of, in Britain, 333, 519, 673; restrictions on importation of, into Britain, 356; Ceresa bubalus on, in France, 689; pests of, in U.S.A., 141, 142, 673; identity and distribution of gall-midges on species of, 673. (See Pyrethrum Plants.)

Chrysanthemum Gall-midge (see

Diarthronomyia).

Chrysobothris femorata, on apple, etc., in U.S.A., 178, 420.

Chrysobothris igniventris, bionomics of, on pines in Turkey, 219.

chrysocephala, Psylliodes.

Chrysocharis gemma, parasite of Phytomyza ilicis in Britain, 669. Chrysocharis syma, parasite of

Phytomyza ilicis in Britain, 669; parasite of, 670.

Chrysochroa, infesting Finnish plywood in India, 465.

Chrysocoris javanus (atricapillus), new parasite of, in Java, 19.

Chrysolina (see Chrysomela).

Chrysomela brunsvicensis, factors preventing establishment of, against Hypericum in Australia, 282.

Chrysomela fastuosa, tests of action of HCN on, 309.

Chrysomela geminata, introduced into Australia against Hypericum, 546.

Chrysomela hyperici, factors preventing establishment of, against Hypericum in Australia, 282.

Chrysomela tremulae (see Melasoma).

Chrysomela varians, factors preventing establishment of, against Hypericum in Australia, 282.

chrysomphali, Aphytis.

Chrysomphalus aonidum, auct. (see C. ficus).

Chrysomphalus aurantii (see Aonidiella).

Chrysomphalus dictyospermi, measures against, on Citrus in Argentina and Transcaucasia, 23, 150.

Chrysomphalus ficus (on Citrus), in Argentina, 23; in Egypt, 509; in Florida, 256, 257, 259; on Elaeagnus pungens in Japan, 91; on coconut in Philippines, 451; other food-plants of, 259, 509; peaks of reproductive activity of, 256; measures against, 23, 256, 509; effect of spray residues on populations of, 257; tests of thiocyanate and oil against, 259.

Chrysomphalus obscurus, measures against, on pecan in Louisiana, 255.

Chrysomphalus personatus, measures against, on Ficus in Egypt, 509; on banana in Haiti, 289.

Chrysomphalus pinnulifer var. diversicolor, on orange in S. Rhodesia, 15.

Chrysomphalus rossi, measures against, on Citrus in New Zealand, 523, 524.

Chrysopa, predacious on Erannis spp. in Poland, 438; predacious on other insects in Russian Union, 215, 453; suitability of, as host for Trichogramma evanescens, 36.

Chrysopa oculata separata (predacious on Macrosiphum onobrychis), bionomics of, in Utah, 30. Chrysophyllum cainito, Dacus passiflorae on, in Fiji, 366.

chrysorrhoea, L., Arctornis.

chrysorrhoea, auct., Euproctis (Liparis) (see Nygmia phaeorrhoea).

Cicadella viridis, on beet in China, 451; measures against, on fruit trees, etc., in Germany, 398; doubtful vector of disease of sugar sorghum in Italy, 472.

Cicadula sexnotata (see Macrosteles). Cicer arietinum, Heliothis armigera

on, in S. Rhodesia, 492.

Cidaria bilineata, suitability of, as host for Trichogramma evanescens, 35.

Cidaria fluctuata, on cauliflower in Denmark, 117.

Cider, affected by arsenic spray residues on apples, 273.

Cienfuegosia hildebrandti, Diparopsis castanea on, in S. Africa, 488, 489; not a food-plant of Dysdercus fasciatus, 489.

Cigarette Beetle (see Lasioderma serricorne).

ciliatus, Dacus.

cilicrura, Hylemyia (Chortophila). Cimex, suitability of, as host for Trichogramma evanescens, 36.

Cinara hattorii, sp. n., on Abies sachalinensis in Japan, 339. cincticeps, Nephotettix bipunctatus.

cinctipes, Lasiosina. cinctus, Cephus.

cinerascens, Cnephalia (Gonia).

cinerea, Apriona; Zeuxia.

cingulata, Herse. cingulatus, Dysdercus.

Cinnabar Moth (see Tyria jacobaeae).

cinnamomeus, Aradus.

Cinnamomum camphora, new Eriophyid on, in California, 288. Cinnamomum glanduliferum, Xylo-

copa latipes in timber of, in India, 337.

Cinnamon Oil, tests of, in baits for Dacus oleae, 98.

circumdata, Chlorops (see Chloropisca notata).

circumflexus, Myzus.

Cirphis latiuscula, parasites of, on sugar-cane in Florida, 657.

Cirphis loreyi, parasites of, on sugarcane in Queensland, 186.

Cirphis unipuncta, in Australia, 57, 294, 389; in Fiji, 213, 641; in Japan, 441; in Manchuria, 361; in U.S.A., 67, 599; on cereals and grasses, 67, 213, 361, 389, 599; on cotton and lucerne, 599; Tachinid parasites of, 57, 294;

probably destroyed by Bufo marinus, 641; measures against, 213, 390; tests of oil baits for. 67.

čisti, Bruchus (Spermophagus).

cistipennis, Fundella.

Cistogaster globosa, parasite Aelia acuminata in Germany, 64. citrana, Eulia (Argyrotaenia).

citri, Dialeurodes; Paratetranychus; Prays; Prontaspis; Pseudococcus; Scirtothrips.

citricola, Coccidophilus.

Citricola Scale (see Coccus pseudomagnoliarum).

citriculus, Pseudococcus.

citrifolii, Dialeurodes.

citrina, Aonidiella; Gyranusa.

citrinus, Aspidiotiphagus.

Citrus, locusts damaging, in Fr. W. Africa, 571; pests of, in S. Africa, 323, 330; pests of, in Argentina, 23, 238; pests of, in Australia, 89, 278, 292, 330, 565, 603; pests of, in Cyprus, 270; Coccids on, in Egypt, 507, 508, 509; Mictis profana on, in Fiji, 641; Pulvinaria polygonata on, in Formosa, 557; Pseudococcus citri on, in France, 266; pests of, in India, 128, 503; pests of, in Japan, 207, 440, 441, 616, 617; pests of, in New Zealand, 21, 56, 277, 278, 523, 524; Coccids on, in Peru, 165, 511; pests of, in Philippines, 452; pests of, in S. Rhodesia, 14, 15, 16; Coccids on, in Russian Union, 148, 149, 150, 308, 454; Dacus on, in Samoa, 365; adaptation of Geometrid to, in Sicily, 630; fruit-piercing moths damaging, in Sierra Leone, 271; Aphids and Coccids on, in Spain, 6; pests of, in U.S.A., 55, 256, 257, 258, 259, 261, 342, 343, 377, 424, 425, 432-434, 459-461, 642, 649, 650, 653; pests of, in W. Indies, 24, 25, 181, 239; illustrations of pests of, 617; effect of spray residues on populations of Coccids on, 257; insects intercepted on fruit and seedlings of, in Japan and Russia, 50, 439, 616; insecticides causing injury to, 15, 150, 425, 433, 460, 508, 649, 650, 653; tests of reactions of sprays used on, 278; effects of lead arsenate and copper carbonate on fruit of, 330, 331.

Citrus Aphis (see Aphis spiraecola). Citrus Borer (see Oemona hirta).

Citrus Gall Wasp (see Eurytoma fellis).

Citrus Red Mite (see Paratetranychus citri).

Citrus Red Scale (see Aonidiella aurantii).

Citrus Thrips (see Scirtothrips citri). Citrus Whitefly (see Dialeurodes

citri). infesting Cladosporium, Astero-

lecanium bambusae in Porto Rico.

claripalpis, Theresia (Paratheresia). claripennis, Coccidoxenus.

Clastocnemis quadrimaculata (see Platygenia).

clavatus, Prochiloneurus.

Clays, as carriers for dusts, 32, 150, 177, 249, 250, 310, 371, 380 spray formula containing, 532; mercury compounds mixed with, for treating seed against Psila rosae, 497.

cleanthes, Rhaconotus.

Clensel, in baits for Trypetids, 98, 270.

Cleonus mendicus, survey of data on, on beet in Italy, 301.

Cleonus piger, bionomics and control of, on artichoke in Italy, 473.

Cleonus punctiventris (on beet), in Czechoslovakia, 470; in Russia, 306, 407, 678, 684; birds destroying, 470; emergence and migration of, 678; experiments with fungi against, 306, 307, 470, 684; other measures against, 407, 470, 678.

Clidemia, attempted introduction of Liothrips urichi into Solomon Is. against, 366.

Clinodiplosis equestris (see Haplodiplosis).

Clisiocampa (see Malacosoma).

clisiocampae, Ablerus; Dibrachys (see D. cavus); Oöencyrtus; Telenomus.

Clistomorpha brasiliensis, parasite of Dysdercus in Brazil, 22.

Closterocerus trifasciatus, parasite of Phytomyza ilicis in Britain, 669.

Clothes Moths, tests of mothproofing preparations and fumigants against, 140, 635.

Clove Tree, pests of, in Nether-

lands Indies, 227.

Clover, pests of, in W. Australia, 131, 292; Apion apricans on, in Czechoslovakia, 470; pests of, in Germany, **519**, **686**; Collembola on, in New Zealand, **550**, 551; Estigmene acraea on, in

Oklahoma, 341; pests of, in Russian Union, 322, 568, 679, 681, 682; Apion spp. on, in Sweden, 117; susceptibility of varieties of, to Apion spp., 118; Aphis fabae transmitting virus disease of, 686.

Clover, Alsike (see Trifolium

hybridum).

Clover, Subterranean (see Trifolium subterraneum).

Clover Weevils (see Apion aestivum, A. apricans and A. flavipes).

clypeata, Lachnosterna (Phyllo-

phaga).

Clysia ambiguella (Vine Moth), in France, 155; in Germany, 102, 111, 458, 465, 496, 582, 689; in Russian Union, 403; bionomics of, 111, 466, 496; breeding of, for testing insecticides, 111; timing of sprays against, 155; sprays against pupae of, 103, 689; baits for, 465, 496, 582; banding against, 458; use of females for trapping males of, 466, 582.

Cnaphalocrocis medinalis, on rice and grasses in Philippines, 364,

451.

Cnaphalodes (see Chermes).

Cneorrhinus plagiatus, on beet and crucifers in Denmark, 117.

Cnephalia cinerascens, parasite of Spodoptera mauritia in Formosa, 363.

Cnephasia longana, on strawberry in Oregon, 498.

Cnidocampa flavescens (see Mon-

Coal, Brown, compared with bentonites for use in nicotine sprays, 545.

Coal-tar Creosote (see Tar Distillates).

Cobana Negra (see Stahlia).

Coca (see Erythroxylon coca).

coccidivorus, Anagyrus.
Coccidophilus citricola, predacious
on Protargionia larreae in

Argentina, 24. Coccidoxenus claripennis, sp. n., parasite of Inglisia conchiformis in Uganda, 40.

coccinellae, Perilitus; Tetrastichus. Coccinia indica, Dacus spp. on, in Madras, 561, 562.

Coccoloba ilheense, new Lamiid on, in Brazil, 290.

Coccomytilus zlocistii (see Lepido-saphes).

Coccophagoides similis, parasite of Targionia vitis in Italy, 471.

Coccophagus gurneyi, causes of immunity of certain mealybugs from, 598.

Coccophagus lycimnia (lecanii), parasite of Aonidiella citrina in Transcaucasia, 149.

Cocculus trilobus, Calpe emarginata on, in Japan, 209.

Coccus hesperidum, in Argentina, 23; in Natal, 503; in S. Rhodesia, 15; parasites of, 15, 503; measures against, on Citrus, 15, 23.

Coccus longulus, measures against,

on Anona in Egypt, 509.

Coccus pseudomagnoliarum, measures against, on Citrus in California, 343, 433, 649; measures against, 433, 649; resistance of strains of, to HCN, 343.

Coccus viridis, on limes in Grenada, 25; Scoliid feeding on honeydew of, on guava in Haiti, 84.

Cochineal Insect (see *Dactylopius*). cochleariae, *Phaedon*. cockerelli, *Paratrioza*.

Cockroaches, destroyed by *Bufo marinus* in Fiji, **641**; sodium fluoride dust against, in Hawaii, **367**; toxicity of *Bacillus cereus* to, **81**; temperature and humidity relations of, **224**.

Coconut, Rhynchophorus palmarum on, in Brazil, 22; Nephantis serinopa on, in Ceylon, 638; pests of, in Fiji, 22, 212, 213, 366; Brassolis sophorae on, in Br. Guiana, 524; pests of, in Hawaii, 367; new thrips intercepted on, in Japan, 50; pests of, in Mauritius, 611; pests of, in New Guinea Territory, 22, 277, 294, 366; Promecotheca opacicollis on, in New Hebrides, 366; pests of, in Philippines, 451; Oryctes rhinoceros on, in Samoa, 641; pests of, in Seychelles, 335; P. reichei on, in Tonga, 366; pests of, in W. Indies, 25, 239; Xyleborus perforans in timber of, 181.

Coconut Caterpillar (see Nephantis serinopa).

Coconut Leaf-roller (see Lampro-sema blackburni).

Coconut Oil (see Oil, Coconut).

Coconut-oil Soap (see Soap, Coconut-oil).

Cocos coronata, pests of, in Brazil, 661.

Cocos nucifera (see Coconut).

Cocos vagans, pests of, in Brazil,

Codiohispa anonicola, on Anonaceae

in Brazil, 567.

Codling Moth (see Cydia pomonella). Coelinius niger, parasite of Chloropids in Britain, 562.

Coeloides, parasite of Pissodes spp.

in Finland, 8.

Coeloides scolyticida, parasite of Scolytids in Holland, 397.

Coeloides stigmaticus, parasite of Pissodes pini in Finland, 8.

Coelonotus, considered a synonym of Aphidius, 615.

Coelonotus japonicus, parasite of Lachnus tropicalis in Japan, 615.

Coelophora inaequalis, introduced into Porto Rico against Sipha flava, 382.

Coelopisthia nematicida (see Trit-

neptis klugi).

Coelopisthia scutellata (see Tritneptis).

coenobita, Panthea.

coerulea, Phytoecia; Zicrona.

coerulescens, Oedipoda; Tetrastichus. coeruleus, Corynetes; Rhynchites. coffeae, Ceratia (Aulacophora); Diarthrothrips; Lygus; Saissetia; Schizobremia; Zeuzera. coffearia, Homona.

Coffee, Stephanoderes hampei not found on, in Italian E. Africa, 175; pests of, in Brazil, 34, 188, 238; Zeuzera coffeae on, in India, 577; pests of, in Netherlands Indies, 161, 522; Xylotrechus quadripes on, in Indo-China, 49; pests of, in Kenya, 40, 503, 665, 666, 667; Anthores leuconotus on, in Nyasaland, 126; pests of, in Sierra Leone, 271; pests of, in Tanganyika, 40, 126; S. hampei on, in Uganda, 188; Leucoptera coffeella on, in W. Indies, 435; relation of Xyleborus and pruning to crop yield of, 522; Aleurites montana a suitable shade-tree for, 49; Rhynchota associated with Nematospora in, 389.

Coffee (Stored), Tribolium spp. imported into Britain in, 666; treatment of, against Stephanoderes hampei, 667.

Coffee Berry Borer (see Stephanoderes hampei).

coffeella, Leucoptera. cognatus, Poeciloscytus. coimbatorensis, Dinarmus. Colabhellus bowringi. against, on vegetables in China, 450.

743

Colaphellus hoefti, on crucifers in Russian Union, 439.

Colaspis hypochlora, on bananas in Br. Guiana, 524.

Colaspis trivialis, measures against, on vines in Brazil, 34.

Cold Storage, uses of, against insect pests, 120, 173, 344, 360, 361, 513; Aphids kept in, for rearing Coccinellids, 81; effect of, on reproduction of parasitic Hymenoptera, 144.

Cold-hardiness, studies on, in insects, 37, 298, 404, 405, 406.

Coleophora fuscedinella, new parasite of, on elm and alder in Britain, 58.

Coleophora hemerobiella, on plum and apple in Czechoslovakia, 468.

Coleophora pruniella, parasites of, on apple and cherry in Oregon,

coleophorae, Apanteles.

Coleoptera, lists of: associated with cultivated fruits in Britain, 160: in potato storage cellars Finland, 392.

Coleus parviflora, pests of, Madras, 129.

colibri, Athalia (see A. rosae). Colletotrichum falcatum, 656.

collinus, Laevicephalus (Deltocephalus).

Collops quadrimaculatus, predacious on Blissus leucopterus in Oklahoma, 261.

Collyria calcitrator, introduced into U.S.A. from Canada against Trachelus tabidus, 381.

Coloborhombus fasciatipennis, on clove trees in Netherlands Indies, 227.

Colocasia, Papuana huebneri on, in Gilbert Is., 366.

Colombia, Monalonion on cacao in, 566; summary of plant quarantines in, 274.

Colorado, Aphids in, 104, 542, 596; breeding areas of Eutettix tenellus in, 593, 594; forest pests in, 377; 528; Paratrioza spp. on tomato and potato in, 72, 408, 422, 423; beneficial insects in, 528.

Colorado Potato Beetle (see Leptinotarsa decemlineata).

Colours, reactions of insects to, 7, 31, 442, 446.

columbariella, Tinea.

communis, Lygus; Melanotus; Panorpa.

comosa, Eucelatoria. complanella, Tischeria.

compressiventris, Apanteles.

Compsilura concinnata, parasite of Vanessa cardui in Transcaucasia, 587; establishment of, against Stilpnotia salicis in Washington,

499. Compsilura concinnata sumatrensis, parasite of Cirphis loveyi in Queensland, 186.

comptana, Ancylis.

Comstockiella perniciosa (see Aspidiotus).

conchiformis, Inglisia.

concinna, Chaetocnema.

concinnata, Compsilura. concinnus, Wesmaëlius.

confertus, Pityophthorus.

confinis, Pityophthorus.

confraternus, Pipunculus.

confusum, Tribolium. confusus, Dactylopius.

Congo, Belgian, Crocidolomia binotalis on cabbage, etc., in, 572; Aleurodids transmitting mosaic of cassava in, 153; Platyedra gossypiella on cotton in, 572; Tinea vastella damaging horns in, 294; locusts in, 476, 482, 483, 486; new parasite of Pseudococcus in, 503; cultivation of insecticidal plants in, 294; parasites imported into Kenya from, against P. kenyae, 665. congregatus, Apanteles.

Conium maculatum, new Aphid on,

in California, 590.

Connecticut, forest pests in, 73, 77, 176; maize pests in, 179, 430, 456; Thrips tabaci on onions in, 71; orchard pests in, 428, 429; pests of tobacco and potato in, 177, 178; Calomycterus setarius in, 600; beneficial insects and biological control in, 73, 74, 176, 177, 178, 428, 429; regulations concerning transportation of nursery stock to other States and Canada from, 176.

Conomorium eremita, parasite of Erannis in Crete, 172.

Conopia exitiosa (see Aegeria).

Conorrhynchus (see Cleonus).

Conotrachelus bondari, on Anonaceae in Brazil, **567.**

Conotrachelus nenuphar (on apple), in Canada, 382, 664; bionomics of, in U.S.A., 382; measures against, 383, 664.

constrictus, Barbitistes.

Contarinia fagi, bionomics and control of, on beech in Germany, 353.

Contarinia humuli, on hops in Britain, 333.

Contarinia mali, sp. n., measures against, on apple in Japan, 556, 557.

medicaginis, factors Contarinia affecting abundance of, lucerne in Germany, 457.

Contarinia nasturtii (on crucifers), Czechoslovakia, 468; in France, 167; bionomics of, in Holland, 514; types of injury to cauliflower by, 167; measures against, 167, 514.

Contarinia pisi, on leguminous crops in Britain, 332.

Contarinia pyrivora, characters distinguishing C. mali from, 556.

Contarinia torquens (see C. nasturtii).

Contarinia tritici, in Holland, 268; in Italy, 3; bionomics and control of, on wheat and Agropyrum, 3, 268.

contraria, Ochrogaster. convergens, Hippodamia.

convexifyons, Dendroctonus.

convexus, Campoplex.

convolvuli, Myzus (see Macrosiphum solani).

Convolvulus, insects attacking, 322, 375, 677.

Cook Islands, insects introduced into, against fruit-flies and Cosmopolites, 212, 365.

cooki, Cremastus.

Copaifera coleosperma, Bostrychids in timber of, in S. Africa, 60.

cophta, Gryllotalpa gryllotalpa.

Copidosoma, parasite of Phthorimaea operculella in Brazil, 289.

Copper Arsenate, properties and action on insects of basic form of, 421, 422.

Copper Carbonate, in sprays against Trypetids and Lepidoptera on Citrus, **330, 331.**

Copper Sulphate, use of, for destroying Oryctes, 347; in dusts against potato flea-beetles, 342; use of, against Rhizopertha in pith helmets, 612. (See Bordeaux Mixture.)

Copper Sulphate, Basic, as a corrective for foliage injury by calcium arsenate, 373; basic copper

arsenate compared with, as a fungicide, 422; not highly toxic to Cylas formicarius, 373.

Copra, insects infesting, 212, 213,

584.

coprosmae, Aphis.

Coptocercus biguttatus, possibly on clove trees in Netherlands Indies, 227.

Coptotermes (in building timber), in Central America, 56; Australia, 545; in Barbados, 56, 384.

Coptotermes acinaciformis, in Australia, 545; in Fiji, 212, 641; damaging plants, 546; damaging timbers, 212, 545, 641.

Coptotermes formosanus, synonymy

and distribution of, 384. Coptotermes frenchi, damaging trees

in New South Wales, 546. Coptotermes javanicus, in Jamaica and Java, 384.

Coptotermes raffrayi, in Australia,

Coptotermes remotus, Hill, 384.

Coptotermes remotus, Silv. (see C. formosanus).

Coptotermes testaceus, damaging furniture and building timber in W. Indies, **55, 56, 384**; characters of, 55; measures against, 55.

corallina, Mecistomela (Alurnus).

corbetti, Craniotectus.

Corchorus, Helopeltis bergrothi on, in Sudan, **574.**

Corcyra cephalonica, in stored cacao in Ceylon, 638.

Cordia myxa, Galerucid on, in Burma, **693.**

Cordyceps barberi, infesting Diatraea saccharalis in Antigua, 660.

Corfu, Apiomyia bergenstammi in,

Coriandrum sativum, Aphids and virus disease of, in California, 590. Corizus hyoscyami (see Therapha). Corks, Borkhausenia pseudospretella damaging, in Scotland, 122.

Corn Borer, European (see Pyrausta

nubilalis). Corn Ear Worm (see Heliothis

armigera). Corn Root Worm (see Crambus caliginosellus).

corni, Lecanium (Eulecanium).

Cornus, cankerworms on, in New Tersey, 74.

cornutus, Bostrychoplites; Phyllocoptes.

Corrosive Sublimate (see Mercury Bichloride).

Corsica, new Coccids in, 48, 235; miscellaneous pests in, 459, 584; Dolichopoda bormansi in caves in, **604.**

corticinella, Setomorpha (see S. rutella).

coruscus, Phalacrus (see P. fimetarius).

coryli, Lecanium.

Corylus avellana (see Hazel).

Corymbites aeneus, feeding habits of, in Germany, 398; on cereals and cabbage in Siberia, 679; ecology of, 679.

Corymbites aeripennis destructor, bionomics of, in Alberta, 539; on potatoes in N. Dakota, 71.

Corymbites latus, on cereals and lupins in Russian Union, 679, 683; ecology of, 679; measures against, 683.

Corynetes coeruleus, in France, 584: predacious on other insects, 584. Corynetes pusillus, in France and

Corsica, 584.

Corythuca gossypii, food-plants of, in Jamaica, 239.

Coscineuta virens, measures against, on cassava in Trinidad, 25.

Cosmolyce boetica, Microbracon greeni ovipositing on, 129.

Cosmophila, measures against, on cotton in Philippines, 184.

Cosmophila flava, new Tachinid parasites of, in Queensland, 57; on cotton in Siam, 50.

cosmophilae, Zenillia. cosmopolitana, Amphorophora (see

A. lactucae).

Cosmopolites sordidus (on banana), in Ćook Islands, 212; in Fiji, 212, 365; in Formosa, 205; in Jamaica, 212, 238; in Java, 205; in Mysore, 291; in New South Wales, 184, 346; Tahiti, 212, 365; experi experiment on migration of, in banana plantations, 346; effect of cultivation on susceptibility of bananas to, 346; utilisation of Plaesius javanus against, 205, 212, 238, 365; other measures against, 184, 291, 346.

costalis, Eutermes (Nasutitermes);

Psylla.

Cotinis longitarsis, in nests of Atta texana in Texas, 288.

Cotton (Gossypium), pests of, in S. Africa, 223, 281, 487, 488, 489, 490, 491; Chalcodermus bondari on, in Argentina, 413; Platyedra gossypiella on, in Belgian Congo,

572; pests of, in Brazil, 22, 34, 86, 136, 413; Thrips tabaci on, in greenhouse in Britain, 672; pests of, in Cyprus, 270; Dysdercus mimus on, in Ecuador, 164; locusts damaging, in Greece, 478; pests of, in India, 127, 128, 228, 560, 575, 576; pests of, in Japan, 225, 555, 557; pests of, in Java, 280; pests of, in Kenya, 666; pests of, in Manchuria, 340; pests of, in Manchuria, 125, 126, 222; pests of, in Peru, 164, 385; pests of, in Philippines, 183, 184; pests of, in Queensland, 280, 281, 603; pests of, in N. Rhodesia, 222, 492; pests of, in S. Rhodesia, 222, 491, 492: pests of, in Russian Union, 43, 147, 150, 215, 304, 313, 314, 552, 609, 625; pests of, in Siam, 49; P. gossypiella on, in Sicily, 174; pests of, in Sudan, 572, 573, 574; pests of, in Tanganyika, 222, 492; pests of, in U.S.A., 82, 165, 242, 243, 244, 245, 246, 247, 340, 341, 372, 380, 381, 413, 416, 417, 500, 599; pests of, in West Indies, 220, 221, 222, 223, 271, 503, 660, 661; legislation 271, 503, 660, 661; legislation against introduction of Anthonomus grandis into Barbados with, 661; susceptibility of varieties of, to pests, 49, 221, 222, 247, 281, 381, 417, 491; type of injury to, by Jassids, 281; effect of thrips on yield of, 413; insects and boll rots in, 34, 222, 290, 360, 389, 416, 417, 491; insects associated with "crazy top" in, 574; Lygus lucorum possibly transmitting virus disease in, 225, 555; general discussion of diseases and pests of, 336; cultural measures against pests of, 221, 222; effect of selenised, on pests, 220, 221, 222; hairiness of leaves of, in relation to Aphis gossypii, 243; calcium arsenate causing injury to, 247; (Baled), fumigation of, against Platyedra gossypiella, 53-55.

Cotton Aphis (see Aphis gossypii). Cotton Boll Weevil (see Anthonomus grandis).

Cotton Bollworm (see Heliothis armigera).

Cotton Bollworm, Pink (see *Platy-edra gossypiella*).

Cotton Bollworm, Red or Sudan (see *Diparopsis castanea*).

Cotton Flea-hopper (see *Psallus* seriatus).

INDEX.

Cotton Leaf Worm (see Alabama argillacea).

Cotton Pruner (see Chalcodermus bondari).

Cotton Stainers (see Dysdercus).

Cottonseed, pests in, and their control, 165, 174, 576; X-rays for detecting Platyedra gossypiella in, 147; Tribolium castaneum infesting insulation material containing, 640; (meal), in baits for cutworms, 287.

Cottony Cushion Scale (see *Icerya* purchasi).

Cotyledon, Aphis on, in Australia, 448.

Couch Grass (see Agropyrum repens). Cowpeas (Vigna), pests of, in Japan, 361, 556, 617; Elasmopalpus lignosellus on, in Porto Rico, 501; Ootheca mutabilis on, in S. Rhodesia, 125; pests of, in U.S.A., 76, 82, 370, 500, 644; susceptibility of varieties of, to Chalcodermus aeneus, 82; insecticides causing injury to, 82.

Crambus caliginosellus, measures against, on tobacco in Tennessee, 380, 652.

cramerella, Acrocercops.

Craniotectus corbetti, on Derris in Malaya, 539.

crassipes, Apiomerus.

on, in U.S.A., 542.

crassissima, Lachnosterna (Phyllophaga).

crātaegi, Aporia; Thomasiniana. Crataegus (Hawthorn), pests of, in Australia, 21, 292; new Thomasiniana on, in Britain, 673; pests of, in Germany, 114, 155; Monosteira unicostata on, in Morocco, 359; Rhopalosiphum prunifoliae

Crataegus azarolus, host-fruit of Ceratitis capitata in France, 195.

Cratosomus bombina, on Anonaceae in Brazil, 567.

Cratosomus punctulatus, on orange in Trinidad, 24.

Cremastus biguttulus (see C. flavo-orbitalis).

Cremastus cooki, parasite of Ancylis comptana in Kansas, 376.

Cremastus flaviventris, sp. n. (parasite of Cydia molesta), bionomics of, in Argentina, 237; technique for breeding, 238.

Cremastus flavoorbitalis, 237; utilisation of, against Lamprosema blackburni in Hawaii, 367; parasite of Chilo simplex in Japan, 442; reactions of, to coloured lights, 442.

Cremasius interruptor, establishment of, against Rhyacionia buoliana in Connecticut, 74.

Cremastus rubeo, sp. n. (parasite of Cydia molesta), bionomics of, in Argentina, 237; technique for breeding, 238.

Crematogaster victima missuriensis, associated with Aphis pawneepae

in Kansas, 376.

Creontiades femoralis, effect of, on

cotton in Arizona, 417.

Creosote, dust containing, against Jassids on cotton, 281; methods of treating timber with, against pests, 181, 337; other uses of, against insects, 366, 420.

Cresol, in spray for destroying potato plants and Aphids, **333**. Crete, Geometrid and its parasites

on olives in, 171, 172.

cretica, Sesamia.

cribrata, Pycnarmon.

Cricket-bat Willow (see Salix alba var. coerulea).

Cricula trifenestrata, on Altingia excelsa in Java, 227.

crinita, Sitona. crinitus, Polistes.

Crioceris asparagi (Asparagus Beetle), measures against, in Utah, 426; bionomics of, in Italy, 471; parasites of, 194, 195, 472.

cristatus, Parabatus; Phloeosinus.

crithmi, Margarodes buxtoni.

Crithmum maritimum, Coccid on, in France, 235.

critolaus, Spathius.

Crocidolomia binotalis (on crucifers), in Belgian Congo, **572**; in Fiji, **641**; in Seychelles, **335**; Polistes predacious on, **336**.

Crossocosmia sericariae (see Stur-

mia).

Crossotus albopunctatus, on cotton in Kenya, 666.

Crotalaria, Anomala expansa on, in Formosa, 363.

Crotalaria juncea, Exora spp. on, in S. Rhodesia, 125.

Croton, Psallus seriatus on, in U.S.A., 245, 538.

Croton glandulosus, Chalcodermus bondari on, in Brazil, 414.

Crows, destroying Cleonus punctiventris in Czechoslovakia, 470. cruciata, Austroicetes.

cruciferae, Phyllotreta.

cruciferarum, Plutella (see P. maculipennis).

crucigerus, Dociostaurus.

cruda, Monima.

Cryolite, in baits, 287, 379, 537, 595; dusting with, 244, 246, 258, 286, 370, 371, 372, 373, 378, 379, 431, 461, 525, 595; carriers for, 258, 286, 371, 379, 431, 461, 525, 595; spraying with, 70, 159, 376, 379, 383, 425, 528, 595, 648; new method of spraying with, 383; tobacco plants dipped in, 380, 652; tests of toxicity of to insects 246, 287 toxicity of, to insects, 246, 287, 373, 422, 528, 537, 627; formulae containing, 258, 286, 379, 425, 431, 461, 525, 595, 652; and Bordeaux mixture, magnesium oxide as a corrective for injury to apple by, 425; and derris, 370; and soy-bean flour, 70; and sulphur, 244, 595; and thiodiphenylamine 379; problems of residues of, 104, 432, 595, 648; other insecticides compared with, 159, 246, 286, 422, 525, 627.

Crypticerya (see Palaeococcus).

Cryptoblabes gnidiella, on vines in Palestine, 191.

Cryptococcus aceris, sp. n., on maple in Caucasus, 680.

Cryptognatha nodiceps (predacious on Coccids), in Fiji, 212; introduced into Hawaii, 366; introduced into other W. Indian Islands from Trinidad, 25, 502.

Cryptogonus orbiculus var. nigripennis, establishment of, against Pinnaspis buxi in Hawaii, **367.**

Cryptohypnus riparius (see Hypnoidus).

Cryptolaemus, utilisation of, against mealybugs in Abkhazia, 454.

Cryptolaemus montrouzieri (predacious on Coccids), ants hindering utilisation of, in S. Africa, 323; utilisation of, in California, 437; establishment of, in France, 266, 267; in Queensland, 345; utilisation of, in Russian Union, 454, 455, 589; relation of development of, to temperature, 589.

Cryptomyzus, status of, 99, 100. (See Capitophorus.)

Cryptoparlatorea pini, sp. n., on pine in India, 224.

Cryptophlebia (see Argyroploce).
Cryptorrhynchus, question of scope

of, **595**.

Cryptorrhynchus gravis, on mango Netherlands Indies, systematic position of, 595.

Cryptorrhynchus lapathi, relation of, to decrease of willow in Britain and Holland, 563; on elm in U.S.A., 499; other foodplants of, 563; systematic position of, 595.

Cryptorrhynchus mangiferae, SVStematic position of, 595.

Cryptostemma calendulaceum, Halotydeus destructor on, in W. in baits for Australia, 131; Listroderes obliquus, 186.

Cryptotermes (see Calotermes).

Cryptus, parasite of Neosyrista similis in Japan, 211.

Cryptus oxymorus, parasite of Setora nitens in Java, 347.

Cryptus sponsor, parasite of Dicranura vinula in Morocco, 196.

Crystal Violet, tests of, in baits for grasshoppers, 134.

Ctenolepisma longicaudata, measures against, in Australia, 510.

Cuba, Tachinid parasite of Diatraea introduced into Louisiana from, 378.

Cubé, dusting with, 71, 177, 179, 249, 250, 286, 417, 426, 435, 497, 498, 584; carriers for, 177, 179; wetting agents improving dusts of, 177, 179; spraying with, 28, 30, 71, 72, 177, 179, 249, 383, 426, 429, 497, 498, 530; spreaders and adhesives for, 28, 30, 71, 72, 177, 179; tests of toxicity of, to insects, 250, 286; effect of weathering on toxicity of, 249; of no value in baits for cutworms, 287; formulae containing, 30, 71, 177, 179, 417, 426, 498, 530; and barium fluosilicate, 177; and sulphur, 71, 72; rotenone content of, 30, 71, 177, 179, 250, 286, 426, 498, 530; determination of rotenone in, 392; other insecticides compared with, 177, 250, 417, 498.

Cucumber, pests of, in Fiji, 366, 641; Ceratia similis on, in Formosa, 618; Myzus ornatus on, in Germany, 688; Megumenum gracilicorne on, in Japan, 441; Dicyphus on, in New South Wales, 449; Diaphania hyalinata on, in Porto Rico, 501; Tetranychus on, in Russian Union, 310; pests on, in greenhouses in U.S.A., 142, 652; Aphids and virus disease of, 672, 686, 687, 688.

cucumeris, Epitrix.

cucurbitae, Dacus (Chaetodacus).

Culex spp., tests of pyrethrum extracts on larvae of, 135.

culinaris, Rhyncolus.

cultratus, Microphanurus.

Cultural Practices, influence of, on insect pests, 221, 346, 389, 655.

Cumin Seed (Stored), Coleoptera in, in Cyprus, 270.

cumingi, Promecotheca. cunea, Hyphantria.

cuneiformis, Saissetia.

cuprea, Anomala.

Cupressus, Phloeosinus bicolor on, in France, 225; Aesiotes leucurus on, in New South Wales, 602.

Cupressus macrocarpa, Phloeosinus spp. on, in France and U.S.A., 225, 324.

Cupressus sempervirens, pests of, in Turkey, 218, 219.

cupreus, Pterostichus (Poecilus); Rhynchites.

Cuprex, 580.

Cuprous Cyanide, dusts and sprays containing. against Lepidopterous tomato pests, 379, 431.

Curculio elephas (on chestnut), bionomics and control of, in France, 358; in Greece, 452; in Italy, 358.

Curculio nucum, on hazel in Greece,

Curculio, Apple (see Tachypterellus quadrigibbus).

Curculio, Cowpea (see Chalcodermus aeneus).

Curculio, Plum (see Conotrachelus nenuphar).

Curled Chervil (see Anthriscus cerefolium).

Curly-top of Beet, relation of Eutettix tenellus to, in Idaho, 286, 412; in other plants, 286; age in beet increasing resistance to, 412.

Currant, Aphids on, in Britain, 333; spray programme for, in Britain, 396; Aspidiotus ostreaeformis on. in Czechoslovakia, 468; pests of, in Germany, 114, 467; pests of, in U.S.A., 178, 179, 596; Parlatoria oleae on, in Uzbekistan, 474: not attacked by Capitophorus korschelti, 100.

Currant, Alpine (see Ribes alpinum).

Britain, Black, pests of, in Britain, 2, 236; new Aphid on, Currant. in in Germany, 100; Aphis varians on, in U.S.A., 596; Dasyneura tetensi confined to, 673; limesulphur causing injury to, 2.

curvipennis, Dacus (Chaetodacus).

Custard Apple (see Anona).

custator, Thyanta.

custos, Auriga (Arma). cuthbertsoni, Analeurops.

Cutworms, baits for, 32, 67, 125, 178, 368, 390; other measures against, 368, 370, 379, 380, 390, 661.

Cyanofumer, 15.

cyanonotus, Cheiloneurus.

cyanura, Mimastra.

Cybocephalus, predacious on Tar-

gionia vitis in Italy, 471.

persicum, Cyclamen Brotolomia meticulosa on, in greenhouses in Austria, 171.

Cyclocephala immaculata, Scoliid parasites of, on sugar-cane in

Florida, 657.

Cyclohexylamine Derivatives, phytocidal and insecticidal properties of, 141.

Cydia funebrana (on plum), in Cyprus, 270; bionomics and control of, in Switzerland, 336.

Cydia glycinivorella (on soy bean), bionomics of, in Japan, 19, 624; in Korea, 19; in Manchuria, 19, 340, 361.

Cydia inopinata, on apple in Man-

churia, 340.

Cydia leucostoma, new parasite of,

on tea in Java, 94.

Cydia molesta (Oriental Peach Moth), in Argentina, 237, 601; in Australia, 293, 382, 544, 639, 640; in China, 450, 667; in Italy, 197, 199, 200, 235, 515; in Manchuria, 340; in Switzerland, 438; in U.S.A., 377, 382; potential distribution of, in Europe, 7; on apple, 340; on pear, 235, 450, 516, 667; bionomics of, 639, 668; parasites and biological control of, 237, 293, 377, 382, 426, 544, 602, 639, 640; other measures against, 199, 235, 450, 515, 516, 639, 668; causes of failure of spraying against, 544; naphthol bands of little value against, 197; survey of data on, 7; attempted utilisation of parasites of, against C. nigricana, 39. Cydia nigricana (Pea Moth), in Canada and Britain, 38; in Russian Union, 109, 589; natural enemies and biological control of. 38, 39, 109, 589.

Cydia pomonella (Codling Moth), 7; in Afghanistan, 575; Algeria, 191-193; in Australia, 20, 57, 186, 293, 390, 554; in Baluchistan, 51; in Canada, 66, 664; in Czechoslovakia, 468; in Denmark, 117; in France, 168, 357, 622; regulations dealing with use of arsenicals against, in France, 168; in Germany, 114, 300; in Holland, 578; in Italy, 4, 104, 197, 235; in Morocco, 283; in Poland, 163, 694; in Russian Union, 109, 587, 589, 610, 623, 683; in U.S.A., 31, 33, 52, 70, 133, 134, 166, 272, 273, 343, 410, 419, 432, 525, 534, 536, 648, 652; road dust apparently increasing infestation of apple by, 33; on pear, 4, 20, 51, 114, 168, 192, 197, 235, 293, 357, 383, 390, 410; on quince, 4, 51, 192; on stone-fruits, 4; on walnut, 4,197; bionomics of, 4,51, 117, 163, 192, 283, 293, 300, 357, 578, 587, 622, 652, 694; application of bioclimatics to study of, 327; parasites of, 4, 109, 293, 579, 588, 589; birds destroying, 114, 588: measures against, 4, 20, 31, 52, 57, 66, 70, 133, 134, 163, 166, 186, 192, 197, 235, 272, 273, 293, 300, 383, 390, 410, 432, 525, 534, 536, 554, 579, 610, 623, 643, 648, 653, 664, 683; tests of insecticides on, 142, 298, 419, 426, 528; resistance to lead arsenate of strains of, 343.

Cydia pomonella putaminana, 197. Cydia splendana (on chestnut), bionomics and control of, in France, 358; in Greece, 452; in Italy, **358.**

Cylas formicarius (Sweet Potato Weevil), risk of introducing, into Japan from Loochoo Islands, 616; factors affecting infestation by, in Porto Rico, 434; measures against, in U.S.A., 373, 379, 539, 643; bionomics of, 373, 379, 616.

cylindrator, Lissonota.

cylindrica, Chaetocnema.

cylindricollis, Sitona.

Cymbidium, Coccid on imported, in Japan, 617.

Cynara scolymus (see Artichoke). Cynipids, records of parasitic, in Germany, 352.

Cynodon, relation of Nomadacris septemfasciata to, in E. Africa, 62, 387, 388.

cyperi, Athesapeuta.

Cyperus, relation of Nomadacris septemfasciata to, in E. Africa, 62, 387, 388; Pseudococcus brevipes on, in Brazil, 23; Nymphula vittalis on, in Korea, 619.

Cyperus rotundus, Spodoptera mauritia on, in Formosa, 363; insects introduced into Australia and Fiji from Hawaii against, 367,546.

Cypress (see Cupressus).

Cyprus, miscellaneous pests in, 270: beneficial insects and biological control in, 270, 271.

Cyrtogaster vulgaris, parasite of Phytomyza ilicis in Britain, 669.

Cyrtomorpha flaviscutellaris, destroying eggs of Austroicetes cruciata in W. Australia, 56.

Cyrtopeltis varians (see Engytatus

geniculatus).

Cyrtorhinus pellicia, ovipositing in egg-cluster of Saccharosydne saccharivora in Florida, 657.

Cytherea infuscata, destroying eggs of Dociostaurus maroccanus in

Spain, 477.

Czechoslovakia, forest pests in, 468, 469, 518, 564; list of leafminers in, 145; miscellaneous pests in, 468-471; popular names of vegetable pests in, 512; beneficial insects in, 145, 518; utilisation of Formica rufa in, 564.

cziceki, Tipula.

D.

Dachicida, types of, in baits for Dacus oleae, 98.

Dachicida P, composition of, 98. Dacivoro, in baits for Dacus oleae, 98, Dacryphalus sumatranus, sp. n., on nutmeg in Sumatra, 512.

dactylidis, Dasyneura.

Dactylis glomerata, Dasyneura dactylidis on, in Britain, 635.

Dactyloctanium aegyptium, Saccharosydne saccharivora on, in Florida, 657.

dactylopii, Leptomastix.

Dactylopius confusus, establishment of Argentine strain of, against Opuntia aurantiaca in Queensland, 345; Cryptolaemus montrouzieri predacious on, 345.

Dactylopius opuntiae, utilisation of strains of, against Opuntia spp. in Queensland, 345.

Dacus, in Fiji, 212, 213, 366; in Hawaii, 366; on Citrus in Samoa, 365; parasites and biological control of, 212, 213, 365, 366, 562.

Dacus ciliatus (brevistylus), bionomics of, on Coccinia indica in

Madras, 561.

acus cucurbitae, on melon in Baluchistan, **577**; experiments on infestation of fruits by, in Dacus cucurbitae, Formosa, 206; relation of temperature to risk of introduction of, into Japan, 615; recently introduced into Kenya, 667; on Coccinia indica in Madras, 562: legislation against importation of, into U.S.A. from Hawaii, 344; measures against, 344.

Dacus curvipennis, rare on banana

in Fiji, 366.

Dacus ferrugineus, fruits infested by, in Baluchistan, 577; parasites of, on Solanum auriculatum in Madras, 577.

Dacus ferrugineus dorsalis, experiments on infestation of fruits by, in Formosa, 206; relation of temperature to risk of introduction of, into Japan, 615.

Dacus ferrugineus tryoni, in Queensland, 345, 365, 602; in grapes, 365: breeding experiments with, 345; Syntomosphyrum indicum New introduced from South

Wales against, 602.

Dacus oleae (Olive Fly), seasonal occurrence of, in Cyprus, 270; in Greece, 453; in Italy, 98, 568; intercepted in Russian Union, 439; baits for, 98, 270; doubtful value of bait-sprays against, 270, 453; survey of data on, 568.

Dacus passiflorae, in Fiji, 212, 366, 605; host fruits and development of, 212; 366; new parasite of,

605.

Dacus psidii, not occurring in Fiji, 366; bionomics and control of. in New Caledonia, 131; on guava and Inocarpus in Society Is., 294.

Dacus xanthodes, in Fiji, 212, 366; on granadilla, 212.

Dacus zonatus, fruits infested by. in Baluchistan, **577.**

dahlemica, Phlugiola.

Dahlia, Ceresa bubalus on, France, 689; Nysius turneri on, in Tasmania, 226.

Dakota, North, potato pests in, 70, 71; grasshoppers in, 33, 655. Dakota, South, Dendrosoter scoly-

tivorus in, 324.

Dalbergia sissoo, Plectoptera reflexa on, in India, 612.

dalmanni, Habrolepis.

Dandelion (see Taraxacum offici-

Danzig, Diprion pini and its natural enemies in, 105, 397.

daricella, Leucoptera.

darwiniensis, Mastotermes.

Dascillus cervinus, on pasture in Scotland, 122.

seriatus, Dasus attacking seed potatoes and sugar-cane Hawaii, 367.

Dasus simplex, on tobacco in S. Rhodesia, 125: measures against, on coffee in Tanganyika,

Dasychira pudibunda, predacious enemies of, in forests in Germany,

Dasyneura airae, on Aira flexuosa in France, **634.**

Dasyneura alopecuri, on Alopecurus pratensis, 635.

Dasyneura alpestris (on Arabis), synonymy of, 673.

Dasyneura arabis (see D. alpestris). Dasyneura brassicae, on swedes in Britain, 332.

Dasyneura dactylidis, on Dactylis glomerata in Britain, 635.

Dasyneura fagicola, sp. n., bionomics and control of, on beech in Germany, 353.

festucae, Dasyneura sp. n., Festuca in Britain, 635.

Dasyneura graminis (on Agrostis), probably in Britain, 635; in New York, **635.**

Dasyneura plicatrix, food-plants and status of, 673.

Dasyneura pyri (on pear), bionomics of, in Italy, 296; in New Zealand, 159; natural enemies of, 159, 296; measures against, 297.

Dasyneura ribicola, food-plant and status of, 673.

Dasyneura schneideri (see alpestris).

Dasyneura tetensi, food-plant and status of, 673.

Dasyneura torquens (see Contarinia nasturtii).

Dasyneura triseti, sp. n., bionomics of, on Trisetum flavescens in Austria, 635.

Dasynus manihotis, new parasites of, in Java, 19.

Dasyscapus parvipennis (parasite of Selenothrips rubrocinctus), in Trinidad, 25, 239; failure to establish, in Grenada, 25; introduction of, into Jamaica, 239.

Date Palm, Pseudococcus brevipes on, in Brazil, 23; new Eriophyid on, in California, 288; pests of fruit of, in Egypt, 506.

Datura stramonium, Phthorimaea operculella on, in Brazil, 289; Protoparce spp. on, in U.S.A., 252, 253.

debilis, Micromalthus.

Decadarchis, on coconut in Fiji, 212. Decadarchis heterogramma, on coconut in Fiji, 213; feeding on pupae of Tirathaba trichogramma, 213.

Decadarchis psammaula, on coconut in Fiji, 212.

decemlineata, Leptinotarsa.

decoratus, Geocoris.

Dectes spinosus, introduced into Australia against Xanthium pungens, 546; liable to attack artichoke, 546.

Decticus albifrons, measures against, in Libya, 478.

Decticus verrucivorus, in Jugoslavia,

defoliaria, Erannis.

Degras (see Wool Grease).

delavoiei, Dicranura vinula.

Delaware, Cydia pomonella in, 534; Popillia japonica and its biological control in, 377, 532.

delhiensis, Azotus.

Delia (see Hylemyia).

delicatus, Macrocentrus.

Delphacodes striatellus (on rice), in Japan, 362; in Manchuria, 340; oviposition of, 362.

deltae, Mesolecanium.

Deltocephalus (see Laevicephalus). Dendrocalamus giganteus, resistance of, to Dinoderus minutus, 501.

Dendroctonus brevicomis, on Pinus ponderosa in U.S.A., 599.

Dendroctonus convexifrons, on Pinus ponderosa in New Mexico, 599.

Dendroctonus frontalis, in U.S.A., 324, 377, 599; on Pinus taeda, 377, 599; new parasite of, 324.

Dendroctonus piceaperda, on spruce in Canada, 275, 276, 369.

Dendroctonus ponderosae, outbreak of, on pines in U.S.A., 599.

Dendroctonus pseudotsugae, on Pseudotsuga taxifolia in U.S.A., 599.

Dendrol, 526, 527. dendrolimi, Rhogas.

Dendrolimus albolineatus, new parasite of, in Sakhalin, 91.

Dendrolimus pini, parasite of, on pine in Germany, 354; studies on cold-hardiness in, in Russia, 404, 405.

Dendrolimus punctatus, bionomics and control of, on pine in China,

449, 451.

Dendrolimus spectabilis (on pine), D. punctatus recorded as, in China, 449; in Japan, 91, 209, 211, 441; effects of temperature and humidity on eggs of, 441; parasites and hyperparasites of, 91, 209, 211.

Dendrosoter, key to species of, in

U.S.A., **324.**

Dendrosoter integer, sp. n., parasite of Scolytids in U.S.A., 324.

Dendrosoter protuberans, parasite of Scolytids in Turkey, **522**.

Dendrosoter scaber, sp. n., parasite of Scolytids in U.S.A., 324.

Dendrosoter scolytivorus, in U.S.A., 324.

Dendrosoter sulcatus, sp. n., parasite of Scolytids in U.S.A., 324.

Denmark, miscellaneous pests in, 117, 518, 673; beneficial insects and biological control in, 117, 518.

denticornis, Cerodontha; Limo-thrips.

denticoxa, Eurytoma (see E. nesiotes).

dentipes, Monodontomerus.

deplana, Lithosia (see Eilema depressa).

depressa, Eilema; Sarcophaga.

depressella, Emmalocera.

depressum, Steirastoma.

depressus, Palorus.

depunctalis, Nymphula.

Derelomus (see Prosoestus).

Dermatodes aeruginosus, on tung nut trees in Java, 523.

Dermestes, Necrobia spp. associated with, **584**.

Dermestes frischi, damaging timber in Germany, 7.

Dermestes lardarius, damaging timber and masonry in Germany, 7, 520; new fumigant against, attacking silkworm cocoons in

Italy, 198; in house insulation in U.S.A., 640; review of measures against, 198.

Dermestes peruvianus, damaging timber in Germany, 7.

derogata, Sylepta.

Derostenus, parasite of Dasyneura

pyri in Italy, 296.

Derris, against Aleurodids, 530, 642; against Aphids, 30, 371, 395, 497, 498; against Jassids, 21, 115, 281, 413, 537; against other Rhynchota, 184, 227, 353, 417, 446, 523, 530; against Coleoptera, 69, 155, 156, 158, 184, 241, 249, 250, 269, 286, 298, 373, 426. 532. 584. 634: against Diptera, 297, 497, 556; against Lepidoptera, 17, 28, 33, 177, 179, 184, 191, 250, 251, 269, 365, 370, 380, 400, 431, 663, 689; in bait-sprays for Cydia pomonella, 300, 301; against Paratetranychus pilosus, 429; against sawflies, 21, 117, 400; with, against Sexava, tests 277; against thrips, 233, 234, 465, 493, 525; action of, on honey bees, 402; tests of toxicity of, to insects, 17, 184, 249, 250, 365, 634, 642; dusting with, 17, 21, 30, 69, 155, 156, 158, 179, 184, 227, 249, 250, 251, 269, 277, 281, 286, 298, 353, 365, 370, 371, 373, 380, 395, 400, 402, 417, 426, 431, 497, 498, 523, 537, 584, 634; carriers for, 69, 179, 249, 250, 371, 380, 523, 537, 634; effect of addition of chloropicrin to dusts of, 471; spraying with, 17, 21, 30, 69, 115, 117, 156, 177, 179, 233, 234, 241, 249, 250, 286, 298, 395, 402, 413, 426, 446, 493, 497, 498, 525, 530, 532, 642, 663, 689; formulae containing, 17, 30, 177, 179, 184, 241, 300, 426, 493, 498, 523, 530, 532, 537, 689; and cryolite, 370; and oil emulsions, 33, 493, 642, 689; intermediary solvents for, when used with oils, 530; and pyrethrum, 155, 156, 158, 298, 471, 634; and soap, 233, 446, 530; other spreaders and adhesives for, 28, 30, 177, 179, 241, 249, 525, 530, 532; and tar distillates 298, 689; thiuram sulphides compatible with, 532; rotenone contents of, 17, 30, 69, 70, 179, 184, 233, 249, 250, 251, 269, 286, 365, 370, 371, 373, 380, 402, 417, 426, 493, 497, 498, 523, 530, 537, 642; determination of

rotenone in, 392, 493; resinate incorrect name for extracts of, 78; tests of effects of rain and light on deposits of, 69, 241; translocation of, in plants, 234; survey of data on, as an insectiother insecticides cide, 104; compared with, 17, 155, 177, 249, 250, 251, 286, 298, 373, 498,

Devris spp., peşts of field and stored, 502, 538; cultivation and preparation for market of, 294, 336, 624; insecticidal properties and constituents of, 250, 294, 623.

Derrisol, tests of, against thrips and

Psila vosae, 127, 497.

destructor, Aspidiotus; Corymbites (Ludius) aeripennis; Halotydeus; (Glycyphagus); Lepidoglyphus Mayetiola; Merisus; Scolytus (see S. scolytus); Tanymecus; Tribolium.

Detal (containing dinitro-o-cresol

q.v.), 297, 298. detersa, Euxoa.

Deudorix bimaculata (see Virachola). Dewberry, pests of, in U.S.A., 31, 79; Dasyneura plicatrix on, 673; susceptibility of varieties of, to Agrilus ruficollis, 79.

Dexia rustica (parasite of Melolontha), in Denmark, 117, 518; in Germany, 608; bionomics of,

Diabrotica soror, not transmitting virus disease of celery in California, 590.

Diachasma aino, sp. n. (parasite of Euphranta), in Japan, 338, 620; bionomics of, 338.

Diacrisia isabella, food-plants of, in Oklahoma, 340.

Diacrisia lubricipeda, Troilus luridus

attacking, 354. Diacrisia obliqua, bionomics of, on lima beans in China, 129; Trichogramma reared on, in India, 612.

Diacrisia virginica, food-plants of, in Oklahoma, 340.

diadema, Stomatodexia.

Dialeurodes chittendeni, on rhododendrons in Britain, 295.

Dialeurodes citri, in U.S.A., 256, 259, 530, 642, 649; on Citrus, 256, 642, 649; on ornamental plants, 259, 530; measures and tests of insecticides against, 259, 530, 642, 649.

Dialeurodes citrifolii, on Citrus in Florida, 256.

Diammonium Phosphate, tests of, in baits for Dacus oleae, 98.

dianaria, Boarmia selenaria.

Diaphania hyalinata, on cucurbits in Porto Rico, 501; water-melon not attacked by, 501.

Diaprepes, probably on limes in

Grenada, 25.

Diaprepes abbreviatus, on sugar-

cane in Barbados, 660.

Diarthronomyia (Chrysanthemum Gall-midge), distribution food-plant range of, 673; legislation against introduction of, into Britain, **356.**

Diarthronomyia hypogaea, 356: midge on commercial chrysanthemums distinct from, 673.

Diarthrothrips coffeae, measures against, on coffee in Tanganyika, 127.

Diaspidiotus (see Aspidiotus). diaspidis, Aphytis (Aphelinus).

Diaspiniphagus similis (see Coccophagoides).

Diaspis bromeliae, utilisation of Chilocorus politus against, pineapple in Mauritius, **611.**

Diaspis piricola (see Epidiaspis

leperii).

Diaspis visci (on Juniperus), in Czechoslovakia, 469; measures against, in New York, 529; on other conifers, 469.

Diastase, accelerating development of Hylotrupes bajulus, 335.

Diatomaceous Earth, unsatisfactory as a carrier for dinitro-ocyclohexylphenol, 261. Celite.)

Diatraea (on sugar-cane), in Br. Guiana, 524; in Grenada, 25; in Trinidad, 24; utilisation of Metagonistylum minense against,

24, 505, 524.

Diatraea canella (on sugar-cane), in Br. Guiana, 505; in W. Indies, 182, 183, 272, 659; relation of parasites of D. saccharalis to, 182, 183, 505, 659; measures against, 272.

Diatraea lineolata, erroneously recorded in St. Lucia, 182.

Diatraea saccharalis (on sugar-cane), in Br. Guiana, 182, 505; in U.S.A., 371, 372, 378, 382, 656, 658; in W. Indies, 25, 182, 183, 378, 382, 424, 435, 502, 658, 659, 660; susceptibility of varieties of sugar-cane to, 238, 372, 378; Cordyceps barlieri infesting, 660; parasites and biological control

of, 25, 182, 183, 238, 272, 378, 382, 424, 435, 502, 505, 656, 658, 660; other measures against, 272, 371, 378, 656.

Diatraea (Argyria) sticticraspis, on sugar-cane in India, 291, 558, 576; parasite of, 291; measures against, 576; damage caused by, to sugar-cane varieties, 558.

Diatraea venosata (on sugar-cane), in Formosa, 206, 363, 447; in India, 558; in Mauritius, 611; natural enemies and biological control of, 363, 364, 447, 611; relation of growth of sugar-cane to oviposition of, 206: damage caused by, to sugar-cane varieties, 558.

diatraeae, Apanteles; Lixophaga; Theresia (Paratheresia) (see T.

claripalpis).

Dibrachys, parasite of spruce sawflies in Czechoslovakia, 518.

parasite Dibrachys affinis, Aegeria typhiaeformis in Italy, 630.

Dibrachys australia (see Pseudomicromelus).

Dibrachys cavus (boucheanus, clisiocampae), hosts of: in Argentina, 238; in Australia, 639, 640; in Germany, 630; in Italy, 629, 630; in Poland, 671; utilisation bionomics of, of, 238; 630: economic status of, 631: synonymy of, 434.

Diceroprocta vitripennis, bionomics of, on cotton in Arkansas, 165.

Dichlorethyl Ether, uses of, against insects, 177, 272, 379, 526.

Dichlorobenzene, and oils. treating timber against carpenter bees, 337.

Dichloropentane, against Aegeria

pictipes, 527.

Dichocrocis punctiferalis, damaging teak seed in Burma, 693; measures against, on peach in China, 450; on papaya and Citrus in Queensland, 603.

Dichroplus, in Argentina, 214.

dicincta, Mylabris.

Dicky Rice Weevil (see Maleuterpes spinipes).

Dicranura vinula var. delavoiei, bionomics of, on Populus nigra in Morocco, 196.

Dictamnia, measures against, on cacao in New Guinea Territory,

dictyospermi, Chrysomphalus.

Dicyphus, food-plants and control of, in New South Wales, 449. differentialis, Melanoplus.

difficilis, Apanteles; Aphaereta. Digitalis, Macrosiphum solani on,

in Germany, 687.

Digitaria sanguinalis, Saccharosydne saccharivora on, in Florida,

Diglycol Laurate and Oleate, as emulsifiers for oils, 642.

dignus, Apanteles.

dilatatum, Piesma quadratum.

dimmocki. Pikonema.

Dimmockia, 289.

dimorphus, Ephialtes (Calliephialtes). Dinapate wrighti, on Washingtonia filifera in California and Mexico,

Dinarmus coimbatorensis, sp. n., parasite of weevils in India, 506.

diniana, Enarmonia (Semasia). Dinitro-cresol, against Aphids, 237, 332, 533; against Capsids, 45, 237; against Coccids, 529, 535; against Coleoptera, 171, 298, 495, 606, 608; against Lepidoptera, 237, 400, 535; against 353; Pentatomids, against Rhagoletis cerasi, 297; against sawflies, 400; dusts containing, 171, 297, 353, 400, 495, 606, 608; sprays containing, 45, 171, 237, 332, 495, 529, 530, 533, 535, 608; and oils, 236, 332; causing injury to trees, 400, 529, 530; toxicity of, to fish, 514; other insecticides compared with, 231, 237, 298, 514, 534. Sodium Dinitro-ortho-cresylate.)

Dinitro-ortho-cyclohexylphenol, against Aphids, 429, 461, 533, 535; against Coccids, 529, 534, 535, 649; against other insects, 535, 649; against mites, 261, 429, 461, 535, 649, 650; destroying predacious mites, 429; preparation of dusts containing, 261, 461, 650; sprays containing, 429, 461, 529, 530, 533, 534, 535, 649; salts of, 649; and oils, 429, 529, 530, 534, 535; causing injury to plants, 529, 530, 535, 649, 650; physical and chemical properties of, 261, 649; compared with tar distillates, 429, 534.

Dinoderus brevis, measures against, in bamboo in Jamaica, 181.

Dinoderus minutus, in bamboo in W. Indies, 181, 423, 501; resistance of certain species of bamboo

to, **501**; Peregrinator biannulipes predacious on, 423; measures against, 181.

Dioctes (see Angitia).

(Holodiomphalia, Lachnosterna trichia).

Dioryctria peyerimhoffi, new parasite of, on Cedrus in Morocco, 58. dioryctriae, Apanteles.

Diospyros kaki (see Persimmon).

Diparopsis castanea (Red Cotton Bollworm), in S. Africa, 488; in Nyasaland, 125, 222; in S. Rhodesia, 492; in Sudan, 573; in Tanganyika, 492; alternative food-plants of, 488, 489, 573; bionomics of, 488, 492, 573; Pheidole attacking, 489; measures against, 222, 573.

Diphenyl, toxicity of, to grass-

hoppers, 134.

Diplachne, relation of Nomadacris septemfasciata to, in Tanganyika, 387.

Diplogaster, parasite of Horistonotus uhleri in S. Carolina, probably infesting Dexia rustica in Denmark, 519.

Diploptera dytiscoides, measures against, on coconut in Hawaii,

Diplosis, on pine in Japan, 339.

Diplosis humuli (see Contarinia). mori(Mulberry Gall-Diplosis midge), bionomics of, in Korea, 337.

Dirhicnus magnicornis Diprion, parasitic on, in Poland, 232.

Diprion abietis, on spruce in Quebec,

464. Diprion frutetorum, records of, in U.S.A., 600.

Diprion lecontei (on pines), in Canada, 53, 464; in U.S.A., 461; measures against, 53, 461.

iprion pinetum (on pines), measures against, in Michigan, Diprion

461; in Quebec, 464.

Diprion pini (on pines in Europe), bionomics and population studies of, 105, 397; natural enemies and diseases of, 105, 232, 397, 398; characters of, 7, 104.

polytomum (European Diprion Spruce Sawfly), in Canada, 176, 275, 276, 368, 381, 382, 464, 647; in U.S.A., 176, 368, 381, 382, 647; extent of damage to spruce by, 275, 276; bionomics and forms of, 176, 369, 647; parasites and biological control of, 276, 369, 381, 382.

Diprion sertifer, on Pinus mugo in Czechoslovakia, 469; new Tritneptis parasitic on, in U.S.A., 434.

Diprion similis (on pines), measures against, in Michigan, 461; natural enemies and diseases of, in Poland, 670; character distinguishing D. pini from, 7.

diprioni, Tetracampe.

diprionis, Tritneptis. dipsacea, Heliothis (Chloridia).

Diptilomiopus prunorum, sp. n., on Prunus spp. in California, 288. Dirhicnus alboannulatus, parasite of Sphinx pinastri in Poland, 677.

Dirhicnus magnicornis, parasite of

Diprion in Poland, 232.

Dirhinus (parasite of Dacus spp.), introduced into Cook Islands and Samoa, 212, 365; bionomics and utilisation of, in Fiji, 212, 213. Dirhinus auratus, parasite of Dacus

ferrugineus in Madras, 577.

Dirhinus giffardi, introduced into Porto Rico against Anastrepha, 502; reared on Musca domestica, 502.

dirhodum, Macrosiphum (Amphorophora).

discedens, Phyllotreta vittata.

discodirus, Opilo. discoidalis, Exora.

dispar, Bothriophryne; Lymantria (Porthesia, Porthetria).

disparis, Ichneumon; Pimpla.

dissimilis, Xylocopa.

Dissosteira carolina, on tobacco in Connecticut, 178.

Dissosteira longipennis, study of flights of, in Colorado, 646.

disstria, Malacosoma.

distigma, Chilocorus.

distincta, Galerucella.

distinguendus, Haplothrips; Lariophagus.

diversicolor, Chrysomphalus pinnu-

lifer.

Diversinervus, bionomics and utilisation against Saissetia oleae of introduced species of, in California, 424; key to species of,

Diversinervus stramineus, sp. n., parasite of Saissetia persimilis in Transvaal, 40.

diversus, Microdus (Bassus).

divinatorius, Liposcelis (Troctes).

divisa, Monolepta.

divisus, Macrosteles.

Dociostaurus crucigerus brevicollis, in Russian Union, 320, 386; ecology and migrations of, 386.

Dociostaurus maroccanus, in Fr. N. Africa, 477, 480; in Cyprus, 270; loss due to, in Greece, 478; in Jugoslavia, 479; in Libya, 478; in Poland, 403; in Portugal, 479; in Spain, 477; in Transcaucasia, 321; crops attacked by, 478; phases of, 321, 480; breeding centres and ecology of, 321, 477, 480; Cytherea infuscata destroying eggs of, 477; measures against, 477, 478, 479, 480.

doddi, Cactoblastis.

Dodecyl Rhodanate (see Thiocyanates).

Dogwood (see Cornus).

Dolba hylaeus, parasitised by Apanteles congregatus in U.S.A., 253.

Doleschallia bisaltide, on Eranthemum in Fiji, 641.

Dolichocolon australe, sp. n., parasite of Laphygma exempta in Queensland, 263.

Dolichocolon orbitale, sp. n., parasite of Hepalia machaeralis in India,

Dolichoderus bituberculatus, in houses in Formosa, 446.

Dolichopoda, in caves in Mediterranean Region, 604.

Dolichopoda bormansi, in caves in Corsica, 604; imported into England, 604; synonymy of, 604.

Dolichos lablab, attraction of strains of, for Heliothis armigera in southern S. Africa, 223, 488, 489, 491; Systates exaptus on, in S. Rhodesia, 124.

Dolichos melanophthalmus, Plodia interpunctella in seed samples of, in Italy, 198.

Dolycoris, on cereals, etc., in Germany, 352, 631, 632; type of injury to cereals by, 352, 353; bionomics and ecology of, 631, 632; measures against, 353.

Dolycoris baccarum, parasites of, on cereals in Germany, 632; on tobacco in Poland, 162.

Dombeya rotundifolia, relation of Dysdercus nigrofasciatus to, in S. Africa, 490.

domestica, Musca.

domesticus, Glycyphagus. dominica, Rhyzopertha.

Dominican Republic, new Coccid on pines in, 392; Coccinellids introduced into, 502.

Doralis (see Aphis).

Dorcadion, and white ear of wheat in Russian Union, 680.

dorsalis, Dacus ferrugineus; Scirtothrips.

Doryctes gallicus, parasite of Chrysobothris igniventris in Turkey, 219.

Dosage-Mortality Curve, 66, 111.

Douglas Fir (see Pseudotsuga taxifolia).

Douglas Fir Beetle (see Dendroctonus pseudotsugae).

Draecenura pelochra, on beans in Fiji, 641.

Dragonflies, destroying Nymphula vittalis in Korea, 619.

Drepanothrips reuteri, bionomics of, on vines and oak in Crimea, 43. Dreyfusia (see Chermes).

dromedarius, Xiphydria (see X. prolongata).

druparum, Callimome.

Dryocoetes abietinus, sp. n., on Abies sachalinensis in Japan and Sakhalin, 615.

Dryocoetes minor, on oak in Sicily, 522; bionomics of, on chestnut in Turkey, 522.

dubia, Anomala. ducens, Feltia.

Ducks, utilisation of, against locusts in China, **450.**

dumetana, Tortrix.

Dung, in baits for locusts and Tenebrionids, 320, 321; trap heaps of, for *Oryctes*, 347.

Dusting, types of equipment for, 247, 266, 299, 325, 371; use of aircraft for, 74, 75, 299, 514, 610, 645; spraying compared with, 177, 179, 249, 266, 495, 497, 498, 514, 610.

Dusts, experiments with new proprietary, against forest pests, 231; uses of oil-impregnated, 75, 308; toxicity of inert, to Bruchus obtectus, 531; cleaning habit of insects in relation to, 139; effect of electric charges and carriers on adhesiveness of, 309; relation of dilution to insecticidal efficiency of, 139.

Dutch Elm Disease (see Ophiostoma ulmi).

Dutox, **70, 425.** (See Barium Fluosilicate.)

Dyk's Method, for estimating prevalence of *Lymantria monacha*, **400.**

dysderci, Hyalomyia (Euphorantha).

Dysdercus (Cotton Stainers), in S. Africa, 489, 490, 491; in Brazil, 22, 34, 86; in Nyasaland, 125; in Peru, 385; in Sudan, 574; susceptibility of varieties of cotton to, 222; bionomics and alternative food-plants of, 489, 490; associated with boll-rots, 34, 222, 290, 389, 491; natural enemies of, 22, 86, 574; measures against, 574.

Dysdercus cardinalis, on cotton in

Kenya, 666.

Dysdercus cingulatus (on cotton), measures against, in Philippines,

184; in Siam, 50.

Dysdercus fasciatus (on cotton), bionomics of, in S. Africa, 489, 490, 491; in Kenya, 666; in N. Rhodesia, 222, 492; in S. Rhodesia, 492; incidence of, in Sudan, 573; in Tanganyika, 492; alternative food-plants of, 222, 489, 490, 491, 492, 573; effects of temperature and humidity on, 491.

Dysdercus honestus, natural enemies of, on cotton in Brazil, 86.

Dysdercus howardi, toxicity of selenium-treated cotton to, in Trinidad, 221.

Dysdercus intermedius, on cotton and Sterculia in southern Africa, 489, 492.

Dysdercus megalopygus (see D. cingulatus).

Dysdercus mendesi, natural enemies of, on cotton in Brazil, 22, 86.

Dysdercus mimulus, and boll rots in cotton in Arizona, 416, 417; measures against, 417.

Dysdercus mimus, on cotton in Ecuador, 164; on wild malvaceae in Peru, 164.

Dysdercus nigrofasciatus (on cotton), bionomics and alternative foodplants of, in S. Africa, 489, 490, 491; in Kenya, 666; in Sudan, 573.

Dysdercus orientalis, in Kenya, 666. Dysdercus ruficollis, natural enemies of, on cotton in Brazil and Peru, 22, 86, 164; measures against, 164.

Dysdercus sidae, new parasite of, in Queensland, **57.**

Dysdercus superstitiosus (on cotton), in N. Rhodesia, 222, 492; in S. Rhodesia, 492; in Sudan, 573, 574; alternative food-plants of, 222, 492, 573, 574.

Dysdercus suturellus, physiological factors affecting abundance of, 258.

Dysoxylon hamiltonii, Xylocopa latipes in timber of, in India, 337. dytiscoides, Diploptera.

E.

Earias, parasite of, on cotton in India, 575; on cacao in New Guinea Territory, 544.

Earias chromataria, bionomics of, on cotton in Philippines, 183; experimentally attacked by Trichogramma minutum, 184; measures against, 184.

Earias fabia, bionomics and control of, in Philippines, 183, 184; on

cotton in Siam, 50.

Earias huegeli, on cotton in Queensland, 603.

Earias insulana (on cotton), in Cyprus, 270; in Sudan, 573.

Earthworms, Lacon murinus feeding on, 398.

Earwig, European (see Forficula auricularia).

Eberhardia, bionomics and identity of species of, on mushrooms in Britain, 1, 2.

Eberhardia alberti, 1.

Echinochloa crus-galli, Chilo simplex on, in Manchuria, **340.**

Echinochloa pyramidalis, relation of Nomadacris septemfasciata to, in Tanganyika, **387**, **388**.

Echinochloa stagnina, 12; Nomadacris septemfasciata not found on, in Fr. W. Africa, 571.

echinopus, Rhizoglyphus.

Echthromorpha intricatoria, parasite of cutworms in New South Wales, 279.

Economic Entomology, general papers on, 261, 407, 531, 600; value of collections in work on, 568; publications on, in Finland in 1938, 694; problems and development of: in India, 391, 448; in Italy, 474; in Russian Union, 214; in U.S.A., 261, 407. Ecpantheria icasia, on vanilla in

Ecpantheria icasia, on vanilla ir Porto Rico, **501.**

ectypus, Pheletes.

Ecuador, Dysdercus mimus on cotton in, 164.

efformata, Anaitis.

Egg-plant, Phthorimaea operculella on, in Brazil, **289**; Epilachna vigintioctomaculata on, in Fiji,

641: Poecilocerus pictus on, in India, 577; pests of, in Jamaica, 239; pests of, in Japan, 225, 446, 555; E. vigintioctomaculata on, in Korea, 362; Paratrioza cockerelli on, in Texas, 651; Lygus lucorum possibly transmitting virus disease in, 225, 555.

Egypt. Acherontia atropos on olive in, 175; new Aphids in, 160; Coccids in, 506, 507, 508, 509; new introduced parasite of Phenacoccus hirsutus in, 506; Achrysopophagus aegyptiacus reared from Coccinellids in, 40; pests of dates in, 506; Gryllotalpa spp. in, 509; locusts in, 481; pests of stored products in, 219, 506; Aphelinus mali imported into Palestine from, 127.

Eilema depressa, characters distinguishing Lymantria monacha

from, 466.

Eirenephilus longipennis, bionomics and control of, in Japan, 618. Elaeagnus, Chrysomphalus spp. on, in Japan and New Zealand, 91, 524.

Elaeis guineēnsis, pests of, in

Dahomey, 347-349.

Elacochlora viridicata, in Argentina, 214.

Elaphria spp., parasites of, on sugar-cane in Florida, 657.

Elasmopalpus lignosellus, on sugarcane in Florida, 657; food-plants of, in Porto Rico, 501.

Elasmus, parasite of Pyrausta aurata in Morocco, 193.

Elasmus zehntneri, parasite of Scir-pophaga spp. in Mysore, 291.

Electric Cables, Xylocopa latipes boring in lead sheathing of, in India, 337.

Electric Charges, effects of, on dusts and sprays, 309, 411.

Electricity, uses of, against pests, 31, 137, 138, 260, 442.

elephas, Curculio (Balaninus).

Elettaria cardamomum, Eupterotid on, in Mysore, 625.

Elgeté, composition and use of, against whiteflies, 530.

Elgetol (containing dinitrocresol q.v.), 529, 530, 535.

elisus, Lygus.

Ellopia athasaria, 143.

Ellopia fiscellaria, in forests in Quebec, 464.

Elm (Ulmus) Galerucella luteola on, in Austria, 324; pests of, in Britain, 58, 65; restrictions on

importation of, into Britain, 356: Alsophila pometaria on, in Canada, 662-664 : G. luteola on, in France, 324; pests of, in Holland, 269, 396; pests of, in Japan, 443, 618; Eriosoma lanuginosum on, in Tasmania, 20; pests of, in U.S.A., 28, 68, 69, 77, 78, 81, 132, 138, 144, 190, 324, 378, 418, 420, 427, 429, 499, 529, 532, 644, 649; bark-beetles and Ophiostoma ulmi on, 65, 68, 77, 138, 180, 378, 396, 529, 645; age and species of, in relation to O. ulmi, 66; Pyrausta nubilalis experimentally fed on, 229; lead arsenate causing injury to, 532. Elm Bark-beetles (see Hylastes

rufipes and Scolytus multistria-

Elm Leaf Beetle (see Galerucella luteola).

Eloeidiphilos aliberti, bionomics and control of, on oil palms in Dahomey, 349.
ongatus, Telenomus (Phanurus)

elongatus, beneficiens.

Erythroxylon Eloria novesi, on coca in Peru, 165.

elutella, Ephestia.

emarginata, Calpe (Oraesia).

Emmalocera depressella, on sugar-cane in India, 558, 559, 577; effect of infestation by, on sugar-cane varieties, 558, 559.

Emmelia trabealis, on ramie in Transcaucasia, 587.

Empoasca, factors affecting attacks of, on cotton in Queensland, 281; characters of species of, allied to E. fabae in U.S.A., 83; measures against, 281.

Empoasca abrupta, in California, 83; shown to be distinct from E. fabae 93.

Empoasca biguttula, on cotton in Manchuria, 340.

Empoasca fabae, in U.S.A., 53, 70, 71, 82, 83, 421, 595; on beans, 82, 83, 595; on potato, 53, 70, 71, 421; bionomics of and species allied to, 82, 83; E. abrupta distinct from, 83; measures against, 53, 70, 71, 421, 422, 595.

Empoasca fabalis, measures against. on lima beans in Porto Rico, 501.

Empoasca filamenta, not transmitting potato viruses in Oregon, 84.

Empoasca flavescens, on tobacco in Japan, 210.

Empoasca lybica, associated with "crazy top" on cotton Sudan, 574.

Empoasca terrae-reginae, on cotton in Queensland, 281.

Empoasca viridigrisea, on tomato in

Queensland, 281. Empusa, infesting Dysdercus spp.

in Brazil, 86.

Empusa grylli, infesting locusts in S. Africa, 91, 92; infesting grasshoppers in U.S.A., 75, 427; studies and attempted utilisation of, 92, 427.

Enaria melanictera, parasite of, in

Madagascar, 611.

Enarmonia caryana (Hickory Shuckworm), bionomics of, U.S.A., 377.

Enarmonia diniana, bionomics of, on larch in Switzerland, 169.

Enarmonia minutana, on poplar and willow in Morocco, 196.

prunivora, Enarmonia measures against, on apple in Quebec, 664. Encarsia formosa (parasite

Trialeurodes vaporariorum), utilisation of, in Britain, 333, 437. probably partenopea,

imported into Britain in pupae of Aleurodes brassicae from Italy, 265.

Encarsia tricolor, probably imported into Britain in pupae of Aleurodes brassicae from Italy, 265.

endogena, Polyodaspis.

Endrosis lacteella, eggs of, 49.

Engytatus geniculatus, predacious on Protoparce sexta in Florida, 251.

Enicospilus antancarus, parasite of Diatraea venosata in Mauritius, 611.

Enicospilus purgatus, parasite of cutworms in Florida, 657.

Enneadesmus forficula subsp. capensis, in S. Africa, 60.

Enneapogon brachystachys, relation of Locustana pardalina to, in S. Africa, 394.

Entedon auronitens var. turcicus, n., parasite of Lithocolletis platani in Turkey, 219.

Entedon pempheridis, sp. n., parasite of weevils in India, 506.

Entomology, books on, 170, 455: preparation of papers on, 138. (See Economic Entomology.)

Entomoscelis adonidus, on crucifers in Russian Union, 439.

Ephestia, suitability of, as host for Trichogramma evanescens, **36.**

Ephestia calidella, measures against, in dates in Egypt, 506.

Ephestia cautella, sources of infestation of stored cacao by, in Ceylon, 638; in dates in Egypt, 506; parasite of, in copra in Fiji, 213; mass rearing of Trichogramma japonicum on, in Japan, 619; measures against, 506, 638.

Ephestia elutella, in Britain, 154; bionomics of, in Ceylon, 638; in Germany, 521; in S. Rhodesia, 125; in stored tobacco, 125, 154; in other stored products, 154, 521, 638; measures against, 125,

154, 521, 638.

figulilella, Ephestia | measures against, in dates in Egypt, 506.

Ephestia kuehniella (in stored products), in Britain, 391, 521; in Germany, 400; in Italy, 172; in Russian Union, 685; in U.S.A., 80; bionomics of, 399; determination of food requirements of, 64, 65; natural enemies of, 264, 265, 400, 521; adaptation of Chelonus annulipes to, 80; suitability of, as host for Trichogramma spp., 35, 391, 589; measures against, in flour mills, 400, 685.

Ephialtes argentinus (parasite of Cydia molesta), bionomics and suggested utilisation of, Argentina, 601, 602.

Ephialtes dimorphus, sp. n., parasite of Platyedra gossypiella in Brazil, 86.

ephippella, Argyresthia.

Epibootania (see Megastigmus).

Epicauta gorhami, measures against, in China, 451.

Epicometis hirta (hirtella), on rape in Czechoslovakia, 468.

Epidiaspis leperii, measures against, on pear in Italy and California, 297, 432; temperature affecting tar distillates efficiency of against, 467.

Epilachna varivestis (Mexican Bean Beetle), in U.S.A., 69, 421, 595, 599; application of bioclimatics to study of, 327; measures against, 421, 595; tests of insecticides on, 69, 78.

Epilachna vigintioctomaculata, on tobacco in Japan, 210; bionomics of, in Korea, 362; on potato in Manchuria, 340, 361.

Epilachna vigintioctopunctata, on solanaceous plants in Fiji, 641.

Epimadiza hirta, on gladiolus in S. Rhodesia, 125.

Epimegastigmus (see Megastigmus). Epinotia meritana, in Br. Columbia, 568; outbreak of, on Abies in

Utah, 567; on pine, 568.

Epitetranychus (see Tetranychus).

epitricis, Perilitus. Epitrimerus myersi, sp. n., on

avocado in California, 288.

Epitrimerus pirifoliae, sp. n., on

pear in California, 33.

Epitrix cucumeris, measures against, on potato and tobacco in U.S.A., 70, 71, 177, 342; bionomics of, 342.

Epitrix parvula, on egg-plant in Jamaica, 239; on tobacco and potato in U.S.A., 254, 342, 380; bionomics and control of, 342,

Epitrix subcrinita, not transmitting potato viruses in Oregon, 84.

Epiurus (see Pimpla).

Epsom Salts (see Magnesium Sulphate).

equestris, Haplodiplosis (Clinodiplosis); Lygaeus (Spilostethus); Merodon.

Eragrostis, relation of Locustana pardalina to, in S. Africa, 394; Nymphula depunctalis on, in Philippines, 364.

Erannis spp., in Crete, 171, 172; in Poland, 438; bionomics of, 172, 438; measures against, on olive, 172.

Evannis auvantiavia, 438.

Erannis bajaria, sorditaria not considered a form of, 171, 172.

Erannis defoliaria, 438.

Erannis leucophaearia, sorditaria considered a form of, 172.

Eranthemum, Doleschallia bisaltide on, in Fiji, 641.

eremita, Conomorium.

erichsoni, Pristiphora (Lygaeonematus, Nematus).

eridania, Xylomyges (Prodenia). Eriobotrya japonica (see Loquat).

Eriobotrya japonica (see Loquat). Eriocampoides (see Caliroa).

Erionota thrax, on coconut and banana in Philippines, 451. Eriophyes, on cotton in India, 128.

Eriophyes avellanae, on hazel in Greece, 452.

Eriophyes carinatus, on tea in India, 128.

Eriophyes fici, on figs in California, 34; question of identity of, 34.

Eriophyes ficus, E. fici possibly identical with, 34.

Eriophyes hibisci, on Hibiscus in Fiji, 641.

Eriophyes lycopersici, sp. n., on tomato in Morocco, 636.

Eriophyes pyri, bionomics and control of, on pear in California, 409.

Eriophyes ribis, measures against, on black currant in Britain, 2.

Eriophyes sheldoni, measures against, on Citrus in California, **55**. Eriophyes vitis (on grape-vine),

measures against, in Australia, 164, 365; in Russian Union, 403. Eriophyids, key to subfamilies and

genera of, 288.

Eriosoma lanigerum (Woolly Apple Aphis), in Britain, 236, 237; in Cyprus, 271; in Denmark, 117; in Egypt, 127; resistance of, to abnormal cold in France, 622; in Germany, 117; in India, 577; in Jugoslavia, 10; in Manchuria, 340; bionomics of, in New Zealand, 548-550; in Palestine, 127; in Poland, 162; in S. Rhodesia, 125; in Russian Union, 588; in Tasmania, 20; in Virginia, 141; susceptibility of varieties of apple to, 100, 141; utilisation of Aphelinus mali against, 10, 20, 117, 127, 163, 271, 548-550, 588; spraying against, 20, 236, 237, 549, 550.

Eriosoma lanuginosum, on elm in Tasmania, 20; not recorded on

pear, 20.

Eriosoma ulmi japonicum, susceptibility of varieties of elm to, in Japan, 443.

Eritrea (see Africa, Italian East). Erromenus simplex, parasite of spruce sawflies in Czechoslovakia, 518.

eruditus, Cheyletus.

Erythmelus psallidis, parasite of Psallus seriatus in U.S.A., 381, 538; bionomics of, 538.

erythrocephala, Acantholyda.

erythrogaster, Campsomeris; Stylocryptus.

Erythroneura pallidifrons, parasite and control of, on greenhouse plants in Britain, 115.

Erythroneura parvula, on vines in Russian Union, 403.

Erythroxylon coca, Linoclostis gonatias on, in Formosa, 447; Eloria noyesi on, in Peru, 165.

Estigmene acraea, food-plants of, in Oklahoma, **340.**

Estola attenuata, on egg-plant in Jamaica, **239.**

Ether, as a solvent, 142.

ethlius, Calpodes.

Ethyl Alcohol (see Alcohol).

Ethylene Dichloride, emulsions of, against Aegeria exitiosa Naupactus leucoloma, 255, 378, 526; and carbon tetrachloride, against pests of stored cereals and plant material, 120, 184, 341, 378, 539; sometimes causing injury to peach, 256; other fumigants compared with, 256, 539.

Ethylene Oxide (as a fumigant), against pests of chestnuts, 359; resistance of strains of Aonidiella aurantii to, 343; curves for expressing toxicity of, to insects, 111; vacuum fumigation with, 359, 605; and carbon dioxide, 359; causing injury to azaleas, 605.

Etiella behri, on Cajanus cajan in

Seychelles, 335.

Etiella zinckenella (on leguminous plants), in Iberian Peninsula, **495**; in India, **129**; in Japan, 556; in Porto Rico, 382, 435, 502; in Russia, 152; in U.S.A., 381; attacking watermelons, 152; parasites and biological control of, 381, 382, 495; alternative host of parasites of lac predators, 129; measures against, 435.

Euacanthus interruptus, on hops in

Britain, 333.

Euarabis (see Arabis).

Euaresta aequalis, introduced into Australia against Xanthium pungens, 546.

Eublemma, damaging teak seed in Burma, 693.

Eublemma amabilis (predacious on Laccifer lacca), parasites and biological control of, in India, 128, 129, 575; longevity of, 128.

Eucalyptus, pests of, in Nyasaland, 126; Coleoptera in timber of, in S. Africa, 60; termite in wharf timber of, in Fiji, 212.

Eucalyptus Oil, spraying with, against Erythroneura pallidifrons, 115.

Eucelatoria comosa, parasite of cutworms in Florida, 657.

Euchlora viridis (see Anomala).

Eucosma ocellana (see Spilonota).

schistaceana, natural Eucosma enemies and biological control of, on sugar-cane in Formosa, 363, 364, 447.

Eudeleboea lopezi, parasite of Cydia molesta in Argentina, 602.

Euderus pempheriphila, parasite of Pempheres affinis in India, 506. Euetheola rugiceps, time of planting maize in relation to, in Arkansas, 178.

Eugenia aromatica (see Clove Tree). Eugenia edulis, Timocratica haywardi on, in Argentina, 188.

Eugenia jambolana, Xylocopa spp. in timber of, in India, 337.

Eugenia uniflora (micheli), Dacus passiflorae on, in Fiji, 212.

Eulecanium (see Lecanium). Eulia citrana, in California, 649; dinitro-o-cyclohexylphenol toxic to, **649.**

Eulia mariana, measures against, on apple in Nova Scotia, 662.

Eulia politana, parasites of, ramie in Transcaucasia, 587.

Eulimneria rufifemur, olfactory responses of, to food-plant of host in Britain, 391.

Eulophus, parasite of Sylepta derogata in Japan, 557.

Euonymus, Aphis fabae on, Britain and Germany, 674, 686.

Eupelmella pedatoria, sp. n., parasite of weevils in India, 506.

Eupelmus, hosts of in Japan, 211,

Eupelmus allyni, emergence holes of, indicating infestation of wheat by Mayetiola destructor in Kansas,

Eupelmus flavicallus, bionomics of, in Australia, 545.

Eupelmus terminaliae, sp. n., parasite of Lasiocampid in India, 385. Eupelmus urozonus, parasite of Pempheres affinis in India, 506.

Eupeodes volucris (predacious on Macrosiphum onobrychis), duration of early stages of, in Utah,

Euphorantha dysderci (see Hyalomyia).

Euphranta, parasites of, on cherry in Japan, 338, 620.

Euphyllura olivina, on olive in Cyprus, 270.

Euplectrus, parasite of Laphygma exempta in Nyasaland, 126.

Euplectrus agaristae (parasite of Phalaenoides glycinae), morphology and bionomics of, in New South Wales, 293.

Euplectrus platyhypenae, 126.

euploeae, Brachymeria.

Eupristina verticillata (pollinating Ficus retusa), introduction of, into Hawaii, 367.

Euproctis chrysorrhoea, auct. (see Nygmia phaeorrhoea).

Eupteromalus, parasite of Nygmia phaeorrhoea in Holland, 205.

Eupteromalus nidulans, parasite of Hormius in Morocco, 193.

Eupteromalus viridescens, parasite of Apanteles congregatus in Tennessee, 254.

Eurhizococcus brasiliensis, on vines in Brazil, 34.

Europe, Tetranychid mites of, 301.

Eurydema, on crucifers in Russian Union, 439.

Eurydema oleraceum, ecology of, attacking cereals in Germany, 631.

Eurydinota lividicorpus, parasite of Coleophora pruniella in Oregon, 643.

Eurydinoteloides gahani, sp. n., parasite of Tyloderma brassicae in Brazil, 183.

Eurydinoteloides montei, sp. n., parasite of Callabismodes tabaci in Brazil, 183.

Eurygaster (on cereals), in Germany, 352; in Poland, 403; type of injury by, 352, 554; measures against, 353, 555.

Eurygaster integriceps, on cereals in Russia, 554; parasite of, 555.

Eurygaster maura, bionomics of, on cereals and grasses in Germany, 63, 64, 632.

Eurytoma denticoxa (see E. nesiotes). Eurytoma fellis (Citrus Gall Wasp), Torymid parasites of, in New South Wales, **564–566**.

Eurytoma monemae, synonymy of, 434.

Eurytoma nesiotes, synonymy of, 434.

Eurytoma parasae (see E. monemae). Eurytoma pistaciae, bionomics of, on pistachio in Greece, **452**, **453**.

Eurytoma promecothecae, sp. n., parasite of Promecotheca papuana in New Guinea Territory, **506.**

Eurytoma saliciperdae, possibly parasitic on Rhabdophaga saliciperda in Britain, 628.

Eurytoma samsonovi, on fruit trees in India, 503.

Euscepes postfasciatus (batatae), synonymy of, **595**.

Euschistus impictiventris; and boll rots in cotton in Arizona, 416, 417.

Eusemion, key to species of, 40.

Eutachina mungomeryi, sp. n., parasite of Laphygma exempta in Queensland, 263.

Eutane terminalis, on grasses in New South Wales, 329.

Eutelia blandiatrix, on Altingia excelsa in Java, 227.

Eutelus, parasite of Phytomyza ilicis in Britain, 669.

Eutelus kojimae, sp. n., parasite of Dendrolimus spectabilis in Japan, 91.

Eutelus subfumatus, parasite of Diprion spp. in Poland, 232, 671.

Eutelus tabatae, sp. n., parasite of Dendrolimus albolineatus in Sakhalin, 91.

Eutermes, in Australia, 545.

Eutermes costalis, bionomics and control of, damaging plants and timber in Porto Rico, 464, 465.
Eutermes exitiosus, in Australia,

Eutermes exitiosus, in Australia, 89, 282, 545; methods of testing resistance of timber to, 545; environment in mounds of, 89, 282.

Eutermes olidus, in Fiji, 641.

Eutermes ripperti, damaging damp timber in Jamaica, 239.

Eutetrapha metallescens, on Quercus mongolica in Russian Far East, 42.

Eutetrapha sedecimpunctata, on Tilia amurensis in Russian Far East, 42.

Eutettix tenellus (Beet Leafhopper), in U.S.A., 286, 409, 412, 593, 596; experiments with curly-top of beet and, 286, 412; on spinach, 594; breeding areas, ecology and migrations of, 286, 593; new varieties of Pipunculus parasitic on, 596; measures against, 287, 412.

Eutochia lateralis, on potato in Hawaii, 367.

Eutrichopoda abdominalis, parasite of Dysdercus in Brazil, 22.

Eutrombidium locustarum, attacking grasshoppers in U.S.A., 75, 376; bionomics of, 376.

Euxesta stigmatias, measures against; on maize in Porto Rico, 328; bionomics and distribution of, 328.

Euxoa detersa personata, baits for, on beans in Wisconsin, 32.

Euxoa messoria, bait for, on tobacco in Connecticut, 178.
Euxoa segetum (see Agrotis).
Euzophera punicaeella, bionomics of, in Baluchistan, 51.
evanescens, Trichogramma.
Evergestis rimosalis, on cabbase in

Evergestis rimosalis, on cabbage in Louisiana, 250.

Louisiana, **250.**Evetria (see Rhyacionia).
ewingi, Persectania.
examinator, Pimpla.
exaptus, Systates.
excarinata, Brachymeria.

exempta, Laphygma (Spodoptera). Exenterus abruptorius, introduction of, against Diprion polytomum

in Canada and U.S.A., 276, 381.

Exenterus adspersus, introduction of, against Diprion polytomum in Canada and U.S.A., 369, 381.

Exenterus laricinus, parasite of Diprion similis in Poland, 671. Exenterus marginatorius, parasite of

Diprion similis in Poland, 671. Exeristes (see Pimpla).

Exeristesoides (see Pimpla). exigua, Hyposoter; Laphygma. exitiosa, Aegeria (Conopia). exitiosus, Eutermes.

Exochomus flavipes, attacking Coccids in Seychelles, 335; establishment of, in Mahé, 335.

Exochomus quadripustulatus, predacious on Chermes spp. in Britain, 262, 263; proposed introduction of, into Australia, 263.

Exochomus ventralis, establishment of, against Coccids in Seychelles, 335; Technomyrmex hindering breeding of, 335.

Exora apicipennis, on Crotalaria juncea in S. Rhodesia, 125.

Exora discoidalis, food-plants of, in S. Rhodesia, 124, 125.

Exorista (see Zenillia).

expansa, Anomala. Experimental Data, analysis of, 65, 66, 240, 242, 243.

F.

fabae, Aphis (Doralis); Empoasca.
fabalis, Empoasca.
fabia, Earias.
fabricator, Ichneumon.
facialis, Catascopus.
fagi, Contarinia; Phyllaphis.
fagicola, Dasyneura.
Fall Cankerworm (see Alsophila pometaria).

Fall Webworm (see Hyphantria

cunea).

farinae, Tyroglyphus.
fascialis, Hymenia (Zinckenia) (see
H. recurvalis).

fasciata, Lophosia.

fasciatipennis, Coloborhombus.

fasciatus, Aeolothrips; Dysdercus; Hercothrips (Heliothrips); Silvanus.

fasciculatus, Araecerus. fastuosa, Chrysomela. fatigans, Culex. fedorovi, Taeniothrips. fellis, Eurytoma.

Feltia annexa, on cabbage and sugar-cane in U.S.A., 250, 657; parasites of, 657.

Feltia ducens, measures against, on

lucerne in Wisconsin, 32.

Feltia malefida, parasites of, on sugar-cane in Florida, 657.

Feltia segetum (see Agrotis). femoralis, Creontiades; Tychius. femorata, Chrysobothris.

femur-rubrum, Melanoplus.

fenestralis, Angitia.

Fennel (Foeniculum), new Aphid on, in California, 590; Thrips tabaci on, in Crimea, 43.

Fenusa pusilla, weight of foliage in relation to, 143.

ferganensis, Rhynchites auratus. Ferns, Calopistria floridensis on, in greenhouses in Canada and U.S.A., **592**.

Ferrisiana virgata, new parasite of, on coffee in Kenya, 40.

Ferrous (Iron) Sulphate, uses of, in sprays, 460, 662, 681; increasing risk of injury to *Citrus* by limesulphur, 460.

ferruginea-striata, Aphis.

ferrugineus, Dacus (Chaetodacus, Strumeta); Laemophloeus.

ferus, Nabis.

Fescue (see Festuca). festina, Stictocephala.

festiva, Chlorida; Poecilonota.

Festuca spp., new Cecidomyiid on, in Britain, 635; Pentatomid cereal pests on, in Germany, 63, 631; Pediculopsis graminum on, in Russian Union, 151.

festucae, Dasyneura.

fici, Eriophyes; Tetranychus.

ficifoliae, Rhyncaphytoptus.

Ficus, new Eriophyid on, in California, 288; Coccids on, in Egypt, 509.

Ficus benjamina, Aleurodids on, in Trinidad, **25.**

Ficus carica (see Fig).

Ficus retusa, Anomala expansa on, in Formosa, 363; Agaonid introduced into Hawaii to pollinate, 367.

ficus, Chrysomphalus; Eriophyes. Field Plots, co-ordination of experiments in, 240, 242, 245, 340, 417.

Fig, mite on, in California, 34; Ceratitis capitata on, in France, 195; Tetranychus fici on, in India, 128.

Fig (Fruit), Lonchaea aristella intercepted in, in Russian Union, 440; (dried), pests in, in Germany, 201, 230; uses of, in baits for Trypetids, 98, 270.

figulilella, Ephestia.

Fiji, miscellaneous pests in, 212, 213, 366, 641; bunchy-top of banana in, 213; natural enemies and biological control of insect pests in, 19, 22, 212, 213, 366, 605, 641; introduction of insect enemies of nutgrass into, 367; introduction of beneficial insects into other countries from, 212, 238, 365, 366, 641.

fijiensis, Biosteres (Opius). filamenta, Empoasca. filamentosus, Pseudococcus. fimetarius, Phalacrus.

Finland, thrips on Alopecurus in, 160; Diarthronomyia on chrysanthemum in, 673; Melolontha hippocastani in, 267; Pissodes spp. and their parasites in, 8; Coleopterous fauna of potato storage cellars in, 392; Indian Coleoptera in timber from 465; entomological publications in, 694.

Fir, Douglas (see *Pseudotsuga taxi-folia*).

Fir, Silver (see Abies).

fiscellaria, Ellopia.

Fish, destroying Nymphula vittalis in Korea, 619; insecticides toxic to, 3, 136, 514; (dried or salt), Coleoptera feeding on, 513, 584.

Fish Oil, as an adhesive and spreader for sprays, 52, 53, 74, 177, 189, 380, 383, 430, 525, 540, 645; as an adhesive for cryolite dips, 652; emulsions of, unsatisfactory against whiteflies, 530; formulae containing, 52, 53, 74, 177, 380, 383, 430, 530, 540; types of, 430, 525.

Fish-oil Soap (see Soap, Fish-oil). flabellicornis, Hylecoetus (Hylecerus). Flamboyant (see Poinciana).

Flame-throwers, against grasshoppers and locusts, **214**, **479**; suggested against *Rhytidoderes* plicatus, **621**.

flammea, Panolis.

flandersi, Tropidophryne.

flava, Cosmophila (Anomis);

Hoplocampa; Sipha. flaveola, Scaptomyza.

flavescens; Empoasca (Chlorita); Monema (Cnidocampa); Phalera.

flavicallus, Eupelmus. flavicauda, Sphaerophoria. flavicornis, Sphegigaster.

flavipes, Apanteles; Apion; Exo-

chomus.

flaviscutellaris, Cyrtomorpha. flavisignatum, Monalonion.

flaviventris, Cremastus; Neurotoma.

flavolineata, Metriona. flavolineatus, Bolax.

flavo-orbitalis, Cremastus.

flavus, Tychius.

(Linseed), grasshopper damaging, in Argentina, 214; Plusia orichalcea on, in Bengal, 577; Thrips spp. on, in Holland, 233, 234, 465; survey of pests of, in Poland, 350; effect of infestation by flea-beetles on yield of, in Russia, 684; pests of, in U.S.A., 180, 652; possible translocation of derris in, 234; pests in stored seed of, 173, 319, 521; insects infesting house insulation material containing,

Flea-beetle, Hemp (see *Psylliodes* attenuata).

Flea-hopper, Cotton (see Psallus seriatus).

Fleas, grain mites spread by, in Russian Union, **552**.

fletcheri, Opius.

Flit, tests of penetration of, into tracheae of insects, 580.

floccifera, Pulvinaria.

floralis, Hylemyia (Chortophila); Nemorilla.

floricolus, Tarsonemus.

Florida, pests of Citrus in, 256, 257, 258, 259; cotton pests in, 242, 247; Anticarsia gemmatalis on soy beans in, 371; sugar-cane pests in, 656-659; Aphids transmitting sugar-cane mosaic in, 658; tobacco pests in, 251, 599; vegetable pests in, 249, 287, 600; legislation against Naupactus spp. in, 275; beneficial insects and biological control in, 143, 251, 261, 461, 656, 657, 658, 659;

parasite of Apanteles congregatus in, 254; climatic conditions unfavourable to Ceratitis capitata in, 327.

floridanus, Horismenus. floridensis, Callopistria.

Flour, pests in, and their control, 27, 52, 64, 65, 173, 175, 223, 234, 319, 350, 378, 399, 400, 513, 584, 612, 640, 685; nutritive value of types of, for pests, 64, 65, 173, 175; method for determining movement of Tribolium confusum in, 52; effect of injury to grain by Pentatomids on quality of, 353,554; effect of grain fumigants on, 319; in baits, 127, 240; as carrier and adhesive for dusts, 185, 367, 371, 674; in repellent paint against Chrysobothris, 420. Flour Beetle (see Tribolium).

Flour Mills, pests in, and their control, 8, 162, 400, 685.

Flour Mite (see Tyroglyphus farinae). Flour Moth (see Ephestia kuehniella).

niella).
Flower Thrips (see Frankliniella moultoni).

fluctuata, Cidaria.

Fluorine Compounds, problems of residues of, on fruit and vegetables, 104, 432, 595, 648; penetration of leaves by, 309; review of use of, as insecticides, 456.

Foeniculum (see Fennel).

Foliage, weight of, in relation to insect control, 143.

foliata, Membracis.

Fomes annosus, 504.

forbesi, Cerosipha (Aphis). Forest Ant, Red (see Formica rufa). Forest Entomology, book on, **527**; recent literature on, **160**.

Forest Tent Caterpillar (see Mala-

cosoma disstria).

Forests, pests of, in Algeria, 226; pests of, in Australia, 20, 21, 185, 262, 545, 546, 602; Galerucella luteola in, in Austria, 324; pests of, in Britain, 58, 65, 262, 295, 391; pests of, in Burma, 612, 692, 693; pests of, in Canada, 25, 53, 275, 276, 368, 369, 381, 382, 463, 540, 647, 662–664; Longicorns in, in Ceylon, 612; pests of, in China, 360, 449, 451, 614; new Coccid in, in Corsica, 235; pests of, in Czechoslovakia, 468, 469, 517, 518, 564; Diprion pini in, in Danzig Territory, 105, 397; new Matsucoccus in, in

Dominican Republic, 392: pests of, in Finland, 8, 267; pests of, in France, 225, 226, 235, 324; pests of, in Germany, 7, 48, 65, 98, 101, 105, 170, 204, 216–218, 230, 231, 298, 299, 334, 335, 353, 354, 397, 398, 399, 400, 401, 466, 512, 521, 522, 563, 579, 580, 581, 507, a suggestions for protection 607; suggestions for protection of Formica rufa in, in Germany, 231; pests of, in Holland, 269, 396; pests of, in India, 224, 612; pests of, in Netherlands Indies, **227, 385, 577, 578;** comparison between gambier plantations and, in relation to pests in Sumatra, 385; pests of, in Italy and Sicily, 516, 522; pests of, in Japan, 19, 91, 209, 211, 339, 441, 443, 615; Dioryctria peyerimhoffi in, in Morocco, 58; Hylastes ater in, in New Zealand, 504; pests of, in Nyasaland, 126; pests of, in Poland, 9, 162, 350, 398, 399, 401, 438, 670, 676; pests of, in Rumania, 355; pests of, in Russian Union, 42, 43, 147, 153, 456, 474, 680, 681, 43, 147, 153, 456, 474, 680, 681, 682; pests of, in Sakhalin, 19, 91, 339, 615; pests of, in Sweden, 122, 123; pests of, in Switzerland, 169, 517; pests of, in Turkey, 218; pests of, in U.S.A., 28, 32, 53, 67, 68, 69, 73, 74, 77, 78, 132, 138, 140, 144, 170, 176, 177, 190, 279, 284, 324, 329, 342, 368, 369, 374, 377, 378, 382, 392, 418, 420, 427, 429, 434, 461, 498, 499, 528, 529, 530, 532, 533, 540, 546, 567, 568, 599, 600, 644, 540, 546, 567, 568, 599, 600, 644, **645**, **647**, **649**, **651**; possibilities of aeroplane dusting and biological control in, 170, 514.

Forficula auricularia, in Ontario, 601; bionomics and control of, in Tasmania, 510; plants attacked by, 510, 601; liberation of Bigonicheta setipennis against, in U.S.A., 379.

forficula, Enneadesmus.

Formica, measures against, in Russia, 44.

Formica rufa, utilisation of, in forests in Germany and Czechoslovakia, 231, 564.

formicarius, Cylas.

Formosa, fruit pests in, 205, 206, 363, 557; miscellaneous pests in, 363, 558, 618; sugar-cane pests in, 206, 339, 363, 364, 444, 446, 447; soil insects in sugar-cane fields in, 206, 446; pests of tea

in, 339, 340, 447; ants of, 446; Aphids producing tannin galls in, 205; new Coccid on Arundinaria in, 103; locusts in, 486; Bruchus rufimanus not occurring in, 208; beneficial insects and biological control in, 205, 363, 364, 447, 558; parasite of Coccinellids in, Rodolia cardinalis introduced into Japan from, 446; risk of introduction of Dacus spp. into Japan from, 615.

formosa, Encarsia.

formosae, Aserica (Autoserica).

formosana, Gryllotalpa.

formosanus, Cardiophorus (Platy-nychus); Coptotermes. fornicatus, Xyleborus.

Fornicia, parasite of Setora nitens in Java, 347.

foveolata, Tyloderma.

Fowls, destroying noxious insects, 253, 377, 557, 608.

fractus, Melanoxanthus.

fragariae, Capitophorus; Taysonemus (see T. pallidus); Tyloderma.

fragilis, Malacosoma.

Cecidomyiid on Aira France, flexuosa in, 635; weevils on artichokes in, 357; pests of chestnuts in, 358; Phloeosinus spp. on conifers in, 225, 226; pests of crucifers in, 167, 458, 459, 621, 622: Pseudococcus citri on orange in, 266; orchard pests in, 168, 195, 357, 390, 622, 689; pests of ornamental plants in, 268, 623, 689; vine moths in, 155; Carpophilus hemipterus in, 167; new Coccids in, 103, 235; Corynetines of, 584; Leptinotarsa decemlineata in, 106, 107, 169, 194, 349, 350, 357, 583, 584, 622, 634, 675; beneficial insects and biological control in, 107, 194, 266, 629; certification of insecticides in, 675: introduction of beneficial insects into other countries from, 282, 324; legislation against introduction of pests into other countries from, 11, 356, 358.

frangulae, Aphis (Doralis). Frankliniella, transmitting disease of tobacco in S. Africa, 360.

Frankliniella fusca, food-plants of, in U.S.A., 177, 372; measures against, 177.

Frankliniella insularis, measures against, on banana in Haiti, 289; transmission of spotted wilt by, 546.

Frankliniella intonsa, bionomics of, on wheat in Sweden, 121.

Frankliniella moultoni, on Citrus in California, 433; characters dis-Scirtothrips tinguishing and, 433.

Frankliniella tenuicornis, bionomics of, on wheat in Sweden, 121.

Frankliniella tritici, food-plants of, in U.S.A., 142, 372, 413; effect of, on yield of cotton, 413; trapped by leaves of tomato, 142.

fraterculus, Anastrepha.

Fraxinus (see Ash).

frenchi, Coptotermes; Lepidiota. frischi, Anomala dubia; Dermestes.

frit, Oscinella (Oscinis).

froggatti, Haplothrips; Typhlocyba. Frogs, destroying noxious insects, 407, 452; possibility of utilising, in Philippines, 452; damaging

trap-ditches in beet fields, 407.

frontalis, Dendroctonus. frugiperda, Laphygma.

Fruit (Dried), pests in, and their control, 64, 154, 167, 173, 201, 225, 230, 521, 584.

Fruit-flies (see Trypetids).

Fruit-fly, Cherry (see Rhagoletis cerasi).

Fruit-fly, Mediterranean (see Ceratitis capitata).

Fruit-fly, Melon (see Dacus cucurbitae).

Fruit-fly, Olive (see Dacus oleae). Fruit-fly, Queensland (see Dacus ferrugineus tryoni).

frutetorum, Diprion.

fullo, Polyphylla.

fulvicornis, Hoplocampa (see H. minuta).

fulvipes, Ichneutes.

fulvoguttata, Melanophila.

fulvum, Stromatium.

fumiferana, Harmologa.

Fumigants, methods of testing and comparing, 111; new type of, 198, 199; types of equipment for using, 53, 54, 174, 260, 272, 605.

Fundella cistipennis, measures against, on lima beans in Porto Rico, 435.

Cvdia funebrana, (Grapholitha, Laspeyresia).

funestus, Lagochirus.

Fungi, Tinea secalella on dried, in Germany, 220; list of animals associated with edible, 511. (See Mushrooms.)

75, 86, 91, 92, 100, 105, 118, 144, 173, 258, 300, 302, 306, 307, 308, 333, 368, 392, 427, 455, 470, 471, 496, 501, 504, 578, 587, 608, 625, 629, 660, 666, 677, 684, 688. Fungi, Injurious, 20, 71, 72, 101, 124, 138, 164, 181, 204, 219, 269, 348, 358, 359, 422, 425, 427, 437, 491, 495, 504, 522, 541, 544, 562, 568, 584, 628, 630, 664, 693; relation of insects to, 61, 62, 65, 68, 77, 186, 222, 242, 284, 290, 360, 365, 378, 388, 389, 396, 401, 452, 453, 524, 527, 529, 564, 642, 645, 656, 657; relation of mites to, 79.

Fungi, Entomogenous, 8, 9, 32, 39,

Fungus, Honey (see Armillaria mellea).

Fungus, Red Rot (see Colletotrichum falcatum).

furcatispinus, Pseudococcus.

furcifera, Sogata.

Furfural, synthetic compounds derived from, as insecticides, 593. Furniture, pests in, and their control, **55**, **160**, **181**, **237**, **493**,

Furs, Coleoptera damaging, 47, 196, 513.

Arcyptera (Stethophyma); fusca, Busseola; Frankliniella; Lachnosterna (Phyllophaga).

fuscedinella, Coleophora.

fuscicornis, Bothriophryne; Tremex. fuscilabris, Ceratomegilla.

fuscipenne, Microplectron.

fuscipunctella, Tinea.

fuscispinus, Carpocoris pudicus.

fuscus, Anthrenus.

fusiformis, Byrsocrypta (Tetraneura).

futilis, Lachnosterna (Phyllophaga).

G.

gahani, Eurydinoteloides; Pseudococcus.

galactinus, Microcryptus.

Galactites tomentosa, weevil on, in France, 357.

galactitis, Apion (Ceratapion) carduorum.

Galeruca tanaceti, on horse-radish in Germany, 110.

Galerucella distincta, bionomics of, on strawberry in Korea, 619.

Galerucella luteola (Elm Leaf Beetle), in Austria and France, 324; in Japan, 144; in U.S.A., 132, 144, 324, 427, 499; parasites and biological control of, 132, 324; other measures against, 428.

Galerucella nymphaeae, bionomics and control of, on Nymphaea

alba in Italy, 3.

Galerucella xanthomelaena (see G. luteola).

gallarum, Byrsocrypta.

Galleria mellonella (Wax Moth), parasitised by Dibrachys cavus in Argentina and Germany, 238, 631; bionomics and fertility of, 515; effect of humidity on infection of, by *Beauveria*, 106; utilisation of bacteria isolated from, 455.

gallicus, Doryctes.

Galvanometer, for detecting Sturmia in pupae of Bombyx mori, 442.

Gambier (see *Uncaria gambir*). Gambrus stokesi, parasite of Cydia spp. in Australia, 293, 382, 640; introduced into U.S.A., 382.

Gammaphytoptus camphorae, gen. et sp. n., on Cinnamomum cam-phora in California, 288.

Gangai Disease, 577.

Gardenia, Dialeurodes citri on, in

Virginia, 530.

Gasterocercodes brasiliensis (and var.), bionomics and control of, on cotton and Hibiscus in Brazil, 136.

Gasterocercodes gossypii, G. brasiliensis mistaken for, 136.

Gastrimargus musicus, phases of, in Australia, 545.

gei, auct., Macrosiphum (see M. solanifolii).

Gelatine, spraying with, against Tetranychus telarius, 583.

Gelechia cercerisella, parasites of, on Cercis canadensis in Kansas, 376. gelechiae, Idechthis; Pteromalus (see Dibrachys cavus).

Gelis asozanus, parasite of Apanteles liparidis in Japan, 211.

geminata, Chrysomela; Solenopsis. gemma, Chrysocharis.

gemmatalis, Anticarsia.

Generation, meaning of term, 137.

geniculata, Bucentes. geniculatus, Engytatus;

cephalus. Genista aetnensis, Icerya purchasi on,

in Sicily, **5.**

genitalis, Hylemyia (Adia, Phorbia). Geocoris decoratus, predacious on Macrosiphum onobrychis in Utah, 30.

Georgia (U.S.A.), cotton pests in, 242, 247, 380; Coleoptera infesting pods of leguminous plants in, 76; miscellaneous pests in, 377, 379, 657; peach pests in, 255, 623; Heliothis armigera in, 179; beneficial insects in, 77.

Geraniol, repellent to parasites of Rhvacionia buoliana, 391.

Geranium (Pelargonium), Coccid on, in Japan, 617.

Germany, Aphis fabae on beans in, 334: Pentatomids on cereals in, 63, 352, 353, 631, 632; pests of crucifers in, 110, 511; forest pests in, 6, 7, 48, 65, 98, 101, 105, 204, 216, 218, 223, 230, 231, 298, 334, 335, 353, 354, 397, 398, 400, 401, 466, 512, 521, 563, 579, 580, 581; Contarinia medicaginis on lucerne in, 457: orchard pests in, 46, 99, 112, 113, 114, 155, 157, 170, 297, 298, 300, 352, 398, 467, 513, 632, 633; pests of small fruits in, 100, 114, 155, 457, 466, 467; pests of ornamental plants in, 218, 267, 583; pests of stored products in, 8, 47, 102, 201, 220, 223, 224, 230, 355, 399, 400, 513, 520, 521, 582; Coleoptera damaging building timbers in, 6, 7, 110, 202, 203, 204, 268, 335, 520; vine pests in, 46, 102, 103, 111, 155, 458, 465, 496, 582, 688, 689; Leptinotarsa decemlineata in, 48, 100, 106, 113, 158, 160, 512, 583; Melolontha in, 112, 171, 299, 300, 334, 457, 495, 606 608; Stenomalus muscarum in houses in, 201; new Tettigoniid 109; Tipulids in, 519: miscellaneous pests, 223, 298, 336, 398, 606; insects and virus diseases of plants in, 156, 157, 267, 301, 333, 633, 636, 686–688; pests imported into, 218, 298, 583; beneficial insects in, 8, 64, 105, 113, 170, 217, 218, 231, 334, 335, 352, 354, 398, 400, 520, 580, 581, 608, 630, 632, 687; lists of popular names of insects in, 437; standards for calcium arsenate in, 115; Aphelinus mali introduced into Denmark from, 117; legislation against introduction of L. decemlineata into Britain from, 356.

gestroi, Hispa.

Geum rivale, Pentatomid on, in Germany, 632.

ghesquierei, Pimelephila.

Giant Palm Borer (see Dinapate wrighti).

Giardomyia rhododendri, sp. n., on rhododendron in New York, 499.

gibbicollis, Hadrobregmus.

gibbosus, Ligyrus.

gibbus, Bruchophagus.

giffardi, Dirhinus; Tetrastichus.

giffardianus, Tetrastichus.

gigas, Sirex.

Gilbert Islands, Dynastid on banana and taro in, 366.

gilvus, Macrotermes. Gipsy Moth (see Lymantria dispar). Gladiolus, Epimadiza hirta on, in S. Rhodesia, 125.

glaseri, Neoaplectana.

Glass, Ground, applied to strawberry roots against Lachnosterna, 425.

Glasshouse Leafhopper (see Erythroneura pallidifrons).

globata, Microgaster.

globosa, Cistogaster; Myiophasia; Xystrocera.

gloveri, Lepidosaphes.

Glue, spraying with, against Tetranychus telarius, 583: as an adhesive for cryolite dips, 380, 652.

Glycerine, increasing toxicity of nicotine sulphate to eggs of Chilo simplex, 19; and tartar emetic, against Scirtothrips citri, 425.

glycinae, Phalaenoides.

glycinivorella, Cydia (Grapholitha). Glycyphagus cadaverum, in seeds of flax, etc., in Russian Union, 319.

Glycyphagus destructor (see Lepidoglyphus).

Glycyphagus domesticus, anatomy and development of, 456.

Glypta, parasite of Tortricids in Trancaucasia, 587.

Glypta haesitator, bionomics parasitising Cydia nigricana in Britain, 38.

Glypta rufiscutellaris (parasite of Cydia molesta), introduced into Australia, 544; in U.S.A., 377,

544. Glyptomorpha nicevillei, parasite of

Scirpophaga spp. in Mysore, 291. Glyricidia, Orthezia praelonga on, in Trinidad, 25.

Gmelina arborea, pests of, in Burma, 692.

Gnathocerus, Corynetes coeruleus probably predactious on, in flour, 584.

gnidiella, Cryptoblabes.

Gnorimoschema, Phthorimaea (q.v.) not considered distinct from, 289. Gobaishia japonica, susceptibility of varieties of Ulmus propingua to, in Japan, 443.

Gold Coast, summary of plant

quarantines in, 592. Gold Mines, Micromalthus debilis in timber in, in S. Africa, 60.

Golf Greens (see Turf).

Gomphocerus sibiricus (see Aeropus).

gonatias, Linoclostis.

Gonatopus, parasite of Saccharo-sydne saccharivora in Florida, 657. Gonia cinerascens (see Cnephalia). goniaeformis, Tritaxys (Anamastax, Blepharipeza).

Gonipterus scutellatus, on Eucalyptus in Nyasaland, 126.

Gonocephalum (see Dasus).

Gooseberry, Aphids on, in Britain, 333, 596; Aspidiotus ostreaeformis on, in Czechoslovakia, 468; Nematus ribesii on, in Germany, 114; pests of, in U.S.A., 178, 179, 596, 597; Dasyneura ribicola confined to, 673.

gorhami, Epicauta.

Gortyna immanis (see Hydroecia). Goryphus javanicus, parasite Setora nitens in Java, 347.

Gossyparia spuria, measures against, on elm in California, 649.

gossypiella, Platyedra (Pectinophora). gossypii, Aphis; Bucculatrix; Corythuca; Gasterocercodes; Macrosiphum (Acyrthosiphon); Phenacoccus.

gossypiperda, Bemisia. Gossypium (see Cotton). gracilicorne, Megymenum.

gracilis, Orthomorpha; Polygraphus.

gradarius, Ichneumon. Grain Noctuid (see Trachea basilinea).

Grain Weevil (see Calandra granaria).

Grain Wireworm, Northern (see

Corymbites aeripennis destructor). Gram (see Cicer arietinum).

graminis, Charaeas; Dasyneura. graminum, Pediculopsis; Steno-Toxoptera (Rhopalothrips; siphum).

Granadilla, Dacus xanthodes on, in Fiji, 212.

granaria, Calandra (Sitophilus). granarium, Macrosiphum (see M. avenae); Trogoderma.

grandis, Anthonomus; Harmolita.

granella, Tinea.

Grape Berry Moths (see Polychrosis botrana and P. viteana).

Grapefruit, Orthezia praelonga on, in Grenada, 25; pests of, in U.S.A., 55, 256.

Grapes, problem of arsenical residues on, 191, 495; juice of, in baits for Trypetids, 270.

Graphium ulmi (see Ophiostoma).

Grapholitha (see Cydia).

Graptolitha antennata, measures against, on apple in Quebec, 664. Grass, Bent (see Agrostis).

Couch (see Agropyrum repens).

Grass, Red-top (see Agrostis vulgaris).

Grass, Rye (see Lolium).

Grass, Yellow Cat (see Trisetum

flavescens).

Grasses, pests of, in Australia, 292, 329, 389, 603; Cecidomyiids on, in Austria, 635; pests of, in Britain, 122, 635; Chirothrips hamatus on, in Finland, 160; Cecidomyiids on, in France, 635; pests of, in Germany, 63, 519; Contarinia tritici on, in Italy, 3; Phytomyza nigra on, in Japan, 208; Laphygma exempta on, in Kenya, 666; planted among clover against Smynthurus viridis in New Zealand, 550; pests of, in Russian Union, 151, 312, 319, 680; Thysanoptera on, Sweden, 121; pests of, in U.S.A., 248, 374, 500, 527, 635; causes of white ear in, 121, 151, 680; cellular structure of, as a protection against Thysanoptera, 121; relations of locusts and grasshoppers to, 12, 178, 214, 321, 387, 388, 394, 477, 479, 483, 618; cereal and sugar-cane pests on, 88, 122, 208, 268, 363, 364, 537, 547, 555, 577, 620, 631, 632; sugar-cane mosaic in, 88; mite in stored seeds of, 319; as base for baits for locusts, 478. Turf.)

Two-striped Grasshopper, (see Melanoplus bivittatus).

Grasshoppers (see Locusts).

Cryptorrhynchus gravis, (Sternochetus).

grayi, Papilio scamander.

Greece, miscellaneous pests in, 452, 453; locusts in, 478; summary of plant quarantines in, 592.

Greenhouse Fumigation, 115, 272, 357, 526.

Greenhouse Stone Cricket (see Tachycines asynamorus).

Greenhouse Whitefly (see Trialeurodes vaporariorum).

greeni, Microbracon.

gregaria, Schistocerca.

Gregarina acridiorum, in Schistocerca gregaria, 486.

Gregarines, in Melolontha, 496.

Grenada, miscellaneous pests in, 25; termite in timber in, 56, 384; Dasyscapus parvipennis introduced into, against cacao thrips, 25.

grisella, Achroia (Meliphora).

griseola, Leucopis.

griseus, Harpalus (Pseudophonus); Hyperodes.

grossulariae, Aphis.

Ground-nut Oil (see Oil, Ground-

nut).

Ground-nuts, Aphis laburni and rosette disease of, in Nyasaland, 126; Dysdercus superstitiosus on, in Sudan, 573.

Ground-nuts (Stored), nutritive value of, for *Ephestia* and *Plodia*, **64**, **173**.

Ground-nuts, Bambarra (see *Voandzeia subterranea*).

Groundsel (see Senecio vulgaris). Gryllica melzeri, sp. n., on Coccoloba ilheense in Brazil, 290.

Gryllotalpa, egg-chambers and

oviposition of, 509.

Gryllotalpa africana, in Egypt, 509; on sugar-cane in Formosa, 363; on tobacco in Japan, 210; in Manchuria, 361; bionomics of, 363, 509.

Gryllotalpa formosana, on sugarcane in Formosa, 206, 363;

bionomics of, 206.

Gryllotalpa gryllotalpa (vulgaris), on barley in Czechoslovakia, 469; in Egypt, 509; on horse-radish in Germany, 110.

Gryllotalpa gryllotalpa f. cophta, in

Egypt, **509**.

Gryllulus (Gryllus) assimilis var. pennsylvanicus, bionomics of, on cotton in Louisiana, **341.**

Guadeloupe, virus disease in banana in, 289; parasite of *Leucoptera coffeella* in, and its introduction into Porto Rico, 435.

Guaiacum, resistance of timber of, to termites, 465.

Guava, Coccids on, in Egypt, 509; Coccus viridis on, in Haiti, 84; Dacus psidii on, in Pacific Islands, 131, 294; Helopeltis bergrothi on, in Sudan, 574; Euxesta stigmatias on, 328.

guddmannella, Phthorimaea (Gnori-

moschema).

Guiana, British, utilisation of Metagonistylum minense against Diatraea in, 505, 524; miscellaneous pests in, 524; Larra americana not found in, 85; beetles boring in rum puncheons exported from, 328, 524; M. minense introduced into St. Lucia from, 182.

Guiana, Dutch, Atta spp. in, 567;

Larra americana in, **85.** guineense, Tetramorium.

gularis, Aphomia (Paralispa).

Gulls, destroying Melolontha in Germany, 608.

guppyi, Tomaspis. gurneyi, Coccophagus. guttulosa, Austracris. gyllenhali, Pissodes.

Gymnoascus ruber, infesting Cydia nigricana in Britain, 39.

Gymnosoma rotundatum, parasite of Pentatomids in Germany, **64.** Gypsonoma minutana (see Enar-

monia).

Gypsum, as a carrier for dusts, 32, 463.

Gyranusa citrina, sp. n., parasite of Coccids in Kenya, **39.**

H.

Habrobracon (see Microbracon).

Habrolepis dalmanni, utilisation of, against Asterolecanium variolosum in Australia and New Zealand, 21, 546.

Hadrobregmus gibbicollis, in building timber and trees in U.S.A., 499.

haematodes, Tibicen.

haematopus, Penthaleus (see P. major).

Haemoglobin, affecting oil deposits of sprays, 411.

haemorrhoidalis, Athous (see A. obscurus); Heliothrips; Omocestus (Stenobothrus).

haesitator, Glypta.

Haiti, pests and diseases of banana in, 288, 289; miscellaneous pests in, 84, 85, 661; Tiphia hispaniolae in, 84; Larra americana apparently not occuring in, 85.

haitiensis, Tetrastichus.

halli, Sturmia.

Halotydeus destructor, bionomics and control of, in Australia, 131. 291, 545.

Haltichella, parasite of Gelechia cercerisella in Kansas, 376.

Halticoptera, parasite of Euphranta in Japan, 620.

Ham, Necrobia rufipes feeding on, **584**.

hamatus, Chirothrips.

Hambletonia pseudococcina, intro-duced into Porto Rico against Pseudococcus brevipes, 502.

Hamelia patens, Monalonion bahiense on, in Brazil, 566.

Hamitermes (see Amitermes).

hampei, Stephanoderes.

Hapalia machaeralis, parasites and biological control of, on teak in India and Burma, 263, 612, 693. Haplodiplosis equestris, on barley

in Czechoslovakia, 468.

Haplothrips aculeatus, in Holland, 269; in Italy, 472; bionomics of, in Sweden, 121; on wheat, 121, 269; doubtful vector of virus disease of sugar sorghum, 472.

Haplothrips distinguendus, bionomics of, on wheat in Sweden, 121. Haplothrips froggatti, on vines in

Queensland, 365, 603; measures

against, 365.

Haplothrips tritici, on wheat in Russian Union, 453, 554; analysis of injury by, 554. harcyniae, Pissodes.

Harlequin Cabbage Bug (see Mur-

gantia histrionica).

Harmolita grandis, on wheat in 280, 375; U.S.A., measures against, 280.

Harmologa fumiferana (Spruce Budworm), strain of, on Pinus ponderosa in Colorado, 377.

Harpalus griseus, measures against, on strawberry in Germany, 457.

Harpalus rufipes (ruficornis) (on strawberry), in Denmark, 117; measures against, in Germany, 457, 466.

Harpobittacus australis, preventing establishment of Tyria jacobaeae

in Australia, 282.

Harvesting Machines, destruction of insects in, 315.

hattorii, Cinara.

Hawaii, miscellaneous pests in, 366, 367; sugar-cane pests in, 90, 367, 368; Tarsoneminae of, 224; termites in, 180, 385; biological control of noxious insects and

plants in, 90, 366, 367, 368; establishment of Agaonid pollinating Ficus retusa in, 367; insects intercepted in aeroplanes in, 368; introduction of beneficial insects into other countries from, 367, 382, 502, 546; pests from, intercepted in U.S.A., 461; legislation dealing with treatments of fruit, etc., for export to U.S.A. from, **344.**

Hawthorn (see Grataegus).

Hawthorn Stem Midge (see Thomasiniana crataegi).

haywardi, Timocratica.

Hazel, pests of, in Greece, 452; Erranis spp. on, in Poland, 438; Lachnosterna hirticula on,

Wisconsin, 32.

Heat, treatment of lily bulbs with, against mites, 419, 617; against pests of seeds and stored products, 102, 120, 137, 220, 225, 230, 235, 319, 513, 521, 573, 576, 638, 648, 667; oven for treating dates with, against pests, 507; treatment of fruit, etc., with, against Trypetids, 344; treatment of timber and furniture with, against Coleoptera, 110, 181, 203. (See Solar Heat, Steam and Water, Hot.)

Heather, Lepidoptera on,

Scotland, 122.

hebetor, Microbracon.

hederae, Aspidiotus.

Heinsia pulchella, destruction of, against Virachola bimaculata in Sierra Leone, 271.

helichrysi, Anuraphis (see A. padi). Helicobia australis, bionomics and taxonomy of, in Australia, 57.

heliopa, Phthorimaea.

Heliothis armigera, in S. Africa, 223, 323, 330, 487, 488, 489; in Australia, 57, 281, 389; in Fiji, 641; in Peru, 385; in Philippines, 452; in Porto Rico, 328; in N. Rhodesia, 492; in S. Rhodesia, 15, 16, 488, 491, 567; in Russian Union, 146, 147, 304, 587, 589, 609; in Sudan, 488, 573; in U.S.A., 133, 179, 244, 245, 250, 370, 379, 392, 417, 423, 430, 431, 432, 525, 595; on cabbage, 250, 370, 379, 641; on *Citrus*, 15, 16, 330; on cotton, 147, 221, 222, 223, 244, 245, 281, 304, 385, 417, 488, 491, 492, 573, 609; susceptibility of varieties of cotton to, 221; on flax and

ramie, 180, 587; on leguminous plants, 179, 223, 431, 488, 489, 491, 492, 595; comparative attraction of strains of Dolichos lablab for, 488, 491, 492; on maize, 133, 179, 323, 328, 389, 392, 430, 431, 452, 488, 491, 492, 595; on solanaceous plants, 379, 389, 431, 432, 525, 589, 609, 641; concentrations of, on plants in flower, 179, 488, 491; bionomics of, 15, 147, 179, 430, 487, 488, 491, 492, 609; effects of temperature and humidity on, 15, 40, 146, 147; metabolism of, 336; food for adults of, in rearing experiments, 427; new fungus infesting, 392; disease in larvae of, 423; parasites of, 16, 57, 431, 567, 589, 609; ants hindering biological control of, by Microbracon brevicornis, 323; experiments in utilising Trichogramma evanescens against, 304; measures against, 15, 133, 244, 245, 328, 330, 370, 379, 380, 390, 417, 431, 432, 525, 595, 609; other Noctuids confused with,

Heliothis assulta (on tobacco), in Japan, 444; in Sumatra, 17; transmitting virus disease in solanaceous plants, 444; tests of insecticides on, 17.

Heliothis dipsacea, on Taraxacum kok-saghyz in Kazakstan, 45.

Heliothis obsoleta (see H. armigera). Heliothis phloxiphaga, on tomato in California, 431.

Heliothis virescens, on cotton in Peru, 385; on tobacco in Philippines, 451.

Heliothrips fasciatus (see Hercothrips).

Heliothrips haemorrhoidalis, in California, 649; measures against, on Citrus in New Zealand, 56; toxicity of dinitro-o-cyclohexylphenol to, 649.

Helix aspersa, failure to establish Lampyris noctiluca against, in Australia, 546.

Hellebore, tests of, against Sciara on mushrooms, 380.

Hellula undalis, measures against, on vegetables in China, 450; on cabbage in Louisiana and Fiji, 250, 641.

Helopeltis antonii, measures against, on cacao in Java, 523.

Helopeltis bergrothi, forms of, on Eucalyptus in Nyasaland, 126; factors affecting infestation of cotton by, in Sudan, 574; other food-plants of, 574.

Helophorus, on wheat in Denmark, 117.

Helophorus nubilus, 117. hemerobiella, Coleophora.

Hemerobius stigma, predacious on Chermes spp. in Britain, 262, 263; proposed introduction of, into Australia, 263.

Hemerocampa leucostigma, in forests in Quebec, 464.

hemerocampae, Tritneptis. Hemiberlesea (see Aspidiotus).

hemipterus, Carpophilus.

Hemiteles carpocapsae, parasite of
Cydia pomonella in Central Asia.

Cydia pomonella in Central Asia, 588.

Hemiteles chosensis, parasite of

Apanteles liparidis in Japan, 211. Hemiteles kumamotensis, parasite of Apanteles liparidis in Japan, 211. Hemlock (see Tsuga).

Hemlock, Poison (see Conium maculatum).

Hemlock Borer (see Melanophila fulvoguttata).

Hemp, Tetranychus telarius on, in India, 127; insect fauna of, in Italy, 173; pests of, in Ukraine, 229, 322; (for smoking), sulphur unsuitable for use on, 127.

Hemp Flea-beetle (see *Psylliodes* attenuata).

heparana, Tortrix (Pandemis).

heraclella, Aphis. herbarium, Catorama.

Hercothrips fasciatus (Bean Thrips), destruction of wild food-plants of, in California, 240; imported into Japan on oranges from Canada and U.S.A., 616.

herculeanus, Camponotus.

heros, Rhynchites.

Herring Oil, **525.** (See Fish Oil.)

Herse cingulata, measures against,
on sweet potato in Jamaica, **239.**hesperidum, Coccus (Lecanium).

hesperum, Coccus (Lecanium). hesperum, Callidium antennatum. hesperus, Lygus; Reticulitermes.

hesperus, Lygus; Reticulitermes. Hessian Fly (see Mayetiola destructor).

Heterobostrychus aequalis, in timber in Burma, 693.

Heterobostrychus brunneus, in timber in S. Africa, 60.

Heteroderes laurenti, in sugar-cane fields in Florida, 656.

heterogramma, Decadarchis.

Heteroschema rugosopunctata (prima), synonymy of, 434.

Heterospilus anobiidivorus, sp. n., parasite of Neogastrallus librinocens in Florida, 461.

Heterospilus prosopidis (parasite of Bruchus chinensis), bionomics and utilisation of, in Japan, 616.

Heterotermes, in Australia, 545. Heterotermes tenuis, in W. Indies,

55, 56, 239, 464; attacking trees, 464; damaging furniture and timber, 55, 56, 465; characters of, 55; measures against, 55, 465.

Hexachlorethane, tests with, against clothes moths, 635, 636. Hexamera, attempted establishment of, against Oncopera in Australia, 545.

Hexamermis, in Trigonophymus arrogans in Argentina, 214; in Leptinotarsa decemlineata in Germany, 100; in grasshoppers in U.S.A., 76.

Hexamermis acridiorum, 214.

Hexamitodera semivelutina, on clove trees in Netherlands Indies, 227.

hibisci, Eriophyes.

Hibiscus spp., relation of Dysdercus spp. to, in Africa, 222, 489, 490, 492, 573; alternative foodplants of other cotton pests, 136, 164, 184, 572, 573, 577, 661; pests of, in Fiji, 641; Corythuca gossypii on, in Jamaica, 239; Aleurodids on, in greenhouse in Virginia, 530.

Hibiscus esculentus, Tetranychus on, in India, 128; Sylepta derogata

on, in Seychelles, 335.

Hibiscus rosa-sinensis, Saissetia nigra on, in Japan, 445.

hickelianae, Chaetophorus.

Hickory, pests of, in U.S.A., 32, 377.

Hickory Shuckworm (see Enarmonia caryana).

hieracii, Myzus.

Hieracium pilosella, Pentatomids on, in Germany, 632.

hieroglyphica, Saperda scalaris.

hilaris, Bagrada.

hindustanicus, Schizotetranychus (Tetranychus).

hippocastani, Melolontha.

Hippodamia convergens, attempted utilisation of, against Aphids in U.S.A., 30, 428; artificial maintenance of egg-production by, 81; HCN not producing protective stupefaction in, 344.

Hippodamia quinquesignata, on mountains in Utah, 30.

Hippotion celerio, on grape-vines in Queensland, 365.

hirsutus, Phenacoccus.

hirta, Epicometis; Epimadiza; Oemona.

hirtella, Epicometis (see E. hirta). hirticula, Lachnosterna (Phyllophaga).

hirtus, Blissus.

Hispa gestroi, bionomics and control of, on rice in Madagascar, 664.

hispaniolae, Tiphia.

histrionica, Murgantia.

hoefti, Colaphellus.

Holcocera pulverea, predacious on Laccifer lacca in India, 128, 575; longevity of, 128; parasites and biological control of, 129, 575.

Holland, Cydia pomonella on apple in, 578; Contarinia nasturtii on cabbage in, 514; pests and diseases of elm and willow in, 205, 269, 396, 563; Thrips spp. on flax and peas in, 233, 234, 465; Leptinotarsa decembineata in, 62, 106, 269, 583; miscellaneous pests in, 269; Merodon equestris on narcissus in, 494; Aphids and virus diseases of potato in, 157, 563, 578; Tetranychid mites of, 301; pests of wheat in, 268; beneficial insects in, 205, 268, 269, 397, 579; legislation against introduction of L. decembineata into Britain from, 356.

Holly (see Ilex spp.).

Hollyhock, Tischeria omissa on, in California, 569.

Hollyhock Leaf-miner (see *Tischeria* omissa).

Holotrichia (see Lachnosterna).

Homalotylus, new parasites of, in S. Africa, 40.

Homona coffearia (on tea), new parasite of, introduced into Ceylon from Java, 19.

homonae, Macrocentrus.

Homorocoryphus vicinus, in Belgian Congo, 476.

honestus, Dysdercus.

Honey, in baits for ants, 44.

Honey Fungus (see Armillaria mellea).

Hoop Pine (see Araucaria cunning-

Hoplia philanthus, damaging lawns, etc., in Holland, 269.

Hoplocampa, on plum in Czechoslovakia, 468.

brevis, measures Hoplocampa against, on pear and apple in Italy, 197.

Hoplocampa flava (Plum Sawfly), measures against, in Germany and Italy, 46, 97, 297.

Hoplocampa minuta (fulvicornis) (Plum Sawfly), in Denmark, 117; in Germany, 46, 297; in Italy, 97: measures against, 46, 97, 117, 297.

Hoplocampa testudinea (on apple), in Britain, 676; in Denmark, 117: in Poland, 675; bionomics of, 676.

Hoplocerambyx spinicornis, adult longevity of, 613.

Hoplothrips liliorus, sp. n., on lily bulbs in Japan and Korea, 50.

Hops, pests of, in Britain, 1, 333; Hydroecia immanis on, in New York, 593.

Borer (see Hydroecia Hop-vine immanis).

(Holo-Lachnosterna horishana, trichia).

Horismenus floridanus, parasite of Apanteles congregatus in Tennessee, 254.

Horistonotus uhleri, bionomics and control of, in S. Carolina, 500.

Hormius, parasite of Pyralid in parasite Morocco, 193; Leucoptera resembling, in Tanganyika, 127; parasite of, 193.

Horns, damaged by Tinea vastella in Belgian Congo, 294.

Horse Beans (see Beans).

Horse-radish, pests of, in Germany, 110.

horticola, Anomala (Phyllopertha). hortulanus, Bibio.

Household Pests, books on, 237, 382; importance of Psocids as, 102, 648; infesting materials used for insulation, 640; birds' nests of little importance as

source of, 101, House-fly (see Musca domestica). hova, Paratamiclea pallida.

howardi, Dysdercus.

hubbardi, Ataxia.

Hudsona anceps, type of injury to wheat grains by, in New Zealand, 548.

hudsonias, Systena.

huebneri, Papuana.

huegeli, Earias.

humeralis, Carpophilus.

Humidity, relations of mites to, 147, 151, 313, 314, 318, 319, 585, 625, 626: relations of locusts to, 11, 12, 92, 145, 146, 147, 214, 388, 479, 480, 483, 484, 487, 570, 603, 604; relations of other insects to, 15, 16, 17, 26, 27, 44, 47, 60, 61, 89, 102, 112, 119, 120, 130, 143, 145, 146, 147, 148, 167, 195, 196, 223, 224, 228, 229, 230, 234, 235, 239, 262, 265, 266, 316, 231, 232, 234, 200, 441, 442, 454 321, 323, 324, 399, 441, 442, 454, 455, 466, 488, 489, 491, 501, 519, 557, 564, 572, 588, 589, 590, 607, 613, 632, 647, 648, 666, 677, 679, 682, 692; formula for indicating local conditions of, 146; control of, in small cabinets, 568.

humuli, Contarinia (Diplosis):

Phorodon.

Hungary, orchard pests in, 458, 690.

hyacinthi, Rhizoglyphus.

Hyadaphis sii (melliferum, xylostei), transmitting virus disease of celery in California, 590: synonymy of, 590.

hyalinata, Diaphania.

Hyalomyia brasiliensis, sp. n., parasite of Dysdercus spp. in Brazil,

Hyalomyia dysderci, sp. n., parasite of Dysdercus spp. in Brazil, 22. Hyalomyia mendesi, sp. n., para-site of Dysdercus mendesi in Brazil, 22.

Hyalomyodes brasiliensis (see Clistomorpha).

Hyalopterus arundinis, measures against, on plum in Italy, 98.

Hybernia bajaria (see Erannis).

Hyblaea puera, on teak in India and Burma, 612, 693.

Hydrangea hortensis, Tetranychus telarius on, in Germany, 583.

Hydriomena bilineata (see Cidaria). Hydrocarbons, Chlorinated, effect of spraying with, against vine moth pupae, 103. (See Xylamon Preparations.)

Hydrochloric Acid, for removing insecticide residues from fruit, 139, 189, 193, 273, 432, 642, 648, 649; effect of addition of other agents to, 189, 190, 648, 649; sodium silicate compared with,

Hydrocyanic Acid, as a greenhouse fumigant, 115, 356, 526; fumigation of mushroom houses with, 260; fumigation of nursery

stock with, 466, 690; tent fumigation of Citrus with, 15, 55, 149, 292, 342, 343, 440, 508; against pests of stored products, 29, 188, **220, 378, 451**; against *Pyrausta* nubilalis in broom corn, 416; against Lepidoptera in baled cotton and cottonseed, 53-55, 165, 174; against Coleoptera in wood, 181, 203; factors affecting resistance of Coccids to, 342, 343, unsatisfactory against Erythroneura pallidifrons, 115; tests of toxicity of, to insects, **309, 605**; protective stupefaction of insects by, 343, 344, 508; effect of temperature on toxicity of, 111; causing injury to plants, 15, 508, 526; vacuum fumigation with, 53, 54, 174, 181, 417; methods of applying, 15, 174, 260, 440, 508; general paper on uses of, 637. (See Calcium and Potassium Cyanides and Zyklon.)

Hydroecia immanis, measures against, on hops in New York, 593.

Hydroecia micacea, bionomics of, on potato in Scotland, 122.

Hydrogen Phosphide, danger of production of, in poisoned baits, 102.

Hydrogen-ion Concentration, affecting deposits and toxicity of sprays, 411, 412, 649.

Hygrophila, relation of Nomadacris septemfasciata to, in Portuguese E. Africa, 62.

hylaeus, Dolba.

hylas, Platytelenomus.

Hylastes ater, Sirex possibly favoured by, on pines in New Zealand, 504.

Hylastes palliatus, bionomics of, in Sweden, 123.

Hylastes rufipes, on elm in U.S.A., 68, 190, 378, 418; bionomics of, 68, 418; associated with Ophiostoma ulmi, 68, 378; factors affecting infestation of traplogs by, 69, 190.

Hylecoetus (Hylecerus) flabellicornis, on Picea jezoensis in Sakhalin, 19.

Hylecoetus matsushitai, sp. n., on conifers in Japan and Sakhalin,

Hylemyia, scope and synonymy of, 113, 114.

Hylemyia antiqua (Onion Fly), measures against, on onions and leeks in Britain, 674, 675.

Hylemyia brassicae (on crucifers), in Britain, 116; in Germany, 110; new parasite of, in Morocco, 605; bionomics and ecology of, in Russia, 316; measures against, 116, 317.

Hylemyia cilicrura, on sweet lupin

in Russia, 150.

Hylemyia floralis, on horse-radish in Germany, 110; on cabbage in Russia, 316.

Hylemyia genitalis, H. tritici recorded as, in Russian Union, 311.

Hylemyia radicum, new parasite of, on crucifers in Morocco, 605.

Hylemyia seneciella, introduced into New Zealand against Senecio jacobaea, 449.

Hylemyia sepia, bionomics of, on wheat in Morocco, 124.

Hylemyia trichodactyla, on sweet lupin in Russia, **150**.

Hylemyia tritici, sp. n., bionomics of, on cereals in Russian Union, 311; measures against, 312.

Hylesinus vestitus, on pistachio in Greece, 452.

Hyloicus kalmiae (see Sphinx).

Hylotrupes bajulus, in timber in Czechoslovakia, 469; in building timbers in Germany, 6, 110, 202, 203, 204, 268, 335, 520; age of timber affecting infestation by, 520; ecology and nutrition of larvae of, 204, 335, 564; measures against, 6, 110, 203; technique for testing wood preservatives against, 202, 203, 204; criticism of tests of Xylamon against, 520; statistical data on prevalence of, 6, 203, 204, 268.

Hylurgopinus (see Hylastes).

Hylurgops (see Hylastes).

Hymenia recurvalis (fascialis), measures against, in China, **451**; on beet and Amarantus in Fiji, **641**; on spinach in Japan, **441**. hyoscyami, Therapha (Corizus).

Hyparrhenia, relation of Nomadacris septemfasciata to, in Tanganyika, 387, 388.

Hypena abyssinialis (ignotalis), parasites and disease of, on Lantana in India, 612.

Hypera transsylvanica, bionomics and control of, on lucerne in Russia, 108.

Hypera variabilis, tests of insecticides on, in Uzbekistan, 627.

Hyperaspis senegalensis, parasitism affecting value of, against Pseudococcus kenyae in Kenya, 665.

hyperici, Agrilus; Chrysomela

(Chrysolina).

perforatum, Aphis Hypericum chloris on, in Britain, attempted establishment of insects against, in Australia, 282,

Hyperodes griseus, bionomics of, in

New Zealand, **547, 548.**

Hyphantria cunea, in U.S.A., 421, 528; parasites of, 528; basic copper arsenate against, 421.

Hypnoidus riparius, in Britain, 96. hypochlora, Colaspis.

hypogaea, Diarthronomyia.

Hypolixus (see Lixus).

Hyponomeuta padellus, measures against, on plum in Italy, 98.

Hyponomeuta padellus malinellus (on apple), in Manchuria, 340; utilisation of bacteria against, in Russia, 455.

Hypopholis sommeri, parasite of, in

S. Africa. **611**.

Hypopteromalus tabacum, parasite of Apanteles congregatus Florida, 254.

exiguae, parasite of Hyposoter Heliothis armigera in California,

Hypsantyx impressus, parasite of Diprion similis in Poland, 671. Hypsipyla robusta, natural enemies and biological control of, on Cedrela toona and mahogany in India, 612.

Hyptis atrorubens, Larra americana feeding on nectar of, in Brazil,

thymus, parasite of Hyssopus Rhyacionia buoliana in Connecticut, 73, 176; bionomics of, 73.

Hysteroneura setariae (see Carolinaia).

I.

Ibalia leucospoides, bionomics of, in Britain, 504.

icasia, Ecpantheria.

Icerya aegyptiaca, measures against, on Citrus and guava in Egypt, 509.

Icerya purchasi, in Argentina, 23; in Cyprus, 270; in Egypt, 509; occurrence of males of, in Italy and Sicity, 5; in Japan, 445; life-cycle of, in Morocco, 691; in Porto Rico, 434; in Russian Union, 589; on Citrus, 23, 270, 509; on other plants, 5, 434, 691; utilisation of Rodolia cardinalis against, 23, 434, 445, 589; oil sprays against, 509.

Icerva sevchellarum, on avocado in Fiji, 641; Rodolia cardinalis introduced into Mauritius against,

611.

Ichneumon disparis, parasite of Lymantria monacha in Germany, 581.

Ichneumon fabricator, parasite of Erannis defoliaria in Poland, 438. Ichneumon gradarius, parasite of Charaeas graminis in Scotland, 122.

Ichneumon pisorius, parasite of Sphinx pinastri in Poland, 677; parasitised by Theronia atalantae, 677.

Ichneutes spp., parasitic on Pikonema alaskensis in Maine, 540.

Ichneutes fulvipes, 540.

Idaho, ecology and relation to curly-top of Eutettix tenellus in, 286, 412; miscellaneous pests in, 31, 379, 456, 596; larch sawflies in, 599; beneficial insects and biological control in, 30, 31, 379.

Idatiella albisignata, bionomics of, in Australia and New Zealand, 159.

Idechthis gelechiae, parasite Gelechia cercerisella in Kansas, 376.

Idgia melanura, predacious Hypsipyla robusta in India, 612. Idiococcus bambusae, gen. et sp. n., on bamboo in Japan, 624.

idiota, Pentodon; Pyrosis (Bhima).

igniventris, Chrysobothris.

ignotalis, Hypena (see H. abyssinia-

Ilex aquifolium, Phytomyza spp. on, in Britain and N. America, 669.

Ilex opaca, Phytomyza ilicicola on, in N. America, 669.

ilicicola, Phytomyza.

ilicis, Opius; Phytomyza (Chromatomyia).

illepida, Argyroploce (Cryptophlebia).

Illiberis pruni (on fruit trees), bionomics and control of, in China, 450, 668; in Manchuria, 340.

Illinoia (see Macrosiphum).

Illinois, Blissus leucopterus cereals in, 284; miscellaneous pests in, 67, 79, 597; orchard pests in, 70, 255, 526, 527; grasshoppers and their natural enemies in, 67, 75; Syrphids of,

immaculata, Cyclocephala (Ochrosidia); Scutigerella. immanis, Hydroecia (Gortyna). impexus, Otiorrhynchus. impictiventris, Euschistus. implicita, Lachnosterna (Phyllophaga).

importata, Chopardina (see Dolichopoda bormansi). impressicollis, Sphedanolestes.

impressus, Hypsantyx. inaequalis, Coelophora. Inareolata (see Angitia).

incertellus, Schoenobius S.(see

bipunctifer). incisi, Opius. incompleta, Carpomyia. inconsequens, Taeniothrips. inconspicua, Sturmia. inconspicuoides, Sturmia. inconstans, Parastasia.

India, cotton pests in, 127, 128, 228, 575, 576; forest pests in, 224, 612; fruit pests in, 128, 385, 392, 503, 577; insects attacking Lantana in, 612; miscellaneous pests in, 494, 577, 584; sugarcane pests in, 128, 395, 558, 575, 576; pests of timber in, 337, 465, 612; cultivation and pests of lac and its host-trees in, 128, 575; locusts in, 291, 484, 485, 624; 577, 560, mites attacking crops in, 127; Rhizopertha damaging pith helmets in, 612; beneficial insects and biological control in, 128, 129, 228, 263, 385, 505, 506, 575, 577, 612; development and importance of economic entomology in, 391, 448; natural enemies of Pseudococcus introduced into Kenya from, 665. (See also under the various Provinces.)

pomonella Indiana, Cydia apple in, 272, 273, 643, 652; cereal pests in, 378, 414; natural enemies of grasshoppers in, 75; Scolytus multistriatus in, 599.

indica, Malacosoma (Clisiocampa); Margaronia.

indicum, Syntomosphyrum (Melittobia).

Indo-China, Xylotrechus quadripes on coffee in, 49; locusts in, 486. induta, Aeolesthes. inferens, Sesamia. infuscata, Cytherea. infuscatellus, Chilo. infuscatus, Orthostethus.

Inglisia conchiformis, new parasite of, in Cajanus in Uganda, 40. innotata, Scirpophaga.

Inocarpus, pests of, in Fiji, 19, 366. Inocarpus edulis, Dacus psidii on, in Society Is., 294.

inopinata, Cydia (Grapholitha). inopinatus, Prorhinotermes.

inops, Sitona.

inornatus, Chilocorus renipustulatus. Inostemma, parasite of Dasyneura triseti in Austria, 635.

inquisitor, Calosoma.

insectella, auct., Setomorpha S. rutella).

insectella, F., Tinea.

Insecticides, methods of testing, 143, 202, 203, 204, 236, 241, 243, 419, 422, 524, 528, 695; tests of penetration of, through membranes and tracheae of insects, 263, 579, 580; determination of particle-size distribution of, 541; plant derivatives and synthetic organic compounds as potential, 104, 134, 285, 331, 392, 419, 593; patents relating to, 224, 392, 694; certification of, in France, 675; general papers on, 100, 102, 456.

Insects, review of injuries plants caused by toxins of, 624; lists of popular names of, 437; preparation of specimens of, 440: Coleoptera attacking specimens of, 47, 196.

insequalis, Pyrausta. insidiosus, Orius. insignis, Apanteles. instabilis, Macrocentrus. instigator, Pimpla. insulana, Earias. insularis, Frankliniella; Scepticus. integer, Dendrosoter. integriceps, Eurygaster. intermedius, Dysdercus. interpunctella, Plodia. interruptor, Cremastus. interruptus, Euacanthus. interstitialis, Baris. intonsa, Frankliniella.

intricata, Oxya.

intricatoria, Echthromorpha. Invermay Bug (see Nysius turneri). inversa, Lachnosterna (Phyllophaga).

io, Vanessa. indicus, Paratetranychus.

F2

Iowa, Alabama argillacea in, 260; grasshoppers in, 134, 138, 537. Ipobracon, in sugar-cane fields in

Florida, 659.

Ipobracon rimac (parasite of Diatraea saccharalis), introduced into Florida from Peru, 658; introduced into Porto Rico, 502.

Ipocoelius (see Tomicobia).

Ipomoea spp., destruction of, against Cylas formicarius in U.S.A., 379; susceptibility of, to Anophia leucomelas, 444. (See Sweet Potato.)

Ips, new parasite of, in U.S.A., 324. Ips japonicus, on Picea jezoensis in Japan and Sakhalin, 615; possibly a race of I. typographus, 615. Ips sexdentatus, parasite of, in

Turkey, 522.

Ips typographus, bionomics and control of, on spruce in Sweden, 122, 123; I. japonicus possibly a race of, 615.

Iran (see Persia).
iridescens, Levuana.
iridipennis, Xylocopa.

Iris, Aphid transmitting virus disease of, in Germany, 686; Aristotelia on, in Japan, 441; mite on, in New York, 79.

Iron Sulphate (see Ferrous Sulphate).

irrorata, Leucopholis.

isabella, Diacrisia (Isia).

Isaria, infesting Sphinx pinastri in Poland, 677; infesting Petrusina marginata in Porto Rico, 501.

Isaria farinosa, infesting Cydia nigricana in Britain, 39.

Ischnaspis longirostris, establishment of Coccinellids against, on coconut in Seychelles, 335.

ishiharai, Chermes (Adelges). Isia isabella (see Diacrisia).

isocrates, Virachola.

Isoneurothrips parvispinus, measures against, on tobacco in Netherlands Indies, 493.

Isostasius punctiger, parasite of wheat gall-midges in Holland, 268.

Italian Millet (see Setaria italica). Italian Pear Scale (see Epidiaspis leperii).

italicus, Calliptamus (Caloptenus).

Italy, cereal pests in, 3, 174, 197, 198, 199, 295, 472; virus disease of sugar sorghum in, 472; pests of chestnuts in, 358; insect fauna of hemp in, 173; Bruchus

cisti on lupins in, 201, 495; miscellaneous pests in, 5, 471, 473; olive pests in, 98, 473, 568; orchard pests in, 4, 5, 97, 98, 104, 115, 197, 199, 235, 296, 297, 515, 516, 629; pests of stored products in, 172, 198, 199, 234; damaging Dermestes lardarius cocoons in, 198; silkworm Aphids of, 224, 512; Coccids of, 48, 511; Galerucella nymphaeae in, 3; Thaumetopoea spp. in, 516; beneficial insects in, 3, 4, 5, 174, 199, 296, 297, 471, 472, 629, 630; importance of economic entomology in, 474; summaries of plant quarantines in, 274, 592; danger of spread of Leptinotarsa decemlineata into, 583; Encarsia spp. probably imported into Britain from, 265; pests imported into other countries from, 160, 218, 265, 623.

itatiayensis, Ceroplastes. Ivory Nut Palm (see Metroxylon).

J.

Jack Pine (see Pinus banksiana). jacobaeae, Tyria.

jacobsoni, Parahydrotaea.

Jamaica, termites in, 181, 239, 384, 693; miscellaneous pests in, 181, 238; beneficial insects and biological control in, 212, 238, 239, 253; summary of plant quarantines in, 592.

jamaicensis, Lachnosterna.

janata, Achaea.

Japan, cotton pests in, 225, 555, 557; pests of crucifers in, 207, 443, 444, 624; forest pests in, 19, 91, 209, 211, 235, 338, 339, 441, 443, 615; pests of fruit trees in, 207, 209, 211, 225, 338, 361, 440, 441, 556, 557, 616, 617, 620, 668; Chrysomphalus ficus on Elaeagnus pungens in, 91; pests of greenhouse and ornamental plants in, 50, 210, 212, 419, 445, 616, 617; pests of leguminous crops in, 19, 103, 208, 210, 338, 361, 362, 556, 616, 617, 624; pests of mulberry in, 103, 555; rice pests in, 207, 209, 225, 338, 362, 440, 441, 442, 555, 556, 619, 620; pests of other cereals and grasses in, 208, 556; pests of solanaceous plants in, 210, 225, 362, 441, 444, 446, 555;

pests of stored products in, 210, 339, 360, 440, 445; pests of sweet potato in, 444, 445, 446, 620; locust outbreak in, 618; miscellaneous pests in, 17, 210, 440, 441; insects and virus diseases in, 362, 444, 555; illustrations of insects in, 446; new Aphids in, 104, 224, 338; new Coccids in, 103, 624; parasite and hyperparasite of Bombyx mori in, 442, 443; beneficial insects in, 91, 207, 209, 210, 211, 212, 327, 338, 339, 441, 442, 556, 557, 615, 616, 617, 619, 620, 624; biological control in, 445, 619; pests intercepted in, or liable to be introduced into, 50, 208, 568, 615, 616; beneficial insects introduced into U.S.A. from, 144, 377, 615.

Popillia Japanese Beetle (see

japonica).

Japanese Empire, distribution and food-plants of Saissetia nigra in, 445.

Mandated Islands, Tapanese Thysanoptera intercepted Japan from, 50, 616.

Japanese Red Pine (see Pinus densiflora).

Tapanese Weevil (see Calomycterus setarius).

Aserica (Autoserica); japonica, Gobaishia; Popillia.

japonicum, Eriosoma ulmi; Tricho-

gramma.

japonicus, Attagenus; Chermes (Adelges); Coelonotus (Aphidius); Ips; Meteorus; Ptinus; Rhizophagus.

Jarring, to dislodge pests from trees, 10, 156, 196, 300, 544.

Jasminum pubescens, Petrusina marginata on, in Porto Rico, 501. javana, Chaetexorista; Pheidole.

javanica, Tessavatoma.

javanicus, Coptotermes; Goryphus. javanus, Chrysocoris; Plaesius.

Jimson Weed (see Datura stramonium).

jourdani, Aspidiotus (Hemiberlesea). Juglans cinerea, Alsophila pometaria on, in Canada, 663.

Juglans nigra, experiments against Chrysobothris femorata on, in Oklahoma, 420.

Juglans regia (see Walnut). juglansiaria, Zamacra. juglans-vegiae, Aspidiotus.

Jugoslavia, Aphelinus mali in, 10: Ceresa bubalus in, 474; locusts in, 479; orchard pests in, 10, 690; Polychrosis botrana in, 494; summary of plant quarantines, in, 592.

junceus, Tychius.

junci, Lixus.

jungi, Austroicetes (see A. cruciata). Juniper (Juniperus), Phloeosinus spp. on, in Central Asia, France and Algeria, 153, 225, 226; Diaspis visci on, in Czechoslovakia and New York, 469, 529. Juniper Scale (see Diaspis visci). juvencus, Sirex (Paururus).

K.

unsatisfactory Kainit. against wireworms on sugar-cane, 656.

Kakothrips pisivorus (robustus), on leguminous crops in Britain, 332; carried by aerial currents, 167.

Kale, Meligethes aeneus on, in Britain, 333.

kalmiae, Sphinx (Hyloicus).

kalmii, Lygaeus.

kamerunicus, Prosoestus (Derelomus).

kamiyae, Calliceras.

Kansas, Cydia pomonella on apple in, 166; Eutettix tenellus on beet in, 594; insects attacking bindweed in, 375; cereal pests in, 274, 341, 374, 375, 376, 542; miscellaneous pests in, 165, 374, 376; Monotoma parallela in stored cereals in, 600; grass-hoppers in, 341, 376; annual insect population records in, 140, 141; beneficial insects in, 374, 375, 376; mite destroying grasshopper eggs in, 376.

Kaolin, as carrier for sulphur dust,

Kapok, new Lamiid on, in Brazil, 290; Dysdercus superstitiosus on, in Sudan, 573.

karafutonis, Chermes (Adelges).

Karaya Gum, uses of, in sprays, 177, 525, 645.

kazanksyi, Pseudococcus.

Kentucky, Lachnosterna spp. in, 592; Protoparce spp. and their natural enemies in, 252, 253; Scolytus multistriatus in, 599.

Kenya Colony, coffee pests in, 40, 503, 665, 666, 667; miscellaneous pests in, 666, 667;

beneficial insects and biological control in, 40, 503, 665, 666; plant pest legislation in, 274, 666.

kenvae, Pseudococcus.

Kerosene, spraying with, against Leptocoris trivittatus, 374; rice-fields treated with, against Nymphula depunctalis, 364; use of, against Pentalonia nigronervosa, 213; injected into horns against Tinea vastella, 295; in mixtures for treating timber against pests, 181, 337; as a solvent, 426, 450, 510, 544; atomised sprays of insecticides in, against Eutettix tenellus, 412; use of, in olive oil emulsions, 6.

Kerosene Emulsions (see Oil Emulsions).

khapra, Trogoderma (see T. granarium).

kiangsu, Ceracris.

kinugasai, Phloeothrips.

Kites, used in studying distribution of insects by wind, 167.

kivuensis, Anagyrus.

klugi, Aelia; Tritneptis (Ptero-malus).

knighti, Monalonion. kockiana, Carcelia. koebelei, Tritneptis. kojimae, Eutelus. kondonis, Rhizoecus.

konoi, Prociphilus.

Korea, miscellaneous pests in, 19, 50; Diplosis mori on mulberry in, 337; pests of pear in, 557, 668; rice pests in, 557, 618; Galerucella distincta on strawberry in, 619; Plodia interpunctella in stored products in, 225; summary of plant quarantines in, 592; beneficial insects in, 225, 338; utilisation of parasites from, against Popillia and Platyedra in U.S.A., 377, 381.

korschelti, Capitophorus (Cryptomyzus).

kuehniella, Ephestia.

kumamotensis, Hemiteles.

kunoense, Lecanium.

Kutane, 431.

kuvanae, Oöencyrtus.

Kuwanaspis arundinariae, sp. n., on Arundinaria in Formosa, 103. Kuwania betulae, in Russia, 235;

characters of, 235.

Kuwania quercus, in Japan, 235; characters of, 235.

Kuwania rubra, sp. n., bionomics of, on Quercus ilex in France and Corsica, 235.

L.

labdacus, Spathius.

labradus, Zizera (Zizina).

laburni, Aphis (Doralis).

Laccifer lacca (Lac), cultivation and natural enemies of, in India, 128, 129, 575; pests of host-trees of, 128, 129.

Lacerta agilis, observations on, in Russia, 407.

Lachnosterna, in U.S.A., 32, 340, 376, 423, 425; depth of pupation of species of, in Kentucky, 592, 593; in W. Indies, 24, 84, 181, 238; plants attacked by, 24, 238, 376, 425; studies on bionomics and ecology of, 32, 376; natural enemies of, 84, 423; measures against, 32, 340, 376, 425.

Lachnosterna bipartita, in Kentucky, 593.

Lachnosterna clypeata, Scoliid parasites of, on sugar-cane in Florida, 657.

Lachnosterna crassissima, on wheat in Kansas, 376.

Lachnosterna diomphalia, on soy beans in Manchuria, 340.

Lachnosterna fusca, in U.S.A., 32, 593.

Lachnosterna futilis, in Kentucky, 592, 593.

Lachnosterna hirticula, in U.S.A., 32, 592, 593; trees attacked by, 32.

Lachnosterna horishana, bionomics of, on sugar-cane, etc., in Formosa, 363.

Lachnosterna implicita, on willows and poplars in Wisconsin, 32.

Lachnosterna inversa, in Kentucky, 592, 593.

Lachnosterna jamaicensis, on eggplant in Jamaica, 239.

Lachnosterna lanceolata, on wheat in Kansas, 376.

Lachnosterna latifrons, Scoliid parasites of, on sugar-cane in Florida, 657.

Lachnosterna morosa, on soy beans in Manchuria, 340.

Lachnosterna rubiginosa, on Sorghum and wheat in Kansas, 376.

Lachnosterna rugosa, on trees and shrubs in Wisconsin, 32.

Lachnosterna sinensis, bionomics and control of, on tea in Formosa, 447.

Lachnosterna smithi, natural enemies and biological control of, on sugar-cane in Mauritius and W. Indies, 611, 660, 670.

Lachnosterna tristis, on oak

Wisconsin, 32.

Lachnus salignus, new parasite of, on willow in Japan, 615.

Lachnus tropicalis, parasite of, on

chestnut in Japan, 615.

Lacon murinus, feeding habits of larvae of, in Germany, 398; cabbage in Siberia, 679.

Lacon musculus, ecology of, in sugar-cane fields in Formosa, 206.

Lacon setiger, ecology of, in sugarcane fields in Formosa, 206.

Lacon variabilis, on sugar-cane in Queensland, 185.

lacteella, Endrosis.

Lactic Acid, for treating flour from grain attacked by Pentatomids, 353.

Lactuca scariola, destruction of, against Hercothrips fasciatus in California, 240.

lactucae, Amphorophora; (see Macrosiphum ribicola).

Laemophloeus, measures against, in oil-palm kernels in Dahomey, 349. Laemophloeus ferrugineus, import-

ance and control of, in flour mills in Russian Union, 685.

Laemophloeus minutus, in stored grain in U.S.A., 341.

Laemophloeus testaceus, importance and control of, in flour mills in Russian Union, 685.

Laevicephalus collinus, on wheat in Central Asia, 453.

Laevicephalus striatus, on wheat in Central Asia, 453.

laevigatus, Camponotus.

lafauryana, Tortrix (Cacoecia). Lagochirus funestus, utilisation of,

against Opuntia spp. in Queensland, 345.

lalori, Apleurotropis.

lambens, Sarcophaga. Lampides boetica (see Cosmolyce).

Lamprosema blackburni, utilisation of parasite of, on coconut in Hawaii, 367.

Lamtoro (see Leucaena glauca).

Lampyris noctiluca, failure to establish, against Helix aspersa in Australia, 546.

Lachnosterna (Phyllolanceolata, phaga).

langei, Apanteles. langi, Buprestis.

lanigera, Oregma (Ceratovacuna).

lanigerum, Eriosoma (Schizoneura). Lantana, insects attacking, in India, 212, 612; distribution of Teleonemia lantanae, against, in Pacific Islands, **212, 366;** Tribolium destructor imported into Britain from Italy in seed of, 623.

Lantana camara, establishment of Teleonemia lantanae against, in

Australia, 546, 603.

lantanae, Agromyza (Ophiomyia); Teleonemia.

lanuginosum, Eriosoma.

lapathi, Cryptorrhynchus (Sternochetus).

Laphygma exempta, measures against, on sugar-cane in Hawaii, damaging cereals pastures: outbreak of, in Kenya, 666; in Nyasaland, 126; in Queensland, 263, 603; in S. Rhodesia, 124; parasites of, 126, 263, 666; fungi and bacteria infesting, 268, 666.

Laphygma exigua, confused with Heliothis armigera on tomato in California, 431; on ramie in

Transcaucasia, 587.

Laphygma frugiperda, natural enemies of, on maize in Porto Rico, 503; on lucerne and sugarcane in U.S.A., 67, 657; tests of oil baits for, 67.

lapidarius, Biscirus. Lapland, Aphids of, 512.

Larch (Larix), restrictions on importation of, into Britain, 356; pests of, in Czechoslovakia, 469; pests of, in Germany, 7, 216-218, **400, 401;** pests of, in Quebec, 464; Enarmonia diniana on, in Switzerland, 169; pests of, in U.S.A., 461, 599; Acantholyda erythrocephala not observed on, in U.S.A., 651; effect of feeding Lymantria moncha on, dinitro-o-cresol dusts causing injury to, 400.

lardarius, Dermestes.

laricinus, Exenterus. Platycampus; laricis, Marlattia; (Lygaeonematus, Pristiphora

Nematus).

laricivorus, Platycampus. Lariophagus distinguendus (utibilis), parasite of Sitodrepa panicea in

Germany, 8; bionomics and rearing of, parasitising Calandra spp. in Italy and Russia, 199, 304, 551; L. utibilis considered distinct from, 199.

Larix (see Larch).

Larra americana (parasite of Scapteriscus vicinus), introduction of, into Porto Rico from Brazil, 85, 435; bionomics and distribution of, 85.

Larrea, Protargionia larreae on, in Argentina, 23.

larreae, Protargionia.

Lasioderma serricorne (in stored tobacco), in 270; Cyprus, 234; bionomics of, in Italy, in Japan, 210; natural enemies of, in Mauritius, 611; in S. Rhodesia, 125; in Virginia, 380; in imported beans in Germany, 521; in stored cumin seed, 270; Tenebroides mauritanicus associated with, 270; measures against, 125, 235, 380.

Lasiophthicus pyrastri (predacious on Aphids), bionomics and morphology of, in Britain, 628, 629;

in Utah, 30.

Lasioptera, on olive in Greece, 453. Lasiosina cinctipes, parasites of, on barley, in Britain, 562.

Lasius niger, in Germany, 112; resistance to arsenicals of females of, 112.

Lasius niger alienus, associated with Aphid on rubber-producing plants in Russia, 44; measures against, 44.

Lasius niger var. sitkaënsis, in rotten wood in U.S.A., 498.

Laspeyresia (see Cydia).

lata, Amorphoidea.

lataniae, Aspidiotus (Diaspidiotus). lateralis, Eutochia.

laterimaculata, Saperda.

Lathyrus odoratus (see Sweet Pea). latifrons, Lachnosterna (Phyllophaga).

latipes, Xylocopa.

latipetiolator, Phygadeuon.

latiscapus, Cheiloneurus.

latistriga, Parasa.

latiuscula, Cirphis.

latus, Corymbites (Selatosomus).

Latvia, Apion seniculum on Trifolium hybridum in, 47; lists of pests in, 47.

latysiphon, Rhopalosiphoninus.

laurenti, Heteroderes.

Lauryl Rhodanate (see Thiocyanates).

Lawns (see Turf).

Lead, Xylocopa latipes boring in, in India, 337; problem of residues of, on fruit, 139, 168, 188–190, 432, 539, 642.

Lead Arsenate, against beet pests, 283, 639; against cereal pests, 376, 441; against pests of Citrus, 330, 331; coffee painted with, against stem borers and termites, 271: against cotton pests, 137, 243, 246, 271, 414, 417, 661; against pests of forest and shade trees, 28, 53, 74, 177, 279, 386, 420, 461, 529, 532, 540, 645, 663, 664; against orchard pests, 20, 21, 52, 66, 70, 89, 97, 98, 132, 133, 166, 168, 186, 188, 192, 197, 207, 235, 272, 273, 279, 292, 293, 300, 301, 361, 376, 383, 390, 419, 420, 430, 432, 450, 525, 534, 536, 579, 642, 643, 653, 662; against pests of solanaceous plants, 17, 63, 125, 186, 379, 390, 431, 583, 634; against Geometrid on tungoil tree, 614; against vine pests, 164, 516; against pests of other crops, 25, 82, 271, 277, 426, 450; against Lamellicorn larvae in soil, 376, 425, 447, 533; tests of toxicity of, to insects, 17, 137, 243, 246, 277, 420, 421, 528, 627, 634; resistance of strains of Cydia pomonella to, 343; in baits, 186, 379, 390; in baitsprays, 300, 301; dusting with, 17, 25, 82, 186, 187, 246, 277, 383, 390, 417, 421, 431, 634, 639; carriers for, 277, 431; formulae for, in sprays, 17, 28, 52, 53, 66, 89, 97, 125, 132, 137, 164, 166, 177, 186, 192, 235, 271, 279, 283, 293, 330, 376, 383, 390, 414, 420, 430, 432, 450, 461, 529, 534, 536, 540, 663; new method of spraying with, 383; autogiro for applying, 74, 75, 645; adhesives and spreaders for, 28, 52, 53, 70, 74, 166, 177, 189, 207, 208, 235, 279, 330, 383, 430, 525, 540, 642, 645, 661; factors concerned in deposits and retention of sprays of, 52, 272, 273, 412, 539; and Bordeaux mixture, 125, 139, 188, 192, 207, 235, 534, 584, 643, 653; and other copper compounds, 421, 534; and lime, 430, 534; and lime-sulphur, 52, 139, 536; and nicotine, 125, 133, 279, 292, 432, 529, 536; and oils, 28, 139, 166, 188, 190, 192, 293, 376, 426, 430, 432, 461, 529, 536; and pyrethrum, 28, 536, 663; and sulphur, 52, 139, 383, 534, 662; thiuram sulphides compatible with, 532; and injury to plants, 82, 532, 536; effect of, on INDEX, 783

quality of orange fruit, 330, 331; problem of residues of, 139, 168, 188-190, 193, 273, 293, 430, 432, 539, 642; question of danger of, to birds, 159; other insecticides compared with, 17, 66, 97, 98, 166, 177, 197, 246, 273, 419, 421, 441, 532, 536, 540, 593, 627, 634.

Leaf-cutting Ants (see Acromyrmex

and Atta).

Leaf-miners of Moravia, 145.

Leafhopper, Beet (see *Eutettix* tenellus).

Leafhopper, Glasshouse (see Erythroneura pallidifrons).

Leafhopper, Potato (see *Empoasca fabae*).

Leafhopper, Six-spotted (see *Macrosteles divisus*).

Leather Jackets (see *Tipula* spp.). Lebeda nobilis, on oaks in China, 360.

lecanii, Coccophagus (see C. lycimnia); Muscideopsis.

Lecanium, on plum in Czechoslovakia, 468.

Lecanium bulgariense, sp. n., bionomics and control of, on roses in Bulgaria, 402.

Lecanium corni, measures against, on vines in Germany, 103; reduced importance of, on plum in Central Europe, 690; on Robinia and fruit trees in Uzbekistan, 474; physiological factors influencing mass increase of, 112, 113, 507.

Lecanium coryli var. turanicum, on fruit trees in Uzbekistan, 474.

Lecanium hesperidum (see Coccus). Lecanium kunoense, susceptibility of varieties of Ulmus propinqua to, in Japan, 443.

Lecanium longulum (see Coccus).

Lecanium persicae, in Australia, 294, 365; on vines, 365; new parasite of, 294.

Lecanium pruinosum, measures against, on walnut in California, 649.

Lecanium prunastri, on fruit trees in Uzbekistan, 474.

Lecanium rugulosum, on fruit trees in Uzbekistan, 474.

Lecanium unifasciatum, on peach and almond in Uzbekistan, 474.

lecontei, Agonoderus; Diprion (Neodiprion).

Leefmansia bicolor, liberation of, against Sexava in New Guinea Territory, 277.

Leek, Hylemyia antiqua on, in Britain, 675.

lefroyi, Microbracon.

legitima, Polia.

leguminosae, Aphis (see A. laburni).

Lema bilineata, on tobacco in S. Africa, **360**.

Lema oryzae, measures against, on rice in Japan, 440; in Manchuria, 361.

Lemon, pests of, in California, 55, 343, 377, 425, 432, 433, 460; Geometrid on, in Sicily, 630; tartar emetic sprays causing slight injury to, 425; (fruit), in baits for cutworms, 387, 390.

Lemon Chrome (see Barium Chromate).

leperii, Epidiaspis.

Lepidiota albohirtum (see Lepido-derma).

Lepidiota caudata, damaging pastures in Queensland, 603.

Lepidiota frenchi, on sugar-cane in Queensland, 88, 186; measures against, 88.

Lepidium alyssoides, relation of Euttetix tenellus to, in U.S.A., 593, 594.

Lepidoderma albohirtum, in Queensland, 88, 185, 294, 603; on sugar-cane, 88, 185; on maize, 603; on forest trees, 185; parasite of, 294; establishment of Bufo marinus against, 185; other measures against, 88.

Lepidoglyphus destructor, in stored grain in Russian Union, 312, 585, 685; not surviving under field conditions, 312; factors affecting production of hypopi in, 585; measures against, 685.

Lepidopterous Larvae, physiology of senses of, 7.

Lepidosaphes, measures against, on Citrus in Egypt, 509; revision of Italian species of, 511.

Lepidosaphes beckii (on Citrus), in Argentina, 23; in Cyprus, 270; in Egypt, 509; bionomics of, in Transcaucasia, 148, 149; in U.S.A., 256, 257, 460; effect of spray residues on populations of, 257; measures against, 23, 460, 509.

Lepidosaphes gloveri, on Citrus in Transcaucasia, 148, 150; bionomics of, 148, 149; measures against, 150.

Lepidosaphes tubulorum, susceptibility of varieties of Ulmus propingua to, in Japan, 443.

Lepidosaphes ulmi (on fruit trees), bionomics of, in W. Australia, 130; on elms in Japan, 443; in Manchuria, 340; in U.S.A., 432, 534; measures against, 130, 432, 534.

Lepidosaphes zlocistii, on fruit trees in Uzbekistan, 474.

Lepisma, infesting house insulation in U.S.A., 640.

Lepisma saccharina, treatment of books against, in Fiji, 366.

Leptacis tipulae, parasite of wheat gall-midges in Holland, 269.

Leptinotarsa decemlineata (Potato Beetle), in Belgium, 106, 583; legislation against introduction of, into Britain, 356; in Br. Columbia, 67; in France, 106, 107, 169, 194, 349, 350, 357, 358, 583, 584, 622, 634, 675; in Germany, 48, 100, 106, 158, 160, 354, 512, 583; in Holland, 62, 106, 269, 583; in Luxemburg, 106, 583; in Switzerland, 106, 350, 583; in U.S.A., 70, 134, 421; danger of spread of, into Italy and Spain, 583; bionomics of, 106, 349, 584, 622; natural enemies of, 100, 107, 194, 354; susceptibility of varieties of potato to, 158, 193; effect of species and hybrids of Solanum on, 107, 158, 169, 357, 358; experiment with, on tomato, 113; definition of focus of infestation by, 107; larval instars of, 584; measures against, 62, 67, 70, 106, 107, 158, 159, 349, 421, 583, 634; international conference on control of, 106; susceptibility of, to hydrocyanic acid gas, 605; magnesium sulphate ineffective against, 134.

Leptispa pygmaea, measures against, on rice in Mysore, 291.

Leptocoris trivittatus, in Canada, 374; bionomics and control of, in Kansas, 374.

Leptocorisa, Catascopus facialis attacking, 642.

Leptocryptus pellucidator, hyperparasite of Diprion similis in Poland, 671.

Leptoglossus, vector of Nematospora, 389.

Leptoglossus australis, on pumpkin in Fiji, 641.

Leptomastidea, African species of, 503.

Leptomastix, African species of, 40, 503.

Leptomastix bifasciatus, sp. n., parasite of Pseudococcus in Tanganyika and S. Africa, 40.

Leptomastix dactylopii, causes of immunity of certain mealybugs from, 598.

Leptomastix nigrum, sp. n., parasite of Pseudococcus in Transvaal, 40.

Leptomastix phenacocci, sp. n. (parasite of *Phenacoccus hirsutus*), introduced into Egypt from Java, **506.**

Leptothrips mali, predacious on insects in Louisiana, 372.

Leptozele (see Macrocentrus).

Leskia aurea, parasite of Aegeria typhiaeformis in Italy, 4, 629, 630; in France and Sweden, 629; doubtfully parasitic on Cydia pomonella, 4; bionomics of, 629, 630.

Lespedeza, Horistonotus uhleri on,

in S. Carolina, 500.

Lestodiplosis pini, predacious on Chermes spp. in Britain, 262, 263; proposed introduction of, into Australia, 263; predacious on Leucopis obscura, 263.

Lethane, 463. (See Thiocyanates.) Lethane Spreader, for derris and

cubé sprays, 30.

Lettuce, restrictions on importation of, into Australia against Pieris rapae, 344; Aphids on, in Britain, 671, 672; Phytomyza atricornis on, in Japan, 444; Macrosteles divisus on, in New Jersey, 536; Forficula auricularia on, in Ontario, 601; experiments with insects and, 398, 423, 459, 519; insects and virus diseases of, 536, 546, 671.

Leucaena glauca, pruning of, against Pseudococcus citri in Java, 161.

Leucodesmia, parasite of Gelechia cercerisella in Kansas, 376.

Leucodiaspis archangelskyae, on fruit trees in Uzbekistan, 474.

leucoloma, Naupactus. leucomelas, Anophia.

leuconotus, Anthores. leucophaearia, Erannis.

Leucopholis irrorata (Sugar-cane Beetle), intercepted in Hawaii in aeroplane from Philippines, 368. leucophthalmus, Philaenus.

Leucopis griseola, bionomics of, attacking Aphids in India, 228.

Leucopis obscura, predacious on Chermes spp. in Britain, 262, 263; proposed introduction of, into Australia, 263; natural enemies of, 263.

Leucoptera, measures against, on coffee in Tanganyika, 126; para-

sites of, 127.

Leucoptera coffeella, (on coffee), ecology of, in Tanganyika, 127;
Mirax introduced into Porto Rico from Guadeloupe against, 435; in St. Lucia, 435.

Leucoptera daricella, ecology of, on coffee in Tanganyika, 127.

Leucoptera scitella, on apple in Uzbekistan, 623.

leucopterus, Blissus. leucospoides, Ibalia.

leucostigma, Hemerocampa.

leucostoma, Cydia (Laspeyresia).

Leucotermes tenuis (see Heterotermes).

leucotreta, Argyroploce.

leucurus, Aesiotes.

leveri, Phaenocarpa.

levis, Trionymus.

Levuana iridescens, parasite of, on coconut in Fiji, 212.

libatrix, Scoliopteryx.

Libocedrus decurrens, Scolytids on, in U.S.A., 324.

librinocens, Neogastrallus.

Libya, locusts in, 478, 481. Licuri Palm (see Cocos coronata).

Licuri Palm (see Cocos coronata).
Licurioba Palm (see Cocos vagans).
Light, influence of: on forms of
Aphis chloris, 59; on feeding of
Lamellicorn beetles, 338; on
locusts, 603, 604; on derris
deposits, 69, 241.

Light (Artificial), insects attracted to, 22, 209, 364, 463, 465; reactions of insects to types of,

31, 441, 442.

Light-traps, for Lepidoptera, 25, 26, 31, 32, 172, 348, 441, 592; for other insects, 50, 95, 164, 340, 501; electrocuting types of, 31.

lignosellus, Elasmopalpus.

Ligyrus gibbosus, on garden plants in Kansas, 376.

Ligyrus tumulosus, on egg-plant in Jamaica, 239.

Lilac (Syringa), Cantharis obscura on, in France, 623; Aphid migrating to Abies from, in Japan, 211.

lilacinus, Pseudococcus.

Lilies (Lilium), Lilioceris lilii on, in France, 268; pests of, in

Germany, 267, 686; pests of, in Japan and Korea, 50, 419, 617; Rhizoglyphus hyacinthi on, in U.S.A., 419; Aphis frangulae transmitting virus disease of, 686; heat treatment causing injury to bulbs of, 617.

Lilioceris lilii, bionomics of, on lily in Germany and France, 267,

268

liliorus, Hoplothrips (Trichothrips). Lima Bean Pod Borer (see Etiella

zinckenella).

Lima Beans, Diacrisia obliqua on, in China, 129; pests of, in Porto Rico, 435, 501; pests of, in U.S.A., 381, 595; susceptibility of varieties of, to pests, 501, 595; problem of fluorine residues on, 595.

limacina, Caliroa (Eriocampoides). Limax agrestis, possible host of Hexamermis in Germany, 100.

limbata, Brontispa.

limbatum, Agonum (see A. pallipes). Lime (Citrus), pests of, in Grenada, 25.

Lime (*Tilia*), Ceresa bubalus on, in France, **689**; Caliroa annulipes on, in Germany, **204**.

Lime, use of, against Aphomia gularis, 521; dusting with, against cacao thrips, 24; as a carrier for dusts, 71, 177, 187, 213, 246, 247, 258, 261, 277, 283, 342, 370, 371, 372, 380, 418, 426, 461, 497, 661; as a repellent, 24, 299, 340, 420, 532, 533, 665; addition of, to arsenical sprays, 158, 172, 189, 283, 421, 430, 438, 451, 534, 634, 664, 683; addition of, to other sprays (see also Bordeaux Mixture), 158, 256, 257, 532, 634; relation of, to spray residues on apples, 189, 190; formulae containing, 71, 172, 177, 187, 189, 213, 246, 247, 256, 257, 283, 342, 370, 372, 380, 418, 425, 426, 430, 438, 461, 497, 532, 534, 644, 683.

Lime-sulphur, unsatisfactory against Bemisia gossypiperda, 560; against Chermes abietis, 529; against Coccids, 23, 34, 128, 130, 432, 433, 460, 529, 530, 690; effect of residues of, on Coccid populations, 257; apparently repellent to Conotrachellus nenuphar, 383; against mites, 2, 57, 164, 187, 256, 365, 409, 429, 433, 460; effects of, against Paratrioza cockerelli on tomato, 72, 73, 79, 651; against

thrips, 14, 433, 460, 666; formulae for, 2, 23, 72, 79, 130, 164, 187, 257, 278, 343, 365, 409, 432, 460, 498, 529, 530, 664, 666, 690; formulae containing, 52, 57, 432, 460, 498, 664; and aluminium sulphate with lime, 257; and arsenicals, 52, 139, 421, 664; effect of, on lead-arsenate spray residues, 139; and ferrous sulphate, 460; and nicotine sulphate, 292, 429, 498, 536; and oil emulsions, 57, 409, 430, 432, 433, 460, 536; and pyrethrum, 536; and sulphur, 57, 433, 460, 534; thiocyanate compatible with, 259; factors affecting injury to Citrus by, 278, 460.

liminaris, Phloeotribus (Phthorophloeus).

Limonius (see Pheletes).

Limothrips cerealium (on cereals), in Holland, 269; bionomics of, in Sweden, 121.

Limothrips denticornis, bionomics of, on cereals in Sweden, 121.

Limothrips schmutzi, doubtful vector of virus disease of sugar sorghum in Italy, 472.

linarius, Thrips (see T. lini). linearis, Atomaria; Lyctus; Oberea. lineata, Celerio; Sitona.

lineatella, Anarsia.

lineaticollis, Antestia.

lineatus, Agriotes.

lineolata, Diatraea.

lineolatus, Adelphocoris.

lini, Thrips.

Linoclostis gonatias, on tea, etc., in Formosa, 447.

Linseed (see Flax).

Linseed Oil (see Oil, Linseed).

Linum usitatissimum (see Flax).

Liocola brevitarsis, in manure in sugar-cane fields in Formosa, 446. Liothrips urichi, attempted intro-

duction of, into Solomon and Shortland Is. against *Clidemia*, **366.**

Liothrips vaneeckei, on lily bulbs in Japan and Korea, 50.

liparidis, Apanteles.

Liparis chrysorrhoea, auct. (see Nygmia phaeorrhoea).

Liparis monacha (see Lymantria). Liposcelis divinatorius, in houses and ground feed in U.S.A., 648, 650;

measures against, 648.

Lissonata cylindrator, parasite of
Trachea secalis in Scotland, 122.

Lissonata sulbhurifera parasite of

Lissonota sulphurifera, parasite of Trachea secalis in Scotland, 122.

Lissopimpla semipunctata, parasite of cutworms in New South Wales, 279.

Listroderes obliquus (on vegetables and tobacco), in Queensland, 186; in U.S.A., 249, 327, 372, 599; bionomics and control of, 186, 372; tests of insecticides on, 249.

Lita solanella (see Phthorimaea operculella).

Litchee, Lobesia aeolopa on, in Formosa, 340.

Lithocolletis, weight of foliage in relation to, 143.

Lithocolletis platani, parasites of, on Platanus spp. in Turkey, 219.

Lithosia deplana (see Eilema depressa).

litura, Prodenia.

livida, Cantharis.

lividicorpus, Eurydinota.

lividigaster, Scymnodes.

Lixophaga diatraeae (parasite of Diatraea saccharalis), further attempts to establish, in U.S.A., 658; distribution and utilisation of, in W. Indies, 182, 183, 502, 658, 659, 660; not parasitic on D. canella, 182; ecology of, 659.

Lixus, new parasite of, in India, 506: Lixus junci, bionomics and control of, on beet in Morocco, 283.

Lixus truncatulus, new parasite of, in India, 506.

Lizards, destroying noxious insects, 48, 394, 407; damaging trapditches in beet fields, 407.

Lloyd's Reagent, nicotine sulphate adsorbed by, 133.

Lobesia aeolopa, on tea and litchee in Formosa, 339, 340.

Loblolly Pine (see Pinus taeda).

Locust, Bamboo (see Ceracris kiang-su).

Locust, Lubberly (see Romalea mocroptera).

Locust, Moroccan (see *Dociostaurus* maroccanus).

Locust, Red (see Nomadacris septem-fasciata).

Locust, Rocky Mountain (see Melanoplus mexicanus).

Locusta migratoria, in India, 291, 484, 559, 560; in Poland, 9, 403; breeding areas and ecology of, in Russia, 109, 146, 553; plants attacked by, 291, 559, 560; phases and migrations of, 109, 284, 559, 560; effects of temperature, humidity and light on, 146, 553, 554, 603, 604; tests of insecticides on, 309, 310.

Locusta migratoria capito, in Madagascar, 14, 62, 478; bionomics of, 14, 62; measures against, 62.

Locusta migratoria manilensis (migratorioides, auct.), in China, 450, 451, 477, 486; in Netherlands Indies, 478, 486; in Philippines, 451, 478, 485, 486; probable range of spread of, from Philippines, 486; plants attacked by, 451; factors affecting outbreaks of, 485; measures against, 450, 477, 478.

Locusta migratoria migratorioides, 480, 483; in Italian E. Africa, 478, 570; in Portuguese E. Africa, 479; in Fr. W. Africa, 11-14, 280, 477, 478, 480, 482, 570, 571; in Belgian Congo, 476; in Sierra Leone, 271; in Sudan, 479, 481; in Uganda, 482; outbreak centres of, 11-14, 480, 482; growth and ecdysis in, 264, 456; other bionomics of, 11-14, 264; experiments with Beauveria bassiana and, 92; measures against, 477, 478.

Locustana pardalina, in Portuguese E. Africa, 479; bionomics of, in S. and S.W. Africa, 92, 93, 393, 394; measures against, 92, 93.

locustarum, Eutrombidium.

Locusts and Grasshoppers, in E. 486, 487, 571; in Arabia, 480, 481, 482; in Argentina, 213, 214, 476, 543; in Australia, 56, 481; in Greece, 478; in India, 291, 484, 485, 559, 560, 577, 624; in Netherlands Indies, 478, 486; in Japan, 618; in Jugoslavia, 479; in Korea, 557; in Libya, 478, 481; in Madagascar, 14, 62, 478, 543, 571; in Paraguay, 543; in Persia, 482; in Philippines, 451, 478, 485, 486; probable range of spread of, from Philippines, 486; in Poland, 9, **403**; in Portugal, **479**; in N. Rhodesia, **483**, **484**; in S. Rhodesia, 124, 125, 637; in Russian Union, 109, 145, 146,

153, 160, 320, 321, 386, 553, 627; key to, of W. Siberia, 694; in Spain, 477; in Anglo-Egyptian Sudan, 479, 480, 481; in U.S.A., 32, 33, 48, 67, 75, 76, 134, 138, 139, 178, 240, 341, 376, 427, 485, 537, 569, 598, 646, 654, 655; international conference on, 475-486; phases of, 11, 12, 13, 48, 93, 321, 387, 388, 393, 394, 480-487, 522, 545, 560, 569, 570, 571; colour variations and anatomy of, 48, 387, 482, 483, 486, 512, 545, 570; breeding centres and ecology of, 11–14, 62, 93, 109, 321, 386, 387, 388, 393, 394, 477, 479-487, 545, 553, 570, 571; migrations of, 33, 109, 126, 291, 386, 387, 388, 394, 478, 479, 481, 484, 485, 487, 559, 560, 646, 666; maps illustrating invasion areas of, 476, 477, 478, 481, 482, 483; effects of temperature, humidity and rainfall on, 11-14, 41, 93, 145, 146, 291, 345, 393, 394, 477, 479, 480, 481, 483, 484, 485, 487, 545, 553, 569, 603; correlation of outbreaks of, with sunspots, 485; reaction of, to light, 603; growth and ecdysis of, 264; effects of deficient diets on, 604; respiration of, 104; birds destroying, 387; insect enemies of, 56, 75, 214, 387, 394, 477, 545; Helicobia australis not a parasite of, 57; mites infesting, 75, 376; fungi infesting, 75, 91, 92, 145, 427; Mermithid parasites of, 75, 76, 214; other organisms in, 486; surveys and estimation of populations of, 76, 153, 160, 320, 387, 393, 569; measures against (see also under Baits), 9, 62, 92, 93, 214, 450, 476, 477, 478, 479, 557, 618, 624, 655; financial loss due to and expenditure on control of, 130, 292, 476, 477, 478, 480, 481; tests of insecticides on, 309, 310, 627; reviews of problems of, 336, 572.

Lolium, Hyperodes griseus on, in New Zealand, **547**; Pediculopsis graminum on, in Russian Union,

Lomaspora (see Arabis).

Lonchaea aristella, intercepted in figs in Russian Union, 440.

Lonchocarpus, cultivation of, for insecticides in Belgian Congo, 294; survey of data on, as an insecticide, 104. (See Cubé.) longana, Cnephasia.

longicaudata, Ctenolepisma. longicaudatus, Biosteres (Opius). longiclavis, Pagiophloeus.

longicorne, Stromatium.

longicornis, Paratrechina (Prenolepis); Scolothrips.

Longicorns, catalogue and data on biology of, in India, Burma and Ceylon, **612**.

longipennis, Dissosteira; Eirenephilus.

longirostris, Ischnaspis; Rhynchocoris.

longispina, Bemisia.

longispinus, Pseudococcus (see P. adonidum).

Longitarsus, on ramie in Transcaucasia, 587.

longitarsus, Amblynotus; Cotinis. longulus, Coccus (Lecanium).

Lonicera, Rhagoletis cerasi not attracted by extracts of, 297.

Lonicera morrowii, Aphids on, in Japan, 104, 211.

Loochoo Islands, pests of sweet potato in, 444, 616.

lopezi, Eudeleboea.

Lophosia fasciata, parasite of Aelia acuminata in Germany, 64; not found in E. Prussia, 632.

Lophyrus (see Diprion).

Loquat, Lepidoptera on, in Formosa and Japan, 206, 207; breeding experiments with Dacus ferrugineus tryoni on, in Queensland, 345.

loreyi, Cirphis. loricata, Ulobaris.

Lorol Alcohol, spray formula con-

taining, 532.

Louisiana, cotton pests in, 246, 341, 372, 538; pests of crucifers in, 250, 254, 370, 371, 372, 373; pests of pecan in, 255, 372; Anticarsia gemmatalis on soy beans in, 258, 371, 373, 422; sugar-cane pests in, 143, 248, 371, 372, 378, 657, 658; Cylas formicarius on sweet potato in, 373, 379, 539, 644; miscellaneous pests in, 248, 370, 537, 599; Listroderes obliquus in, 250, 372; legislation against Naupactus spp. in, 275; Thysanoptera of, 372; beneficial insects and biological control in, 143, 372, 373, 378, 538, 656, 658, 659.

lounsburyi, Aspidiotiphagus.

Loxostege sticticalis, in Kansas, 376; migrations of, in Saskatchewan, 27; in Russian Union, 44, 109, 146, 306, 307, 313, 455, 680;

food-plants of, 44, 307, 313, 376, 680; effects of temperature and humidity on, 146, 455; race of *Trichogramma evanescens* parasitising, 109; experiments with fungi against, 306, 307; beet protecting *Scorzonera* against, 680; other measures against, 313, 376.

Lubberly Locust (see Romalea

microptera).

lubricipeda, Diacrisia (Spilosoma). Lucerne, grasshopper damaging, in Argentina, 214; Contarinia medicaginis on, in Germany, 457; Smynthurus viridis on, in New Zealand, 550; pests of, in Russian Union, 108, 215, 216, 305, 474, 475, 552; bubalus on, in Switzerland, 169; pests of, in U.S.A., 30, 31, 32, 67, 323, 370, 378, 381, 423, 427, 462, 599; cutworms on, in Victoria, 390; relation of Lygus spp. to seed production of, 462; time of sowing and harvesting of, in relation to weevils, 108, 109, 371. Lucerne Flea (see Smynthurus viri-

ais).
Lucerne Flower Gall-midge (see

Contarinia medicaginis). Lucerne Leaf Bug (see Adelpho-

Lucerne Leaf Bug (see Adelphocoris lineolatus).

Lucilia, suitability of, as host for Trichogramma evanescens, 36.

lucorum, Lygus; Zenillia (Carcelia). luctuosus, Oxycarenus.

Ludius aeripennis (see Corymbites). luminosus, Pyrophorus.

lunata, Membracis (see M. foliata). luniger, Syrphus.

Luperina testacea, on wheat in Denmark, 117.

Luperodes quadriguttatus, on soy beans in Manchuria, 340.

Lupins (Lupinus spp.), Cantharis obscura on, in France, 623; Aphids and virus disease of, in Germany, 686, 687, 688; Etiella zinckenella on, in Iberian Peninsula, 495; Bruchus cisti on, in Italy and Sicily, 201, 495; pests of, in Russia, 150, 151, 683; susceptibility to pests of species and varieties of, 150, 151, 495.

luridula, Anomis. luridus, Troilus. lutaria, Sialis. luteola, Galerucella. luteovaria, Urochela. lutescens, Phonoctonus. luteum, Trichogramma.

Luxemburg, Leptinotarsa decemlineata on potato in, 106, 583; legislation against introduction of L. decemlineata into Britain from, 356.

lybica, Empoasca.

lycimnia, Coccophagus.

Lycium spp., Paratrioza spp. on, in U.S.A., 422, 423, 651.

lycopersicella, Phthorimaea (Gnori-

moschema).

lycopersici, Eriophyes; Phyllocoptes. Lycophotia praecox, on soy beans in Manchuria, 340.

Lycophotia saucia (see Peridroma). Lyctus (in timber), in Burma, 693; in U.S.A., 499.

Lyctus brunneus (in timber), in S. Africa, 60; in Australia, 509; treatment of timber against, 509. linearis (canaliculatus),

damaging floors in Czechoslo-

vakia, 469.

Lyctus tomentosus, measures against, in furniture in Jamaica, **181.**

Lydella nigripes, parasite of Lepidoptera in Poland, 438.

Lygaeonematus (see Pristiphora). Lygaeonematus pini (see Pristiphora abietina).

Lygaeonematus stecki (see Pristi-

phora pallida).

Lygaeus equestris, on Taraxacum kok-saghyz in Kazakstan, 45. Lygaeus kalmii, measures against,

in California, 649.

Lygidea mendax (on apple), in New York, 533; measures against, in Nova Scotia, 662.

Lygocerus testaceimanus, parasite of Leucopis obscura in Britain, 263. Lygus, measures against, on cotton

and lucerne in U.S.A., 246, 417, 463; effect of, on cotton, 417.

Lygus coffeae, on coffee in Tanganyika, 127.

Lygus communis, measures against, on apple in Nova Scotia, 662.

Lygus elisus, in U.S.A., 417, 462; on cotton, 417; bionomics of, on lucerne, 462.

Lygus hesperus, in U.S.A., 417, 462; on cotton, 417; bionomics of, on lucerne, 462.

Lygus lucorum, possibly transmitting virus disease in cotton and egg-plant in Japan, 225, 555.

Lygus pratensis, in Czechoslovakia, 469; in Italy, 472; in Poland, 162, 163; in Russian Union, 45, 303; in U.S.A., 84, 246, 533, 590; bionomics of, on apple, 533; doubtful relation of, to virus diseases of plants, 84, 163, 303, 472, 590; miscellaneous foodplants of, 45, 162, 246, 469.

Lygus pratensis oblineatus, on cotton

in Arizona, 417.

Lygus spinolae, measures against,

on vines in Austria, 45.

Lymantria dispar, in Czechoslovakia, 463; in Germany, 354; in Japan, 209, 327, 442, 615; in Poland, 163; in Russian Union, 313, 314, 404, 405, 406, 456, 627; in U.S.A., 138, 139, 170, 180, 645; on fruit trees, 163, 468; studies on nutrition of, 143, 404, 456; effects of temperature on, 313, 314, 404, 405, 406; application of bioclimatics to study of, 327; natural enemies of, 209, 302, 327, **354**, **442**, **615**; utilisation of Calosoma sycophanta against, 170; other measures against, 138, 139, 645; tests of insecticides on, 310, 627.

Lymantria monacha (in forests), 223; in Czechoslovakia, 469; in Germany, 99, 170, 218, 223, 231, 298, 335, 398, 400, 401, 467, 512, 563, 581; in Poland, 398; physiology of, 99, 298, 398, 399, 564; natural enemies of, 170, 218, 335, 581; proposed utilisation of Calosoma sycophanta against, 170; Barbitistes constrictus not predacious on, 401; infected with polyhedral disease, 582; use of scent of females for estimating prevalence of, 400; estimation of age and abundance of, from larval excreta, 563; insecticides against, 231, 400, 512; characters of, 466.

Lyncestis acontioides, on Poinciana

in Jamaica, 181.

Μ.

machaeralis, Apanteles; Hapalia. Mackiella phoenicis, gen. et sp. n., on date palm in California, 288.

Macrocentrus, scope and synonymy of, **263**.

Macrocentrus abdominalis, parasite of Tortricids in Transcaucasia, 587.

Macrocentrus amicroploides, parasite of Alsophila pometaria in Canada, 663.

Macrocentrus ancylivorus (parasite of Cydia molesta), introduced into Australia, 544,

attempted utilisation of, against *C. nigricana* in Canada, **39**; other hosts and utilisation of, in U.S.A., **376**, **377**; parasitised by *Dibrachys cavus*, **640**.

Macrocentrus calacte, sp. n., parasite of Argyroploce illepida in

Fiji, 19, 213.

Macrocentrus delicatus, parasite of Cydia molesta in U.S.A., 377.

Macrocentrus homonae, sp. n., introduced into Ceylon from Java against Homona coffearia, 19; bionomics of, 20.

Macrocentrus instabilis, parasite of Cydia molesta in U.S.A., 377.

Macrocentrus thoracicus (parasite of Cydia molesta), attempted utilisation of, against C. nigricana in Canada, 39.

Macrocentrus trimaculatus, systema-

tic position of, 263.

Macroglenes penetrans, parasite of wheat gall-midges in Holland, 268.

Macropsis munda (see Bythoscopus).

macroscopa, Brachmia.

Macrosiphoniella (see Macrosiphum).

Macrosiphum aucubae (see M.

solani).

Macrosiphum avenae, on sugar-cane in Hawaii, 367; Leucopis griseola predacious on, in India, 228; possible vector of disease of sugar sorghum in Italy, 472; distribution and food-plants of, 472.

Macrosiphum dirhodum, possible vector of disease of sugar sorghum in Italy, 472; distribution and food-plants of, 472.

Macrosiphum gei, auct. (see M.

solanifolii).

Macrosiphum gossypii turanicum, on cotton in Transcaucasia, **552.** Macrosiphum granarium (see M. avenae).

Macrosiphum maculata, sp. n., on lucerne in Central Asia, **552.**

Macrosiphum moltshanovi, on peas and vetch in Central Asia, 552.

Macrosiphum onobrychis (on peas and lucerne), in Russia, 305; in U.S.A., 29-31, 33, 323, 370, 415, 497, 531, 645; susceptibility of varieties of peas to, 33; experiments with pea virus in beans and, 370; physiological conditions producing wing development in, 323; natural enemies and biological control of, 30, 31, 305; other measures against, 29, 415, 645; importance of timing measures against, 415, 531.

Macrosiphum pelargonii, and virus diseases of potato in Russia, 303. Macrosiphum pisi (see M. onobrychis).

Macrosiphum pseudosolani (see M.

solani)

Macrosiphum ribicola, food-plants of, in Britain, 333, 671, 672; transmitting lettuce mosaic, 671.

Macrosiphum rosae, and mosaic disease of narcissus in New York, 652; on rose in Tasmania, 21.

Macrosiphum rubiellum, on black-

berry in Britain, 333.

Macrosiphum solani, in Britain, 563, 671; in Germany, 157, 267, 634, 686, 688; in Holland, 578; in U.S.A., 83, 369, 590, 652; and virus diseases of potato, 83, 84, 157, 267, 563, 578, 634, 687; and virus diseases of other plants, 84, 369, 590, 652, 671, 672, 686, 688; flight habits of, 634; synonymy of, 590.

Macrosiphum solanifolii, measures against, on Passiflora edulis in Australia, 448; in Britain, 563, 671, 672; in Germany, 157, 267, 634, 686; in Holland, 563, 578; in U.S.A., 83, 369, 370, 563, 652; bionomics of, 578, 634; overwintering on rose, 563, 686; and virus diseases of potato, 83, 84, 157, 267, 563, 578, 634, 686; and virus diseases of other plants, 84, 369, 370, 448, 652, 671, 686; digestive system of, 455.

Macrosiphum sonchi, transmitting lettuce mosaic in Britain, 671,

672.

Macrosiphum taraxaci, on Taraxacum kok-saghyz and dandelion in Kazakstan, 44.

Macrosteles divisus, measures against, transmitting lettuce yellows in New Jersey, **536.**

Macrosteles sexnotata, possible vector of virus diseases of potato in Russia, 303.

Macrotermes gilvus, feeding on injured roots of tung nut trees in Java, 523.

macula-alba, Ceuthorrhynchus.

maculata, Macrosiphum (Macrosiphoniella); Nephrotoma (Pales). maculator, Pimpla.

maculatum, Piesma.

maculatus, Bruchus; Camponotus.

maculipennis, Plutella.

Madagascar, Aleurodids and virus diseases in, 153, 637; Hispids on rice in, 664; locusts in, 14, 62, 478, 543, 571; Tachinid introduced into Mauritius from, 611.

Madeira, Coccids of, 392.

madens, Tribolium.

Madras, miscellaneous pests in, 129, 561; beneficial insects and biological control in, 562, 693.

Magicicada septendecim, application of bioclimatics to study of, 327. Magnesium Arsenate, uses of, as

an insecticide, 82, 425.

Magnesium Carbonate, action of dusts of, on *Bruchus obtectus*, **531**; tests of, in baits for grasshoppers, **134**.

Magnesium Fluoride, tests of, in baits for grasshoppers, 134.

Magnesium Oxide, as a corrective for foliage injury by fluorine insecticide and Bordeaux mixture, 425.

Magnesium Sulphate, spraying tests with, against Heliothis armigera, 133; doubtful value of, in sprays against Leptinotarsa decembineata, 133, 158, 634; ineffective against other insects, 133, 368, 425, 431. magnicornis, Dirhicnus.

magus, Tremex.

Mahé, establishment of Exochomus

flavipes in, 335.

Mahogany, pests of, in India, 612; timber of, practically immune from termites in Porto Rico, 465.

maidis, Aphis (Rhopalosiphum).

Maine, sawflies on birch and spruce in, 275, 540; Liposcelis divinatorius in, 650; beneficial insects in, 540.

Maize, Heliothis armigera on, in S. Africa, 488; pests of, in Australia, 88, 389, 603; pests of, in Canada, 25–27, 295, 601; locusts damaging, in China, 451; Aphis maidis on, in Fiji, 641; pests of, in Italy, 174, 198, 295; pests of, in Kenya, 666; locust damaging, in Madagascar, 62; Pyrausta nubilalis on, in Manchuria, 340; pests of, in Philippines, 451, 452; pests of, in Porto Rico, 328, 501, 503; pests of, in S. Rhodesia, 124, 491, 492; P. nubilalis on, in Russian Union, 109, 148, 152, 229, 587; pests of, in U.S.A., 32, 133, 138, 178, 179, 180, 248, 284, 295, 340,

341, 376, 379, 392, 414, 430, 431, 456, 500, 533, 595, 600, 645; insects associated with, in Ohio, 280, 455; methods of protecting ears of, against pests, 328; varieties and time of planting of, in relation to pests, 178, 341, 533; virus diseases in, 88, 472; little affected by soil treatment with calcium arsenate, 380.

Maize (Stored), pests of, and their control, 24, 119, 667; (cobs), in baits, 67, 260; (meal or flour), in baits, 124, 125, 127, 287, 379, 595; reducing toxicity of cryolite dust to Heliothis armigera, 525.

Maize Aphis (see Sipha maydis).
major, Penthaleus; Polistes.

malabaricus, Phassus.

Malacosoma americana, mating and

oviposition of, 456.

Malacosoma disstria, in U.S.A., 377, 597, 646; on Acer saccharum, 377; parasites of, 597, 646.

Malacosoma fragilis, bionomics of, on aspen in Colorado, 528.

Malacosoma indica, on fruit trees in India, 577.

Malacosoma neustria, in Czechoslovakia, 145, 468, 469; in Holland, 205, 269; in orchards in Poland, 163; on forest and shade trees, 269, 468, 469; parasites of, 145, 205; measures against, 269.

Malaya, pests of *Derris* in, **539**; locusts in, **486**; summaries of plant quarantines in, **274**, **592**; Carabid sent to Samoa from, against *Oryctes rhinoceros*, **641**.

malefida, Feltia (Agrotis). Maleuterpes spinipes (phytolymus),

on Citrus in Australia and New Zealand, 21.

mali, Aphelinus; Contarinia; Leptothrips; Psylla (Psyllia).

malinellus, Hyponomeuta padellus.
Manchuria, miscellaneous pests in,
19, 340, 361; Aphomia gularis
imported into Germany from,
520.

Mandarin Orange (see Orange,

Mandarin).

Mangel, pests of, in Britain, 332; restrictions on importation of, into Britain, 356; Piesma quadratum and crinkle disease of, in Poland, 163, 351, 352.

mangifera, Symitha.

mangiferae, Cryptorrhynchus (Sternochetus).

Mango, Coccid on, in Egypt, 509;
Anomala expansa on, in Formosa,
363; pests of, in Netherlands
Indies, 227; Dacus psidii on, in
New Caledonia, 131; Symitha
mangifera on, in Seychelles, 335;
fruit-piercing moths damaging,
in Sierra Leone, 271; pests of,
in W. Indies, 239, 464.

mangoensis, Zizera (Zizina) labra-

dus.

Mangrove (see Rhizophora).

manicatus, Chirothrips.

Manihot dulcis var. aipi, new mite on, in Brazil, 290.

Manihot tweediana, Coccid on, in

Argentina, 23.

Manihot utilissima (see Cassava). manihotis, Bemisia; Dasynus. Manila Hemp (see Musa textilis). manilae, Ceraphron (Calliceras). manilensis, Locusta migratoria. Mantis, predacious on Monema

flavescens in Japan, 207.

Maple, new Coccid on, in Caucasus, 680; Alsophila pometaria on, in Canada, 663; Cantharis obscura on, in France, 623; pests of, in U.S.A., 74, 374.

Maple, Broad-leaf (see Acer macro-

phyllum).

Maple, Sugar (see Acer saccharum). Marasmia trapezalis, on maize in S. Rhodesia, 124.

margalaestriata, Setomorpha (see S. rutella).

Margarodes, new species of, in Brazil, 623.

Margarodes buxtoni crithmi, subsp. n., bionomics of, on Crithmum maritimum in France, 235.

Margaronia indica, on cucurbits in Fiji, **641.**

Margaronia pyloalis, liberation of Cedria paradoxa against, on mulberry in India, 612.

marginata, Petrusina (Ormenis). marginatorius, Exenterus.

marginella, Campsomeris.

mariana, Eulia.

maritimus, Pseudococcus.

Marlattia laricis, on larch in Quebec, 464.

marmorata, Parastasia.

maroccanus, Dociostaurus.

marshalli, Chalcodermus.

Maruca testulalis (on beans), in Fiji, 641; in Mauritius, 611; in Porto Rico, 435.

Maryland, miscellaneous pests in, 377, 599, 600; mushroom pests

in, 79, 380; Macrosiphum onobrychis on peas in, 497; Popillia japonica and its biological control in, 377, 533.

Massachusetts, Calomycterus setarius in, 600; hibernation of Heliothis armigera in, 430, 431; Hylastes rufipes in, 418; Psila rosae on carrot in, 496; Tetranychus telarius disseminating carnation rust in, 79.

Mastotermes darwiniensis, in

Australia, 545.

Matsucoccus, on pines in U.S.A. and Dominican Republic, 378, 392; new species of, 392.

matsushitai, Hylecoetus.

maura, Eurygaster. mauritanicus, Tenebroides.

mauritia, Spodoptera.

Mauritius, miscellaneous pests and biological control in, 611, 670.

maxwelli, Oxypleurites. maydis, Sipha.

Mayetiola destructor, in Morocco, 691; in New Zealand, 547, 548; in Russian Union, 145, 680; in U.S.A., 83, 327, 375, 378, 381; date of sowing and varieties of wheat in relation to, 83, 327, 375, 378; on other cereals, 691; bionomics of, 145, 547, 691; parasites and biological control of, 145, 375, 381.

mayri, Sphaerakis (Bruchobius). Meat, in baits for Carabid on strawberry, **466**.

Mecistomela corallina, on Cocos coronata in Brazil, 661.

medicaginis, Aphis; Contarinia. Medicago sativa (see Lucerne).

medinalis, Cnaphalocrocis.

Mediterranean Fruit-fly (see Ceratitis capitata).

Medlar, Cydia molesta on, in Argentina, 601.

megacephalus, Microphanurus (see M. basalis).

megalopygus, Dysdercus (see D. cingulatus).

Megaselia, on tung nut trees in Java, 523.

Megastigmus, species of, bred from galls in Australia, 565; study of, infesting conifer seeds in Germany, 230; subgenera of, 565.

Megastigmus abietis, in spruce seeds in Germany, 230.

Megastigmus ballestrerii, bionomics of, on pistachio in Greece and Sicily, **452**, **453**.

Megastigmus brevivalvus, bionomics of, in New South Wales, 564-566.

Megastigmus nonvitta, bionomics of, in New South Wales, 565.

Megastigmus spermotrophus, introduced into Germany in seeds of Douglas fir, 230.

Megastigmus trisulcus, bionomics of, in New South Wales, 565.

Megymenum gracilicorne, bionomics of, in Japan, 441.

Meigenia mutabilis, possible parasite of Leptinotarsa decemlineata in France, 194; parasite of Crioceris asparagi in Italy, 472; bionomics of, 194.

melanictera, Enaria.

melanis, Pipunculus confraternus. melanocephalum, Tapinoma.

Melanophila fulvoguttata, on Tsuga canadensis in U.S.A., 378.

Melanoplus (in U.S.A.), relations of outbreaks of, to rainfall and temperature, 469; experiments on flight of, 33, 646; natural enemies of, 75; baits for, 134, 537; plants damaged by species of, 178.

Melanoplus bivittatus, in U.S.A.,

537, 569.

Melanoplus differentialis, in U.S.A., 33, 75, 178, 537, 569.

Melanoplus femur-rubrum, in U.S.A.,

75, 178, 537, 569.

Melanoplus mexicanus (spretus), in Canada, 476, 485; in U.S.A., 33, 134, 178, 485, 569, 599, 646, 655; migrations and phases of, 476, 569; effect of tillage on, 655; organisation of measures against, 476.

melanorhodon, Anacridium moestum.

Melanostoma, attacking Aeolarchis sphenotoma in Fiji, 213.

Melanotus communis, on sugar-cane in Florida, **656.**

Melanotus tamsuyensis, bionomics of, in Formosa, 363.

Melanoxanthus fractus, on tung oil trees in Java, 523.

melanura, Idgia.

Melaphis chinensis, bionomics of, producing tanningalls in Formosa, 205.

Melasoma tremulae, on aspen in U.S.A., 599; penetration of fluids into tracheae of, 579, 580.

Melia azadirachta, mite on, in India, 128.

Meligethes aeneus (on crucifers), in Britain, 333; mobile traps for, in Germany, 511.

Meliphora grisella (see Achroia).

Melittia satyriniformis (Squash Vine Borer), measures against, in Wisconsin, 33.

Melittobia indicum (see Syntomosphyrum).

melliferum, Hyadaphis (Rhopalosiphum) (see H. sii).

mellitor, Microbracon. mellonella, Galleria.

Melolontha, in Czechoslovakia, 469, 471; in Germany, 171, 299, 300, 334, 398, 495, 496, 606; in Italy, 200; effect of temperature and humidity on, 300, 471; oviposition sites of, 299, 608, 676; natural enemies and diseases of, 100, 117, 300, 306, 307, 334, 398, 496, 518, 608; organised collection of adults of, 9, 171, 299, 606, 608; insecticides against adults of, 171, 299, 495, 606, 608; reactions of sexes of, to poisons, 406; other measures against, 299, 608; biological bases of measures against larvae of, 606-608; tests of calcium cyanamide against, 200; effect of flooding on, 457; action of HCN on, 309.

Melolontha hippocastani, distribution of, in Finland, 267; in Germany, 171, 299, 300, 606, 607, 608; in Poland, 676; in Ukraine, 406; life-cycle of, 9, 171; other bionomics and physiology of, 406, 607, 608; larva of, 224, 694.

Melolontha melolontha (vulgaris), parasite of, in Denmark, 117, 518; in Germany, 112, 171, 300, 457, 607, 608; in Holland, 269; in Poland, 9, 676; in Russia, 306; bionomics of, 112, 171, 607, 608; fungi infesting 306, 307.

Melon, Trypetids on, in Hawaii and Baluchistan, 344,577; Diaphania hyalinata on, in Porto Rico, 501; pests of, in U.S.A., 340, 500.

Melon Fruit-fly (see Dacus cucurbitae).

melvillei, Tropidophryne.

melzeri, Gryllica.

Membracis foliata (lunata), or Anonaceae in Brazil, 567.

membranaceus, Brachytrypes.

Membranes, method for investigating permeability of, 263.

mendax, Lygidea.

Hyalomyia mendesi. Dysdercus; (Phoranthella).

mendicus, Cleonus (Conorrhynchus). menippus, Rhaconotus.

Mentha (see Mint).

Mercurous Chloride, uses of, against Diptera infesting vegetables, 116, 497, 674, 675; applied to strawberry roots against Lachnosterna, tests of, in baits for 425; grasshoppers, 134; against termites, 181.

Mercury Bichloride, uses of, against Diptera infesting vegetables, 116, 317, 497; repellent to grass-

hoppers, 134.

Merhynchites, subgenus of Rhynchites (q.v.), 595.

meridionalis, Pyrausta aurata.

Merisus destructor, emergence holes of, indicating infestation of wheat by Mayetiola destructor in Kansas,

meritana, Epinotia. Mermis, in Dysdercus in Sudan, 574; in grasshoppers in U.S.A.,

Mermis subnigrescens, 76.

Merodon equestris (Narcissus Fly), bionomics of, in Holland, 494.

Meromyza saltatrix, and white ear in cereals in Russian Union, 680.

Meroprion anonicola, n., measures against, on Anonasquamosa, etc., in Brazil, 567.

Mescinia peruella, irrigation against, on cotton in Peru, 385.

Mesochorus, parasite of Vanessa cardui in Transcaucasia, 587; hyperparasite of Diprion similis in Poland, 671.

possibly Mesochorus petiolaris, hyperparasitic on spruce sawflies in Czechoslovakia, 518.

Mesolecanium deltae, measures against, on orange in Argentina,

Mesoleius tenthredinis, parasite of Pristiphora erichsoni in Quebec, 464.

messoria, Euxoa.

Metagonistylum minense (parasite of Diatraea), in Brazil, 505; utilisation of, in Br. Guiana, 505, 524; distribution and utilisation of, in W. Indies, 24, 182, 183, 505, 659; bionomics of, 182, 505.

Metal Shields, against termites, 181, 465, 694,

metallescens, Eutetrapha (Saperda). Metamasius sericeus, on sugar-cane in Jamaica, 238.

Metarrhizium anisopliae (infesting noxious insects), in Hawaii, 368; in Russia, 306, 684; in Sweden, 118; experiments on utilisation of, 118, 306, 684; penetration of chitin of insects by, 307.

Conditions Meteorological Humidity, Rainfall and Tem-

perature).

Meteorus, hosts of, in Holland, 205; parasite of Setora nitens in Java,

Meteorus japonicus (nipponensis), parasite of Lymantria dispar in Japan, 615; failure to establish, in U.S.A., 615; characters and synonymy of, 614.

Meteorus pulchricornis, parasite of Lymantria dispar in Europe, 615; parasite of Pyralid in Morocco, character distinguishing 193:

M. japonicus and, 615.

Meteorus versicolor, bionomics of, in Germany, 354; establishment of, against Stilpnotia salicis in Washington, 499.

Meth-allyl-chloride, temperature not

affecting toxicity of, 111.

Methyl Bromide, tests and uses of, as a fumigant, 359, 539, 541, 605: resistance of strains of Aonidiella aurantii to, 343; effects of, on apples and azaleas, 541, 542, 605; regulations dealing with fumigation with, against Popillia and Naupactus in U.S.A., 274. 591.

Methyl Salicylate, in repellent paint against pests of books, 366.

Methylated Spirit, in spray against Tingid, 359, 570; in repellent paint against pests of books, 366; tests of penetration of, tracheae of insects, 579, 580.

meticulosa, Brotolomia, metricus, Polistes.

Metriona bicolor, on bindweed in Kansas, 375.

Metriona bivittata, on bindweed in Kansas, 375.

Metriona flavolineata, measures against, on sweet potato in Jamaica, 239.

Metroxylon, Promecotheca reichei on, in Fiji, 366.

Metzneria, bionomics of, on Pyrus sinensis in China, 669.

Mexican Bean Beetle (see Epilachna varivestis).

mexicana, Paratrioza. mexicanus, Melanoplus.

Mexico, miscellaneous insects in, 143, 285, 569.

Mezium americanum, infesting house insulation in U.S.A., 640. micacea, Hydroecia.

micans, Stenomalus.

Mice, destroying noxious insects, 105, 217.

Michigan, Cirphis unipuncta in, 599; forest pests in, 28, 378, 461; Pyrausta nubilalis in, 414.

Microbracon, parasite of Pissodes notatus in Finland, 8.

Microbracon abscissor, parasite of Cydia glycinivorella in Japan, 624.

Microbracon brevicornis, ants hindering control of Heliothis armigera by, in S. Africa, 323.

Microbracon brevicornis, auct. (see

M. hebetor).

Microbracon chinensis, establishment of, against Chilo simplex in

Hawaii, 367.

Microbracon greeni, parasite of lac predators in India, 128, 129, 575; rearing and other hosts of, 128, 129; M. lefroyi considered a race of, 575.

Microbracon hebetor, in Fiji, 213; Germany, 400, 520, 521; liberation of, against lac predators in India, 129; in Korea, 225; parasitising Ephestia and Plodia, 213, 225, 400, other hosts of, 129, 520.

Microbracon lefroyi, parasite of cotton bollworms in India, 575; considered a race of M. greeni,

575.

Microbracon mellitor, probably parasitic on Chalcodermus aeneus in U.S.A., 77.

Microbracon nigrorufum, introduced into U.S.A. from Korea against

Platyedra gossypiella, 381. Microbracon pectinophorae, intro-duced into U.S.A. from Korea against Platyedra gossypiella, 381.

Microbracon piger, introduced into U.S.A. against Etiella zinckenella,

381.

Microbracon pygmaeus, parasite of Coleophora pruniella in Oregon,

Microbracon triangularis, parasite of Aegeria typhiaeformis in Italy, 630.

Microcerotermes, in Australia, 545. Microcerotermes arboreus, measures against, damaging furniture and timbers in Trinidad, 55; characters of, 55.

Microcryptus, introduced into U.S.A. from Canada against Diprion polytomum, 381.

Microcryptus basizonus, parasite of Diprion similis in Poland, 671.

Microcryptus brachypterus aphyopterus, parasite of spruce sawflies in Czechoslovakia, 518.

Microcryptus galactinus, parasite of spruce sawflies in Czechoslovakia,

518.

Microcryptus similis, parasite of spruce sawflies in Czechoslovakia, 518.

Microcryptus subguttatus, parasite of Diprion similis in Poland, 671.

Microdus diversus, introduced into Australia against Cydia molesta, 544.

Microdus nigrotrochantericus, sugar-cane fields in Florida, 659.

Microdus sanctus, possible parasite of Diatraea in Florida, 659.

Microdus stigmaterus (parasite of Diatraea saccharalis), utilisation of, in Florida, 656, 659; question of establishment of, in Porto Rico, 502.

parasite Microdus texanus, Lepidoptera in sugar-cane fields

in Florida, 657, 659.

Microgaster globata, parasite Tortricids in Transcaucasia, 587.

Microgaster psarae, parasite of Pycnarmon cribrata in Madras, 129. Microgaster subcompleta, parasite of

Vanessa cardui in Transcaucasia, 587.

Micromalthus debilis, bionomics of, in timber in gold mines in S. Africa, 60; morphology of, 61.

Micromyzus oliveri, food-plants and control of, in greenhouses in Washington, 541.

Microphanurus, keys to species of, 19, 632.

Microphanurus anitus, sp. n., parasite of Aelia acuminata in Germany, 632.

Microphanurus basalis, utilisation of, against Nezara viridula in

Australia, 292, 603.

Microphanurus choaspes, sp. n., parasite of Aelia acuminata in Germany, 632.

Microphanurus cultratus, parasite of Carpocoris pudicus in Germany, 632.

Microphanurus megacephalus (see M. basalis).

Microphanurus priapus, sp. n., hosts of, in Java, 19.

Microphanurus semistriatus, parasite of Pentatomids in Germany and Britain, 64, 632.

Microphanurus seychellensis, parasite of Cantheconidea robusta in Ceylon, 19.

Microphanurus sulmo, sp. n., parasite of Cantheconidea robusta in

Ceylon, 19.

Microphanurus vindicius, sp. n., parasite of Dasynus manihotis

in Java, 19.

Microplectron fuscipenne (parasite of Diprion spp.), utilisation of, in Canada, 276, 369; in Poland, 671; introduced into U.S.A., 381.

Microplitis cerurae, parasite Arctornis chrysorrhoea in Japan, 338.

microptera, Pararcyptera (Arcyptera); Romalea.

Microtermes, on injured tung oil trees in Java, 523.

Microterys, key to African species of, **503**.

Microterys bizanensis, sp. n., parasite of Saissetia cuneiformis in Eritrea, 503.

Microtervs nicholsoni, sp. n., parasite of Coccids in Natal, 503.

Microterys umbrinus, sp. n., parasite of Ceroplastes in Kenya, 503.

Mictis profana, food-plants of, in Fiji, 641.

middletonii, Aphis.

migratoria, Locusta (Pachytylus). migratorioides, Locusta migratoria. mikado, Miramella (Podisma).

miliaris, Asterolecanium.

Milionia basalis, on Pinus mercusi in Netherlands Indies, 578.

Milk (Dried), in sprays, 52.

Milky Disease (in Popillia japonica), methods of dissemination of, in U.S.A., **377.**

Millepedes, attacking tomatos in greenhouses in Britain, characters of economic species of,

Millet, Italian (see Setaria italica). Millettia, cultivation of, for insecticides in Belgian Congo, 294.

Millettia reticulata, insecticide prepared from, 450.

Mimastra cyanura, on fruit trees in India, 503.

mimulus, Dysdercus.

mimus, Dysdercus. minense, Metagonistylum.

Minnesota, forest pests and their parasites in, 434, 597, 599, 646; Cirphis unipuncta in, 599.

minor, Dryocoetes.

Mint, Jassid on, in greenhouses in Pyralid on, in Britain, 115; Morocco, 193.

minuta, Hoplocampa.

minutana, Enarmonia (Gypsonoma).

minutum, Trichogramma

minutus, Arrhenodes (Patysystro-phus); Dinoderus; Laemophloeus. Miramella mikado, in Japan, 618. Mirax, parasite of Leucoptera coffe-

ella in Guadeloupe, 435; introduced into Porto Rico, 435.

Miscanthus, sugar-cane pests on,

in Formosa, 363.

Mischocyttarus phthisicus, predacious on Laphygma frugiperda in Porto Rico, 503.

misella, Paroxyna; Tinea (see T.

insectella).

Mississippi, Chalcodermus aeneus in, 644; cotton pests in, 243, 599; Mymarid parasite of Psallus seriatus on Croton in, 538; Cylas formicarius in, 379; legislation against Naupactus spp. in,

Missouri, grasshopper outbreaks in, 569: miscellaneous pests in, 133.

597, 599.

missuriensis, Crematogaster victima. Mites, studies on, infesting stored grain, etc., in Russian Union, 312, 318, 319, 551, 585, 686; list of, on plants in Sweden, 355; disseminating carnation rust fungus, 79; attacking insects, 75, 86, 131, 169, 292, 400, 496, 672; attacking other mites, 319; natural enemies of, 410, 429, 603, 626; classification and new species of, 33, 34, 224, 288, 290, 301, 636, 672.

mitsukurii, Telenomus.

Mnium, Melaphis chinensis migrating to, in Formosa, 205.

modesta, Campsomeris marginella.

moestum, Anacridium.

Molasses (Treacle), in baits, 67, 68, 89, 98, 130, 240, 270, 287, 291, 293, 476, 478, 495, 510, 537, 602, 682; in bait-sprays, 232, 330; in sprays against thrips, 177, 425, 525; as an adhesive for lead arsenate, 661; attracting females of Cydia glycinivorella, 19: effect of, on fecundity of Agrotis segetum, 682.

Moles, destroying Protoparce spp.

in U.S.A., 251, 253.

molesta, Cydia (Grapholitha. Laspeyresia).

molestae, Angitia (Dioctes). molitor, Tenebrio.

moltshanovi, Macrosiphum (Acyrthosiphon).

mombin praeoptans. Anastrepha fraterculus.

Apate; monacha, Lymantria (Liparis, Psilura).

Monalonion, identity, bionomics and control of species of, on cacao, etc., in Brazil, 566.

Monalonion bahiense, 566. Monalonion bondari, 566.

Monalonion flavisignatum, sp. n., 566.

Monalonion knighti, sp. n., 566. Monalonion xanthophilum, 566. Monema flavescens, bionomics and control of, in Japan, 207.

monemae, Eurytoma.

Monima cruda (pulverulenta), feeding habits and parasites of, in Poland, 438.

Monoblastus, parasite of Pikonema alaskensis in Maine, 540.

monoceros, Oryctes.

Monochlornaphthalene, (See Naphthalenes, Chlorinated.) monodactylus, Oidaematophorus.

Monodontomerus, parasite of Nygmia phaeorrhoea in Holland, 205. Monodontomerus dentipes, hosts of, in Japan, 211; parasite Diprion similis in Poland, 671. parasite of

Monolepta divisa, on vines in

Queensland, 365.

Monolepta nigrobilineata (on soy beans), bionomics of, in Japan, **361**: in Manchuria, **340**.

Monolepta rosea, on vines in Queensland, 365.

Monolepta varicornis, on maize in Philippines, 452.

Monomorium pharaonis, in houses in Formosa, 446.

bionomics Monosteira unicostata, and control of, in Morocco, 359, 570.

monostigma, Scirpophaga.

Monotoma parallela, in stored cereals in Kansas, 600.

Montana, Melanoplus mexicanus in,

montanus, Pachynematus (Nematus). montei, Eurydinoteloides.

Monterey Cypress (see Cupressus macrocarpa).

montrouzierei, Cryptolaemus.

Moodna ostrinella, fumigation of cottonseed against, in Peru, 165. mori, Bombyx; Diplosis; Trionymus.

morio, Syagrus; Teleonemia.

Mormon Cricket (see Anabrus simplex).

Mormoniella vitripennis, bionomics

of, in Japan, 443.

Moroccan Locust (see Dociostaurus

maroccanus).

Morocco, beet pests in, 283; cereal pests in, 124, 691; Pyralid on mint in, 193; miscellaneous pests in, 58, 604, 691; orchard pests in, 283, 359, 570; Lepidoptera on poplars and willows in, 196; new mite on tomato in, 636; new Coccid in, Schistocerca gregaria in, beneficial insects in, 58, 193, 194, 196, 283, 284, 604; summary of plant quarantines in, 274.

morosa, Lachnosterna (Holotrichia).

morstatti, Xyleborus.

Mosaic (see Virus Diseases).

mosellana, Sitodiplosis (Thecodiplosis).

Mosquito Larvae, tests of insecticides on, 134, 135, 142, 241,

Moth-proofing Preparations, tests of. 140.

moultoni, Frankliniella.

Mucor, pathogenic to locusts in S. Africa, 92; infesting chestnuts in France, 358.

Mugho Pine (see Pinus mugo). Mugwort (see Artemisia vulgaris).

Mulberry, Aulacaspis pentagona on, in Brazil, 86; pests of, in Japan, 103, 555; Diplosis mori on, in Korea, 337; Margaronia pyloalis on, in India, 612; experimental preference of Diacrisia obliqua for, 130.

Mulberry Gall-midge (see Diplosis mori)

multistriatus, Scolytus.

Mundulea, survey of data on, as an insecticide, 104.

mundus, Bythoscopus (Macropsis).

mungomeryi, Eutachina.

Murgantia histrionica, measures against, on crucifers in Virginia, 530; alimentary system of, 280. murinana, Tortrix (Cacoecia).

murinus, Lacon (Brachylacon).

Murraya, Pulvinaria polygonata on,

in Formosa, 557.

Musa textilis (in Philippines), Pentalonia nigronervosa transmitting bunchy-top of, 365, 667; indicating areas favourable to Locusta migratoria manilensis, 485.

Musca, Dirhinus reared on, 212, 502.

Musca domestica, parasitised by Mormoniella vitripennis in Japan, 443.

Musca domestica vicina, parasitised by Spalangia cameroni in Fiji, 366.

Muscardine Fungus, White (see Beauveria bassiana).

muscarum, Stenomalus.

Muscideopsis lecanii, sp. n., parasite of Lecanium persicae in W. Australia, 294.

musculosa, Oria.

musculus, Lacon.

museorum, Anthrenus.

Mushrooms, characters and identity of mites on, in Britain, 1; pests of, in U.S.A., 79, 380; fumigation against pests of, 260.

musicus, Gastrimargus.

Mussel Scale (see *Lepidosaphes ulmi*). Mustard, pests of, in Lousiana, **250**, **373**.

mutabilis, Meigenia; Ootheca.

Myelois ceratoniae, measures against, in dates in Egypt, **506.**

Myelophilus piniperda, spring activity of, in Sweden, 123; frass and débris in galleries of, 391; forest management maintaining natural enemies of, 437.

myersi, Epitrimerus.

Myiopardalis pardalina, on melon in
Baluchistan, 577; intercepted in
cucurbits in Russian Union, 439.

Myiophasia globosa, parasite of Chalcodermus aeneus in U.S.A., 77, 82.

Mylabris dicincta, probably destroying eggs of Nomadacris septemfasciata in Tanganyika, 387.

Myllocerus, on tung oil trees in Java, 523.

Myristica fragrans (see Nutmeg). myristicae, Xyleborus.

Myrmelachista ambigua ramulorum, baits for, in Porto Rico, 435.

myron, Ampeloeca.

Mysore, miscellaneous pests in, 291, 625; beneficial insects in, 291.

mytilaspidis, Aphytis (Aphelinus). Mytilaspis (see Lepidosaphes). Mytilococcus (see Lepidosaphes). Myzella, considered distinct from Capitophorus, 99, 100.

Myzocallis ononidis, on cotton in

Russia, 215.

Myzocallis ulmicola, on elms in Japan, 443.

Myzodes (see Myzus).

Myzus cerasi (on cherry), in Australia, 20, 279; in Britain, 333; in New York, 535; measures against, 20, 279, 535.

Myzus circumflexus, relation of, to virus diseases of plants in U.S.A.,

83, 84, 369, 590.

Myzus convolvuli (see Macrosiphum solani).

Myzus hieracii, experiments with lettuce mosaic and, in Britain, 671.

Myzus lactucae (see Macrosiphum ribicola).

Myzus ornatus, transmitting virus disease of cucumber in Germany,

Myzus persicae, in S. Africa, 360: in Australia, 20, 448; in Britain, 236, 563, 578, 672; in Czechoslovakia, 468; in France, 622; in Germany, 157, 267, 301, 334, 633, 686, 687; in Holland, 578; in India, 228; in Netherlands Indies, 493; in Japan, 210; Indies, 493; in Japan, 210; in S. Rhodesia, 125, 226; in Russia, 215, 302; in U.S.A., 83, 142, 274, 369, 590; on cotton, 215; on peach and apricot, 20, 157, 236, 267, 468, 578, 633, 687; and potato virus diseases, 83, 157, 236, 267, 301, 302, 334, 563, 578, 633, 686, 687; on tobacco, 125, 210, 360, 633; and virus diseases of tobacco, 125, 226, 302, 493; on other plants, 125, 578, 622, 672, 687; and virus diseases of other plants, 274, 282, 448, 500, 674, 686, 687, 369, 448, 590, 671, 686, 687; characteristics of bean and tomato leaves unfavourable to, 142; flights of, 226, 267, 301, 578, 633, 687; other bionomics of, 267, 578, 622, 687; insects predacious on, 226, 228, 687; measures against, 20, 125, 157, 226, 236, 448, 578, 687.

Myzus polygoni-yonai, sp. n., on Polygonum in Japan, 104.

Myzus pseudosolani (see Macrosi-phum solani).

N.

Naaki, **120, 582.**

Nabis alternatus, predacious on Macrosiphum onobrychis in Utah, 30.

Nabis ferus, predacious on noxious insects in Utah, 30, 463.

Nacoleia octasema, measures against, on banana in Fiji, 213.

nana, Trichogrammatoidea.

Naphthalene, against carpet beetles and clothes moths, 188, 635, 636; against mites in beet seed, 686; uses of, against wireworms, 97, 177, 178, 378; killing eggs and larvae of *Psila rosae*, 497; as a repellent, 71, 116, 211, 299, 420, 497, 522, 608; increasing effect of wool grease emulsions on eggs of Cydia pomonella, 426; plants injured by, 71, ovicidal properties of derivatives of, 331, 332.

Naphthalenes, Chlorinated, tests of, as ovicides, 1, 332; bands with, against treated Cydia molesta, 639.

napi, Pieris.

Narcissus, pests of, in Britain, 512; Merodon equestris on, in Holland, 494; pests of, in U.S.A., 652, Aphids transmitting 654; mosaic disease of, 652.

Narcissus Fly (see Merodon

equestris).

nasturtii, Contarinia.

nasuta, Prorops.

Nasutitermes (see Eutermes). natalensis, Tropidophryne.

Naupactus, legislation against, in U.S.A., 275, 591.

Naupactus leucoloma, in U.S.A., 259, 274, 275, 378, 591; bionomics of, 275, 378; measures against, 259, 378, 591.

navale, auct., Tribolium (see

castaneum).

Nebraska, Aphids on wheat in, 542. Necrobia ruficollis, habits and distribution of, 584.

Necrobia rufipes, habits and distribution of, 212, 213, 584.

Necrobia violacea, habits and distribution of, 584.

Nectarine, Tortrix sarcostega on, in Baluchistan, 51; Dicyphus on, in New South Wales, 449.

negligens, Acritocera.

(Pteronematicida, Coelopisthia malus) (see Tritneptis klugi).

Nematodes, infesting insects, 38, 39, 76, 92, 100, 165, 214, 338, 347, 348, 364, 500, 519, 574; method of sampling soil for, 317.
Nematospora, Rhynchota associated

with, on plants, 388, 389.

Nematospora gossypii, experiment with, in cotton bolls, 491.

Nematus abietinus, erichsoni laricis (see Pristiphora).

Nematus montanus (see Pachynematus).

Nematus proximus, characters of, 6. Nematus purpureae, on Salix purpurea in Germany, 6; characters of, 6.

Nematus ribesii, bionomics of, on currant and gooseberry in Ger-

many, 114.

Nemeritus canescens, parasite of Ephestia kuehniella in Germany, 400; bionomics of, parasitising Plodia in Italy, 173; olfactory conditioning and host selection in, **264**, **265**.

Nemoraea, parasite of Laphygma exempta in Kenya, 666.

nemorata, Phyllotoma.

Nemorilla floralis, parasite Ancylis comptana in Kansas, 376; parasite of Pyrausta aurata in Morocco, 193.

nemorum, Phyllotreta. nenuphar, Conotrachelus.

Neoaplectana, 39.

Neoaplectana glaseri, in Popillia japonica in U.S.A., 165.

Neocerata rhodophaga, on rose in U.S.A., 600.

Neochryopus savagei, predacious on Oryctes in Dahomey, 347. Neodiprion (see Diprion).

Neodiscodes, African species of, 503. Neogastrallus librinocens, new para-

site of, in Florida, 461. silveirai, measures Neolecanium

against, on vines in Brazil, 34. Neomaskellia bergi, not transmitting dwarf disease of sugar-cane in Queensland, 186.

neomexicanus, Aphis.

Neosyrista similis, bionomics and control of, on rose in Japan, 210, 211.

Neowashingtonia Washing-(see tonia).

Nephantis serinopa, liberation of Trichospilus pupivora against, on coconut in Ceylon, 638.

Nephopteryx pirivorella (see Numo-

nia).

Nephotettix bipunctatus cincticeps, transmitting dwarf disease of rice in Japan, 362; oviposition of, 362.

Nephrotoma maculata, damaging cereals and pastures in Germany, 519.

Nephrotoma virgata, on tobacco in Japan, 210.

nesiotes, Eurytoma.

Nestlera, relation of Locustana pardalina to, in S. Africa, 394.

Netherlands Indies, coffee pests in, 161, 522; cotton pests in, 280; Stromatium longicorne in furniture in, 493; Drepanid on gambier in, 385; miscellaneous pests in, 48, 227, 523; new Scolytids on nutmeg in, 512; pests of Pinus mercusi in, 577, 578; Lepidoptera on tea in, 94, 347; tobacco pests in, 17, 493, 637; Tineid in stored tobacco in, 50; locusts in, 478, 486; termites in, 384, 523; insects and virus diseases in, 153, 493, 637; beneficial insects in, 19, 94, 347, 385, 386; beneficial insects introduced into other countries from, 19, 205, 366, 506, 665.

Neurotoma flaviventris, bionomics

of, in Germany, 114.

neustria, Malacosoma (Clisiocampa). New Britain (see New Guinea

Territory).

New Brunswick, Alsophila pometaria on elm, etc., in, 662-664; Blissus hirtus in, 601; utilisation of parasites against Diprion polytomum in, 369.

New Caledonia, Dacus psidii in, 131; Teleonemia shipped from Fiji to, against Lantana, 212.

New Guinea Territory, cacao pests in, 277, 544; parasites and biological control of coconut pests in, 22, 277, 294, 366, 506.

New Hampshire, Cirphis unipuncta in, 599; Malacosoma disstria in, 377.

New Hebrides, beneficial insects introduced into, 212, 366; failure to introduce *Pleurotropis parvulus* into New Guinea from, 366.

New Jersey, apple pests in, 536, 642; forest pests in, 74, 77, 279, 600, 651; European Cerambycid in building timber in, 600; Macrosteles divisus and lettuce yellows in, 536; Pyrausta nubilalis in, 179; Tiphia popilliavora and its utilisation against Popillia japonica in, 84, 377; Tortilia viatrix in stored senna in, 29.

New Mexico, Cydia pomonella in, 31, 273; Dendroctonus convexifrons in, 599; breeding areas of

Eutettix tenellus in, 593, 594; Platyedra gossypiella in, 381.

New South Wales, Cosmopolites sordidus on banana in, 184, 646; Eurytoma fellis on Citrus in, 564; forest pests in, 546, 602; orchard pests in, 57, 279, 293, 345, 449, 544, 640; insects and virus disease of Passiflora edulis in, 448; vegetable pests in, 58, 449, 545, 639; Eutane terminalis in, 329; grasshoppers in, 56, 130; beneficial insects in, 56, 279, 293, 564, 640; Angitia fenestralis introduced into, 545; establishment of Teleonemia against Lantana in, 546; Syntomosphyrum indicum introduced into Queensland from, 602.

New York, forest pests in, 68, 69, 77, 190, 279, 529, 530, 600, 651; Dutch elm disease in, 77; Hydroecia immanis on hops in, 593; miscellaneous pests in, 425, 599, 635; orchard pests in, 77, 255, 382, 533, 534, 535; pests of ornamental plants in, 79, 499, 652; vegetable pests in, 53, 274, 531; Aphids and virus diseases in, 273, 274, 652; new mealybugs

in, 48.

New Zealand, cereal pests in, 547, 548; Citrus pests in, 21, 56, 277, 278, 523, 524; orchard pests in, 159, 292, 548, 550; pests of pines in, 504; Tineid on raspberry and blackberry in, 37; Smynthurus viridis in, 550; host range and transmission by thrips of spotted wilt in, 546; American Buprestid in, 329; natural enemies and biological control of insects in, 37, 159, 262, 504, 548, 551; biological control of noxious plants in, 449; summary of plant quarantines in, 274; beneficial insects introduced into Australia from, 545, 546.

Nezara, vector of Nematospora, 389. Nezara chloris (see Aethemenes).

Nezara viridula, utilisation of Microphanurus basalis against, in Australia, 292, 603; on tobacco in Netherlands Indies, 493.

Nicandra physaloides, Myzus persicae on, in S. Rhodesia, 125; possible reservoir of tobacco rosette, 125.

nicevillei, Glyptomorpha (Bracon). nicholsoni, Microterys.

nicolasi, Campylomma.

Nicotiana spp., Phthorimaea operculella on, in Brazil, 289 : experiments with virus diseases and, **444, 546.** (See Tobacco.)

Nicotine, 593; against Aphids, 29, 157, 226, 236, 257, 332, 333, 353, 371, 395, 428, 448, 458, 645, 674, 687; against Aspidiotus perniciosus, 690; against Jassids, 115, 281, 413; against other Rhynchota, 232, 359, 449, 530; against 156, Coleoptera, 383, 623; against Diptera, 297, 353, 514, 519; effect of drenching mushrooms with, against Sciara, 380; against Lepidoptera, 103, 516, 663, 689; studies on fixed compounds of (including nicotinebentonite), against Cydia spp., 133, 272, 273, 525, 534, 536, 544, 545, 643, 653; against thrips, 233, 357, 365, 526; tests of ovicidal properties of, action of, on bees, 101, 332; 402; dusting with, 101, 115, 365, 371, 395, 428, 449, 674; as a greenhouse fumigant, 115, 357, 526; applied by vaporisation, 29, 395, 413; formulae containing, 103, 115, 233, 281, 333, 353, 359, 365, 371, 395, 448, 458, 514, 530, 674, 689; and methylated spirit, 359; and oils, 103, 332, 413, 458, 530, 534, 689; and pyrethrum, 449; and soaps, 115, 233, 332, 359, 395, 448, 458, 514, 519, 530; and sodium chlorate, 333; and tar distillates, 689; new activator for, 645; effects of, on carnations and azaleas, 526, 605.

Nicotine Sulphate, against Aphids, 125, 151, 258, 274, 278, 279, 371, 428, 429, 536, 541, 645, 662; against Coccids, 278, 432; against Jassids, 292, 413, 429, 432, 501; against other Rhynchota, 115, 125, 463, 561, 570, 662, 690; against Cydia pomonella, 89, 133, 186, 272, 432, 525, 536, 643, 653; against other Lepidoptera, 33, 126, 179, 376, 529, 535, 662; action of, on eggs of Chilo simplex, against Diptera, 58, 380, 555, 556, 628; against mites, 410, 429; against sawflies, 21, 275, 540; against thrips, 56, 71, 72, 127, 177, 365, 410, 425, 498, 525; action of, on bees, 101; dusting with, 21, 179, 371, 463, 470, 561; carriers for, 179, 371; formulae containing, 33, 56, 71, 89, 115, 125, 133, 177, 179, 186, 278, 279,

292, 365, 429, 432, 498, 525, 529, 535, 536, 540, 541, 570, 690; and bentonite, 133, 179, 272, 525, 643, 653; adsorption of, by other siliceous earths, 133; and Bordeaux mixture, 125, 501, 536; and calcium arsenate, 529; addition of chloropicrin to dusts Ceuthorrhynchus of, against macula-alba, 470; and lead arsenate, 125, 279, 292, 432, 529, and lime-sulphur, 292, **429, 498, 536**; and methylated spirit, 570; and molasses, 127, 425, 525; and oil emulsions, 56, 58, 89, 186, 278, 376, 413, 429, 432, 498, 529, 535, 536, 690; and colloidal sulphur, 186; and soap, 115, 151, 177, 258, 278, 365, 498, 540, 541, 570; other spreading and wetting agents for, 33, 71, 89, 179; activators for, 177, 645; adhesives for, 272, 525, 643, 653; other insecticides compared with, 71, 72, 127, 177, 536, 540.

nidulans, Eupteromalus.

Anagyrus; Coelinius: niger, Lasius; Orius (Triphleps).

Nigeria, Aleurodids transmitting cassava mosaic in, 153; summary of plant quarantines in, 592. nigeriensis, Bemisia.

Nightshade, Black (see Solanum nigrum).

nigra, Phytomyza; Saissetia.

nigrescens, Anagyrus.

nigricana, Cydia (Laspeyresia).

nigricornis, Phytomyza (see affinis); Trioza.

nigridens, Phaeogenes. nigrifemur, Perisierola.

nigripennis, Cryptogonus orbiculus.

nigripes Lydella (Anetia).

nigrobilineata, Monolepta (Para-

luperodes). nigrofasciatus, Dysdercus.

nigronervosa, Pentalonia. nigrorufum, Microbracon.

nigrotrochantericus, Microdus (Bassus).

nigrum, Leptomastix.

Nilaparvata oryzae, oviposition of, on rice in Japan, 362.

nipponensis, Meteorus (see

japonicus). nipponica, Acantholyda.

nirecola, Chromaphis.

nitens, Setora. nitida, Revena (Amercedoides).

Nitrobenzene, in oil emulsions against vine moth pupae, 103.

nivella, Scirpophaga.
niveus, Oecanthus.
nobilis, Lebeda; Rhizophagus.
noctilio, Sirex (Paururus).
noctiluca, Lampyris.
noctuae, Zenillia.
nodiceps, Cryptognatha.

Nomadacris septemfasciata, in Fr. W. Africa, 570, 571; in Portuguese E. Africa, 62, 479, 484; in S. Africa, 91, 92; in Belgian Congo, 476, 482, 483, 486; in Kenya, 666; in Madagascar, 478, 543, 571; in Nyasaland, 62, 126, 388; in N. Rhodesia, 483, 484; in S. Rhodesia, 124, 637; in Tanganyika, 387, 480, 484; in Uganda, 480, 666; breeding centres and ecology of, 62, 387, 388, 479, 480, 483, 484, 570, 571; phases of, 387, 388, 482, 483, 486, 570, 571; migrations of, 126, 387, 388, 476, 479, 484, 666; fungi infesting, 91, 92; other natural enemies of, 92, 387; measures against, 92, 479, 484, 637; international organisation for control of, 483, 484.

nonvitta, Megastigmus (Epibootania). Noogoora Burr (see Xanthium pungens).

nordmannianae, Chermes.

Norfolk Island, introduction of Teleonemia into, against Lantana, 546.

Norta altissima, relation of Eutettix tenellus and curly-top virus to, in Idaho, 286.

norvegicus, Calocoris.

Norway Spruce (see *Picea abies*). *Nosema*, in insects in S. Africa and Brazil, **459**; host relations of, **459**; study of new species of, in

Cactoblastis spp., 459. Nosema cactoblastis, sp. n., 459. Nosema cactorum, sp. n., 459. notabilis, Aesiotes.

notata, Chloropisca. notatus, Pissodes.

Notodacus (see Dacus).

Nova Scotia, apple pests in, 662; Blissus hirtus damaging lawns in, 601; Alsophila pometaria on elm, etc., in, 662-664.

Novius cardinalis (see Rodolia). Novosil (containing dinitro-orthocresol q.v.), 400.

noxius, Brachycolus. noyesi, Eloria. nubilalis, Pyrausta. nubilus, Helophorus. nucicolora, Elaphria. nucleorum, Pachymerus. nucum, Curculio (Balaninus); Pimpla.

Numonia pirivorella, on Pyrus sinensis in China and Korea, 668. Nun Moth (see Lymantria monacha). Nurudea shiraii, producing tannin galls on Rhus in Formosa, 205.

nüsslini, Dreyfusia (see Chermes nordmannianae).

Nutgrass (see *Cyperus rotundus*). Nutgrass Armyworm (see *Laphygma exempta*).

Nutmeg, new Scolytids on, in

Sumatra, 512.

Nyasaland, cotton pests in, 125, 126, 222; miscellaneous pests in, 126; Nomadacris septemfasciata in, 62, 126, 388; beneficial insects in, 126.

Nygmia phaeorrhoea, on oak and fruit trees in Czechoslovakia, 468, 469; parasites of, in Holland, 205; in Poland, 163; measures against, in U.S.A., 139, 180.

Nymphaea alba, Galerucella nymphaeae on, in Italy, 3.

nymphaeae, Galerucella.

Nymphula depunctalis (on rice), in Mysore, 291; in Philippines, 364, 451; bionomics of, 364; measures against, 291, 364.

Nymphula vittalis, bionomics and control of, on rice, etc., in Korea,

618, 619.

Nysius, type of injury to wheat by, in New Zealand, 548.

Nysius turneri, bionomics and control of, in Tasmania, 226, 227.

Nysius vinitor, measures against, on vegetables and stone-fruits in New South Wales, 449.

O.

Oak (Quercus), pests of, in Australia, 546; adaptation of pests of, to rhododendrons in Britain, 295; Alsophila pometaria on, in Canada, 663; pests of, in Czechoslovakia, 468, 469; chestnut pests on, in France, 358; pests of, in Germany, 204, 299, 354, 522; Thaumetopoea processionea on, in Italy, 516; pests of, in Poland, 438; pests of, in Russia, 43, 456; Dryocoetes minor on, in Sicily, 522; pests of, in U.S.A., 32, 645; weight of foliage of, in relation to insect control, 143; insects experimentally fed on, 229, 399, 406.

Oak Silkworm, Chinese (see Antheraea pernyi).

Oak Timber Worm (see Arrhenodes

minutus).

Oats, pests of, in Australia, 21, 545; pests of, in Britain, 96, 97, 122; Tipulids not completing development on, in Germany, 519; pests of, in Japan, 208, 556; Calliptamus italicus damaging, in Poland, 9; pests of, in Russian Union, 322, 454, 680; Limothrips cerealium on, in Sweden, 121; pests of, in U.S.A., 284, 599; insects and white ear in, 680; Mayetiola destructor not developing on, 691; soil treatment with calcium arsenate affecting yield of, 380.

Oats (Stored), pests of, and their control, 64, 119, 318, 319.

Oberea linearis, on hazel in Greece,

oberthüri, Catocala.

oblineatus, Lygus pratensis.

obliqua, Diacrisia.

obliquus, Listroderes. oblongus, Phyllobius.

obscura, Cantharis; Leucopis;

Rhabdocnemis. obscurata, Brachymeria.

obscurator, Orgilus.

obscurella, Brachonyx pineti.

obscuripennis, Rhyparida.

obscurus, Agriotes; Anaphothrips; Athous; Chrysomphalus.

obsoleta, Heliothis (Chloridea) (see H. armigera).

obtectus, Bruchus (Acanthoscelides).

ocellana, Spilonota (Eucosma).

ocellatus, Paniscus; Pristomerus; Smerinthus.

Ochrogaster contraria, natural enemies of, on Acacia in Australia, 545.

Ochrosidia immaculata (see Cyclo-

cephala).

Ochsenheimeria taurella, causing white ear in cereals and Bromus in Russian Union, 680.

octasema, Nacoleia.

octocinctus, Stenaraeoides.

octomaculata, Saperda.

oculata, Chrysopa.

oculiperda, Thomasiniana.

Odontopus, vector of Nematospora,

Odontotermes (see Termes).

Occanthus niveus, feeding habits of, in Utah, 30; tests of barium fluosilicate on, 139.

Oedaleus australis, in Australia, 545.

Oedipoda, in Jugoslavia, 480.

Oedipoda coerulescens, in Poland, 9.

Oemona hirta, bionomics of, on Citrus in New Zealand, 21.

Oenophthira pilleriana (see Sparganothis).

oestracea, Trichopsidea.

Ohio, pests of beans in, 82, 421; cereal pests in, 82, 414; insects associated with maize in, 280, 455; Scolytus multistriatus and Dutch elm disease in, 378, 599; spray programmes for fruit crops in, 104, 623; potato pests in, 342, 421; Chelonus annulipes introduced into Florida from, against Diatraea saccharalis, 656.

Oiceticus abboti, on Citrus in

Jamaica, 181.

Oidaematophorus monodactylus, on bindweed in Kansas, 375.

Oil, Camphor Sassafras Base, as intermediary solvent for derris in oil sprays, 530.

Oil, Castor, tests of penetration of, into tracheae of insects, **579**, **580**.Oil, Cedar-wood (see Cedar-wood Oil).

Oil, Cinnamon (see Cinnamon Oil).
Oil, Coconut, and derris extract
against thrips, 493.

Oil, Cottonseed, emulsions of, as carrier for paradichlorobenzene,

Oil, Ground-nut, as an insecticide, 168, 530, 593; and derris extract against thrips, 493.

Oil, Linseed, emulsion of, against *Tetranychus telarius*, **45**; effect of emulsions of, on pear buds, **94**; timber treated with, against carpenter bees, **337**; as adhesive for lead arsenate, **53**, **430**, **540**.

Oil, Olive, emulsions of, against Aphids and Coccids, 6; in bait

for Dacus oleae, 98.

Oil, Pine-tar, as an insecticide, 593. Oil, Soy-bean, forms of nicotine-bentonite containing, 272, 643, 653.

Oil Emulsions, against Aleurodids, 25, 259, 530, 560, 642, 649; against Aphids, 20, 236, 237, 429, 450, 458, 460, 467, 517, 529, 534, 535, 536, 550, 662; against Coccids, 23, 25, 34, 55, 66, 103, 128, 130, 164, 238, 255, 256, 259, 278, 292, 297, 432, 433, 435, 460, 467, 508, 509, 524, 529, 534, 535, 553, 649, 662, 681, 690; against Capsids, 45, 236, 237, 566;

against other Rhynchota, 25, 353, 413, 432, 467, 535, 690; against Coleoptera, 426, 450; against Cydia spp., 20, 57, 89, 166, 186, 192, 293, 432, 534, 536; against other Lepidoptera, 28, 33, 103, 236, 237, 376, 450, 529, 535, 689; against Diptera, 58, 297, 450, 497; against mites, 2, 55, 56, 57, 103, 257, 409, 429, 430, 432, 460, 535, 536, 550, 649; against sawfles, 461; against thrips, 56, 433, 408; against sawfles, 56, 438, 460; against sawfles, 56, 488, 480; against sawfles, 56, 488, 56, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, 57, 56, destroying beneficial insects, 430, 549, 662; types of oils for, 2, 20, 28, 56, 57, 58, 89, 94, 103, 128, 130, 164, 166, 186, 188, 192, 237, 255, 259, 278, 292, 293, 297, 332, 353, 376, 409, 426, 429, 430, 432, 435, 450, 458, 460, 498, 509, 524, 525, 526, 527, 529, 530, 534, 535, 536, 553, 560, 566, 626, 627, 642, 649, 653, 662, 681, 689, 690; factors affecting deposits of, 409, 411, 508; preparation of types of, 432, 508, 626; emulsifiers and spreaders for, other than soap, 70, 278, 332, 411, 529, 626, 627, 642, 681; formulae for, 2, 20, 23, 28, 33, 56, 57, 66, 89, 94, 103, 130, 255, 278, 292, 293, 409, 429, 430, 432, 435, 458, 460, 509, 524, 526, 527, 529, 534, 535, 536, **549, 550, 681**; formulae containing, 28, 56, 57, 89, 103, 192, 236, 259, 278, 293, 409, 429, 432, 450, 458, 460, 461, 497, 498, 509, 529, 530, 534, 535, 536, 553, 690; and aniline, 103; and Bordeaux mixture, 188, 192, 497, 536; and calcium arsenate, 529; and carbon bisulphide, 553; derris, 33, 530, 642, 689; and dinitro-cresol, 236, 332, 534; and dinitro-o-cyclohexyl-phenol, 429, 529, 530, 534, 535, 649; and lead arsenate, 28, 139, 166, 188, 190, 192, 293, 376, 426, 430, 432, 461, 525, 529, 536; and limesulphur, 57, 409, 430, 432, 460, 536; and nicotine, 56, 58, 89, 103, 186, 278, 332, 376, 413, 429, 432, 458, 498, 529, 530, 534, 535, 536, 689, 690; and nitrobenzene, 103; as a carrier for paradichlorobenzene against borers, 526, 527 and pyrethrum, 28, 376, 450, 689; and sulphur, 57; and tar distillates, 236, 534, 535; and thiocyanates, 236, 237, 259, 332; value of, in combined postdormant washes, 396; effects of, on plants, 45, 94, 103, 255, 430, 435, 458, 460, 508, 509, 529, 536, 653, 680; interactions of, with other sprays on Citrus, 278, 279; effect of sprays of, on removal of spray residues from fruit, 189, 192, 430; tar distillates compared with, 429, 467, 468, 509, 529; other insecticides compared with, 2, 237, 458.

Oil Palm (see Elaeis guineënsis).

Oils, in baits, 67, 260; spraying with, against locusts, 618; water in rice fields treated with, against pests, 365, 613, 665; microtechnique for testing action of, on Coccids, 143; in mixture for treating timber against pests, 181, 337; atomised sprays of insecticides in, 154, 155, 412, 413; autogiro for applying formulae containing, 74, 75; dusts impregnated with, 75, 308; as an adhesive for cryolite dips, 380, 652; and beta-naphthol, for treating bands against Cydia spp., 197; in formula for removing spray residues from fruit, 190. (See Kerosene.)

Oils, Sulphonated (see Ultrawet).
Oklahoma, Eutettix tenellus on beet in, 594; cereal pests in, 261, 274, 599; miscellaneous pests in, 79, 340; Pemphigus on Poinsettia in greenhouse in, 257; Chrysobothris femorata in, 420; Melyrid predacious on Blissus leucopterus in, 261; Empusa grylli infesting grasshoppers in, 427.

Okra (see Hibiscus esculentus).

oldenlandiae, Theretra.

oleae, Dacus; Parlatoria (Syngenaspis); Saissetia.

oleellus, Prays.

Oleic Acid, and ammonia in nicotine spray, 514; Stantex Dispersing Oil containing, 530.

oleivorus, Phyllocoptruta (Phyllocoptes).

olens, Staphylinus.

oleracea, Tipula.

oleraceum, Eurydema. Olfactometers, 264, 265, 391.

Olibrus bicolor, measures against, on Taraxacum kok-saghyz in Kazakstan, 45.

olidus, Eutermes.

Oligonychus ulmi, auct. (sée Paratetranychus pilosus).

Oligotrophus bergenstammi (see Apiomyia).

Olive, new Eriophyid on, in California, 288; identity of Geometrid on, in Crete, 171; pests of, in Cyprus, 270; Acherontia atropos on, in Egypt, 175; pests of, in Greece, 453; pests of, in Italy, 98, 473, 568; bulletin on pests of, in Turkey and other countries, 452; Dacus oleae intercepted in fruit of, in Russian Union, 439.

Olive Fruit-fly (see *Dacus oleae*). Olive Moth (see *Prays oleellus*).

Olive Oil (see Oil, Olive).

oliveri, Micromyzus.

olivieri, Polyphylla. olivina, Euphyllura.

Omaseus vulgaris (see Pterostichus). omissa, Tischeria.

Omotestus haemorrhoidalis, in Poland, 9; ecology of, in Russia, 386.

Oncideres amputator, on ornamental trees in Jamaica, 181.

Oncocephalus geniculatus, predacious on Horistonotus uhleri in S. Carolina, 500.

Oncopera, attempted establishment of *Hexamera* against, in Australia, 545.

Onion, Hylemyia antiqua on, in Britain, 674; Aphids transmitting virus diseases of, in Germany, 686; Thrips tabaci on, in Porto Rico, 436; pests of, in U.S.A., 71, 372, 378, 541; planted among carrots against Psila rosae, 497; insecticides causing injury to, 71, 72; legislation dealing with fumigation of bulbs of, against Popillia japonica in U.S.A., 274.

Onion Fly (see Hylemyia antiqua). Onion Thrips (see Thrips tabaci).

Onobrychis sativa, source of infestation of cotton by Aphis laburni in Russia, 216.

onobrychis, Macrosiphum.

ononidis, Myzocallis (Therioaphis).
Ontario, Pyrausta nubilalis on
maize in, 25-27; Cerosipha
forbesi on strawberry in, 276;
Forficula auricularia in, 601.

Ocencyrtus clisiocampae, parasite of Malacosoma disstria in Minne-

sota, **597.**

Ocencyrtus kuvanae, factors determining choice of unparasitised eggs of Lymantria dispar by, 302. Octheca mutabilis, on cowpeas and soy beans in S. Rhodesia, 125.

opacicollis, Promecotheca.

opalus, Pachnaeus.

Opatrum sabulosum, baits for, on Perilla, etc., in Ukraine, 321.

805

Opatrum triste, bionomics and control of, on tobacco, etc., in Crimea, 41, 42.

operculella, Phthorimaea (Gnori-

moschema).

Operophtera brumata, in Britain, 236; in Germany, 99, 112, 398, 513; on oaks in Poland, 438; on fruit trees, 236, 398, 513; unable to develop on peach, 513; bionomics of, 99, 112; reactions of, to temperature, 40; measures and tests of insecticides against, 236, 237, 332, 398, 438.

Ophiomyia (see Agromyza).

Ophiostoma ulmi (causing Dutch elm disease), distribution of, in Britain, 65; in Holland, 396; in U.S.A., 65, 68, 77, 138, 180, 378, 529, 645; relation of Scolytids to, 65, 68, 77, 378, 396, 529; destruction of elms against, 378, 645.

Ophonus pubescens (see Harpalus

rufipes).

Opilo discodirus, predacious on Hypsipyla robusta in India, 612. Opius, parasite of Anastrepha

fraterculus in Brazil, **34.**Opius aino (see Diachasma).

Opius anastrephae, parasite of Anastrepha spp. in Porto Rico, 502.

Opius fijiensis (see Biosteres).
Opius fletcheri, parasite of Dacus
in Fiji, 213.

Opius ilicis, sp. n., parasite of Phytomyza ilicis in Br. Isles, 669. Opius incisi, parasite of Dacus in Madras, 562, 577.

Opius longicaudatus (see Biosteres). Opius persulcatus (see Biosteres). Opsimus quadrilineatus, in building

timber in U.S.A., 499.

Opsophyto arteagai, parasite of Trigonophymus arrogans in Argentina, 214.

Opuntia spp., failure to establish Cactoblastis doddi against, in S. Africa and Australia, 459; utilisation of other insects against, in Queensland, 344, 345.

opuntiae, Dactylopius.

Oraesia emarginata (see Calpe).
Orange, locusts damaging, in Fr. W.
Africa, 571; pests of, in S.
Africa, 330; Mesolecanium deltae on, in Argentina, 238; Ceratitis capitata on, in Cyprus, 270;

Pseudococcus citri on, in France, 266; Chrysomphalus rossi on, in New Zealand, 524; C. pinnulifer diversicolor on, in S. Rhodesia, 15; pests of, in Trinidad, 24; pests of, in U.S.A., 55, 256, 343, 377, 432-434, 460, 653; insects intercepted on fruit and seedlings of, in Japan and Russia, 50, 439, 616; insecticides and injury to, 6, 278, 653; effects of lead arsenate and copper carbonate on fruit of, 330, 331.

Orange, Mandarin, Coccids on, in Russian Union, 148, 308; males of *Icerya purchasi* reared on, 6.

orbitale, Dolichocolon. orbitalis, Cheiloneurus.

Orchestes (see Rhynchaenus).

Orchids, Coccid on imported, in Japan, 617.

Oregma lanigera, resistance of varieties of sugar-cane to, in Formosa, 444.

Oregon, pests of forests and timber in, 498, 499, 599; miscellaneous pests in, 596, 599; orchard pests in, 377, 498, 643; Bruchus pisorum on peas in, 325; potato pests in, 83, 84; list of strawberry pests in, 498; Aphids and virus diseases in, 83, 84, 369; beneficial insects in, 499, 599, 643; liberation of parasite of Forficula auricularia in, 379.

Oreta carnea, bionomics and control of, on Uncaria gambir in Sumatra, 385, 386.

Orgilus obscurator, establishment of, against Rhyacionia buoliana in Connecticut, 74, 177.

Orgyia antiqua, not parasitised by Trichogramma evanescens, 35; tests of ovicides on, 332.

Oria musculosa, causing white ear in wheat in Russian Union, 680.

orichalcea, Plusia.

oriens, Prociphilus.
Oriental Peach Moth (see Cydia molesta).

orientalis, Anomala; Aonidiella (Aspidiotus); Blatta; Bruchocida; Dysdercus.

Orius, predacious on Erythmelus psallidis in U.S.A., 538.

Orius insidiosus, predacious on Paratetranychus pilosus in Connecticut, 429.

Orius niger, in wheat fields in Central Asia, 453.

Orius tristicolor, predacious on

Macrosiphum onobrychis in Utah, 30.

Ormenis marginata (see Petrusina). Ornamental Plants, book on control of diseases and pests of, 437.

ornatum, Trogoderma.

ornatus, Myzus; Phytodietus. ornigis, Xenosternum.

ornigis, Xenosternum. ornithogalli, Prodenia.

Ornithopus spp., pests of, in Portugal and Spain, 495.

Orthezia praelonga, measures against, on Glyricidia and Citrus

in W. Indies, 25.

Orthodichlorobenzene, in mixture for impregnating wood against Coleoptera, 182; soil treatment with, against termites, 422, 694; emulsion of, against Tipulids, 95; and carbon bisulphide, impractical against Lepidiota frenchi, 88. Orthol K, 529.

Orthomorpha gracilis, on greenhouse tomatos in Britain, 333.

Orthorrhinus patruelis, on caçao in New Guinea Territory, **544.**

Orthostethus infuscatus, in sugarcane fields in Florida, 656.

Oryctes, bionomics and control of, on oil palms in Dahomey, 347.

Oryctes agamemnon, in Dahomey,

347. Oryctes boas, in Dahomey, 347.

Oryctes monoceros, in Dahomey, 347. Oryctes rhinoceros, habits of, in Formosa, 363, 446; in Philippines, 451; Catascopus facialis introduced into Samoa against, 641; on coconut, 451, 641.

Oryza barthi, relation of locusts to,

in Fr. W. Africa, 570.

oryzae, Calandra (Sitophilus); Chlorops; Lema; Nilaparvata;

Pachydiplosis.

Oryzaephilus surinamensis, in oil palm kernels in Dahomey, 349; in flour mills in Russian Union, 685; in stored grain in U.S.A., 341.

oryzella, Oscinis.

Oscinella, on wheat in Czecho-

slovakia, 468.

Oscinella frit (on cereals), wind in relation to spread of, in Britain, 167; in Denmark, 117; bionomics and distribution of, in Russian Union, 454; forms of, in Switzerland, 270; causing white ear in wheat, 680.

Oscinella frit var. pusilla, status of,

270.

Oscinis frit (see Oscinella).

Oscinis oryzella, on rice in Japan, 225.

oscus, Platygaster.

ostreaeformis, Aspidiotus.

ostreata, Pseudoparlatoria.

ostrinella, Moodna.

Otiorrhynchus, list of species of, on vines in Crimea and Caucasus, 43. Otiorrhynchus asphaltinus, 43. Otiorrhynchus brauneri, 43.

Otiorrhynchus caucasicus, 43.

Otiorrhynchus impexus, 43.

Otiorrhynchus ovalipennis, 43. Otiorrhynchus rugosostriatus, 43.

Otiorrhynchus singularis, on rhodo-

dendron in Britain, 295. Otiorrhynchus sulcatus, on rhododendron in Britain, 295.

Otiorrhynchus turca, on vines in Russian Union, 43, 403.

oudemansi, Tenuipalpus. ovalipennis, Otiorrhynchus.

ovatus, Platycampus.

Oxya chinensis, measures against, in China, 477.

Oxya intricata, measures against, in China, 477.

Oxya velox (on rice, etc.), bionomics and control of, in Korea, 557; in Philippines, 451.

Oxycarenus luctuosus, on fruit trees

in Queensland, 603.

parasite Oxyglypta rugosa, Apiomyia bergenstammi, 5. oxymorus, Cryptus.

Oxypleurites maxwelli, sp. n., on olive in California, 288.

Oyster-shell Scale (see Lepidosaphes ulmi).

P.

Pachnaeus opalus, on beans Florida, 600.

Pachydiplosis oryzae, parasites of, on rice and grasses in India, 577. Pachymerus nucleorum, in seeds of

Cocos spp. in Brazil, 661. Pachynematus montanus, on spruce in Czechoslovakia, 469, 517, 518;

parasites of, 518.

Pachynematus scutellatus, Syrphid predacious on, on spruce in Germany, 334.

Pachyneuron aphidis, hyperparasite of Eriosoma lanigerum in Jugoslavia, 10.

Pachyprotasis variegata, on potato in Holland, 269.

Pachytylus (see Locusta).

(1875)

pacificus, Tetranychus.

padellus, Hyponomeuta.

padi, Anuraphis.

padi, auct., Priophorus (see P. pallipes); Siphonaphis Rhopalosiphum prunifoliae).

Pagiophloeus longiclavis, on mahogany in India, 612.

Pagria signata, on soy beans in Japan, **556**.

Pagyda salvalis, damaging teak seed in Burma, 693.

RodoliaPalaeococcus, cardinalis predacious on, on Casuarina in Porto Rico, 434.

Palaeococcus rosae, 434.

Paleacrita vernata, measures against, in forests in U.S.A., 28,

Pales maculata (see Nephrotoma).

Palestine, miscellaneous pests in, 127, 512; vine moths in, 191; Aphelinus mali imported into, from Egypt, 127; Icerya purchasi on Citrus imported into Cyprus from, 270.

Hylastes (Hylurgops); palliatus, Tanymecus

pallicornis, Rhynchaenus (Orchestes). pallida, Paratamiclea; Pristiphora (Lygaeonematus).

pallidifrons, Erythroneura.

pallidus, Tarsonemus.

pallipes, Priophorus. pallipes, F., Agonum (Carabus).

pallipes, auct., Agonoderus (see A . lecontei) .

Palm, Ivory Nut (see Metroxylon). Palm, Licuri (see Cocos coronata).

Palm, Licurioba (see Cocos vagans). Palm, Oil (see Elaeis guineënsis).

Palm Borer, Giant (see Dinapate wrighti).

palmae, Asterolecanium. palmarum, Rhynchophorus.

Palms, pests of, in Brazil, 661; Dinapate wrighti on, in California and Mexico, 569; Promecotheca reichei on, in Fiji, 366; Oryctes rhinoceros on, in Formosa, 363; Brontispa limbata on, in Mauritius, 611; thrips imported into Japan on seedlings of, (See Coconut, Date Palm and Elaeis.)

Palomena prasina, parasite of, on cereals in Germany, 64, 632; effect of temperature and humidity on, 632.

Palomena viridissima, ecology of, attacking cereals in Germany, 631.

Palorus depressus, in oil-palm kernels in Dahomey, 349.

palpigera, Brachyacma.

Palpostoma aldrichi, sp. n., parasite of Lepidoderma albohirtum in Queensland, 294.

Palpostoma testacea, P. aldrichi

recorded as, 294. paludosa, Tipula.

Panana Canal Zone, termites in timber in, 180.

pancratii, Brithys.

Pandanus spp., Aeolarchis sphenotoma on, in Fiji, 213.

Pandemis (see Tortrix).

pandorus, Pholus satellitia.

panicea, Sitodrepa (Stegobium). Panicum spp., Nymphula depunctalis on, in Philippines, **364.**

Paniscus ocellatus, parasite of cutworms in Florida, 657.

Panolis flammea, on pines in Germany, 65, 354, 579; pupal weight and egg-production in, 65; parasite of, 354; forecasting outbreaks of, 65, 579.

Panorpa communis, predacious on Erannis spp. in Poland, 438.

Panthea coenobita, in forests in Germany, 466; characters distinguishing, from Lymantria monacha, 466.

Panthorhytes plutus, on cacao in New Guinea Territory, 277, 544; bionomics and control of, 277.

papaveris, Aphis.

Papaya, relation of *Dacus* spp. to, in Formosa, **206**; pests and disease of, in Porto Rico, **435**, **502**; pests of, in Queensland, **603**; injured by oil emulsion, **435**.

Papers, treatment of, against Ctenolepisma, 510; Psocids damaging, 102.

paphus, Protoparce sexta.

Papilio scamander grayi, on avocado in Brazil, 23.

Papilio xuthus, in forests in China, 360.

papua, Calotermes.

Papuana huebneri, measures against, on banana and taro in Gilbert Island, 366.

papuana, Promecotheca.

Parabatus cristatus, parasite of Monima cruda in Poland, 438.

Paradichlorobenzene, uses of, against borers, 256, 277, 420, 526, 527, 544; wood impregnated with mixture containing, against Coleoptera, 181; against clothes

moths and carpet beetles, 188, 635, 636; ineffective in spray against Cydia pomonella, 134; against Hylemyia brassicae, 317; against termites, 89, 544; as a soil fumigant, 88, 89, 366; fumigation of sweet potato tubers with, against Cylas, 379, 643; solutions and emulsions of, 134, 526, 527, 544; new method of volatilising, 643; and carbon bisulphide, 88; causing injury to plants, 88, 134, 256, 420.

paradoxa, Cedria.

Paraffin (Liquid), tests of penetration of, into tracheae of insects, 579, 580.

Paraffin Wax, as repellent for Chrysobothris femorata, 420.

Paraguay, Schistocerca paranensis in, 543.

Parahydrotaea jacobsoni, on tea and bamboo in Formosa, 339, 340.

Paralispa gularis (see Aphomia). parallela, Monotoma.

Paraluperodes nigrobilineata (see Monolepta).

paranensis, Schistocerca.

Paranthrene regalis, on vines in China, 450.

Paraphorantha peruviana, parasite of Dysdercus ruficollis in Peru, 164.

Paraphorantha politana, sp. n., parasite of Dysdercus mendesi in Brazil, 22.

Parapteromalus, parasite of spruce sawflies in Czechoslovakia, 518.

Pararcyptera microptera, ecology and migrations of, in Russia, 386.

Parasa latistriga, on plum in S. Rhodesia, 125.

Parasa sinica, measures against, on Platanus orientalis in China, 449.

parasae, Eurytoma (see E. monemae). Parasetigena segregata, auct. (see

Phorocera silvestris).

Parasitic Hymenoptera, factors concerned in host-finding by, 166, 302, 391; physiological causes of immunity from, 598; effect of host size on sex ratio of, 499; control of sex in, 595, 596; effect of cold storage on reproduction of, 144; agar preparation for feeding, 426.

parasitica, Rhodesina.

Parastasia inconstans, on cacao in New Guinea Territory, 544.

Parastasia marmorata, on cacao in New Guinea Territory, 544.

Paratamiclea pallida var. hova, parasite of Enaria in Madagascar, 611; introduced into Mauritius against Lachnosterna smithi, 611.

Paratetranychus bioculatus, on tea

in India, 128.

Paratetranychus citri(measures against on Citrus), in New Zealand, 56; in U.S.A., 256, 257, 261, 433, 459-461, 649, 650.

Paratetranychus indicus, life-cycle of, on Sorghum and sugar-cane

in India, 128.

Paratetranychus pilosus (on fruit trees), in New Zealand, 550; in Nova Scotia, 662; in Tasmania, 21; in U.S.A., 429, 432, 535; other food-plants of, 21, 429; sprays against, 21, 429, 432, 535, 536, 550; laboratory tests of ovicides against, 332, 535; predacious enemies of, and sprays destroying them, 429, 662.

Paratetranychus punicae, on pomegranate and vines in India, 128. Paratheresia, considered distinct from Theresia (q.v.), 143.

Paratrechina longicornis, in houses in Formosa, 446; baits for, in Porto Rico, 435.

Paratrioza cockerelli (on potato and tomato), in Alberta, 176; in U.S.A., 72, 79, 408, 422, 651; causing Psyllid yellows, 72, 176, 651; stimulating flowering and fruiting of tomato, 73; on other solanaceous plants, 422, 651; bionomics of, 408, 422; measures against, 72, 79, 651.

Paratrioza mexicana, in U.S.A., 422. pardalina, Locustana; Myiopardalis (Carpomyia).

Paregle radicum (see Hylemyia).

parietaricola, Phenacoccus.

Paris Green, in baits, 32, 67, 68, 127, 185, 287, 321, 323, 380, 390, 520; dusting with, 244, 245, 271, 277, 291, 370, 381, 417, 418, 463, 661; carriers for, 277, 291, 370, 661; spraying with, 137, 172, 376, 414, 438, 525, 583, 683; and calcium arsenate, 243, 244, 417, 418; and lime, 172, 438, 683; and sugar, 525; and sulphur, 244, 245, 381, 417, 463; against ants, 44; uses of, against termites, 56, 89, 181, 465; tests of toxicity of, to insects, 137, 243, 277, 528, 627; other insecticides compared with, 243, 244, 245, 417, 627.

Parlatoria oleae, bionomics of, in Uzbekistan, 474, 552, 553.

Parlatoria zizyphus, measures against, on Citrus and Ficus in Egypt, **509.**

Paroxyna misella, measures against, on chrysanthemum in Britain,

Parsley, new Aphid and disease of, in California, 590, 591.

Parsnip, Aphids on, in Tasmania and California, 21, 590.

partenopea, Encarsia.

parvipennis, Dasyscapus.

parvispinus, Isoneurothrips.

parvula, Epitrix; Erythroneura (Zygina).

parvulus, Pleurotropis.

Paspalum, Nymphula depunctalis

on, in Philippines, 364.

Paspalum dilatatum, planted round banana plantations Cosmopolites in New South Wales, 346.

Paspalum urvillei, insects on, in U.S.A., 248, 537, 657.

passerinii, Phloeomyzus.

Passiflora edulis, pests and disease' of, in Australia, 448.

passiflorae, Dacus (Chaetodacus). Passion Fruit (see Passiflora edulis). Patanga succincta, in India, 559;

on Citrus in Japan, 441. Patchiella réaumuri, in Italy, 512.

patrona, Casmara.

patruelis, Orthorrhinus.

pauperans, Termes.

Paururus (see Sirex). pauxillus, Rhynchites.

pawneepae, Aphis.

Pea, Heliothis armigera on, in S. Africa, 489; Cydia nigricana on, Britain and Canada, 38; Bruchus pisorum on, in Czechoslovakia, 468; Aphids on, in Germany, 686; Thrips angusticeps on, in Holland, 233; Etiella zinckenella on, in India, 129; pests of, in Japan, 361, 443, 556, 617; pests of, in Porto Rico, 501; pests of, in Russian Union, 454, 552, 679; pests of, in U.S.A., 29-31, 33, 323, 325, 370, 381, 415, 456, 497, 500, 531, 600; susceptibility of varieties of, to 33: Macrosiphum onobrychis, virus diseases in, 370, 671, 686; (stored), not attacked by Calandra granaria, 119; Plodia inter-punctella in, 173; X-rays for detecting Bruchus pisorum in, 147.

Pea, Pigeon (see Cajanus cajan). Pea Aphis (see Macrosiphum onobrychis).

Pea Bruchid (see Bruchus pisorum). Pea Moth (see Cydia nigricana). Pea Weevil (see Sitona lineata).

Peach, pests of, in Argentina, 188, 237, 602; pests of, in Australia, 20, 226, 449, 544, 639; pests of, in Baluchistan, 51, 577; Aulacaspis pentagona on, in Brazil, 86; Myzus persicae on, in Britain, 236; pests of, in China, 450, 668; M. persicae on, in Czechoslovakia, 468; pests of, in France, 195, 623; pests of, in Germany, 157, 267, 633, 687; M. persicae on, in Holland, 578: Aphids on, in Hungary, 458; pests of, in India, 503; Cydia spp. on, in Italy, 4, 197, 199, 235, 515; Dacus psidii on, in New Caledonia, 131, 132; Coccids on, in Russian Union, 316, 474, 552; Ceratitis capitata intercepted in fruit of, in Russian Union, 439; Cydia molesta on, in Switzerland, 438; pests of, in U.S.A., 34, 77, 255, 261, 288, 340, 377, 429, 430, 526, 527, 532, 623, 650; not attacked by Aegeria typhiaeformis, 630; susceptibility of, to Aspidiotus perniciosus, 690; Operophtera brumata unable to develop on, 513; insecticides and injury to, 255, 256, 272, 309, 425, 426, 458, 526, 527.

Peach Aphis (see Myzus persicae). Peach Borer (see Aegeria exitiosa). Peach Moth, Oriental (see Cydia

molesta).

Pear, Cydia pomonella on, in Algeria, 192; pests of, in Australia, 20, 21, 89, 292, 293, 390; pests of, in Baluchistan, 51, 577; pests of, in China, 450; pests of, in France, 168, 357, 622, 689; pests of, in Germany, 114, 155, 398, 632; pests of, in India, 503, 577; pests of, in Italy, 4, 115, 197, 235, 296, 297, 516; pests of, in Japan, 207, 225, 361, 557, 629, 630; gall-midges on, in Korea, 557; pests of, in Manchuria, Monosteira unicostata on, in Morocco, 359, 570; pests of, in New Zealand, 159; Psylla spp. on, in Rumania, 357: pests of, in Russian Union, 43, 316, 474, 552 : Ceresa bubalus on, in Switzerland, 169, 438; pests of, in U.S.A., 377, 383, 409, 410, 432, 434, 498, 535; resistance of varieties of, to Apiomyia bergenstammi, 5; susceptibility of, to Aspidiotus perniciosus, 690; types of injury to, by Monosteira unicostata and Eriophyes pyri, 359, 409; review of Cecidomyiids on, 557; effects of oil sprays on, 6, 94; problem of arsenic residues on fruit of, 168, 192, 193, 293.

Pear, Sand (see Pyrus sinensis). Pear Blossom Weevil (see Anthonomus piri).

Pear Leaf Blister Mite (see Eriophyes pyri). Pear Leaf Midge (see Dasyneura

pyri). Pear Psyllid (see Psylla pyricola).

Pear Sawfly (see Caliroa limacina). Pear Scale, Italian (see Epidiaspis leperii).

Pear Thrips (see Taeniothrips inconsequens).

Peat, for preparing fixed nicotine sprays, **545**.

Pecan, pests of, in U.S.A., 255, 372, 420; oil emulsions causing injury to, 255.

Pecan Nut Case-bearer (see Acrobasis caryae).

Pectinophora gossypiella (see Platy-

pectinophorae, Chelonus; Microbracon (Habrobracon).

pectoralis, Platycampus. pedatoria, Eupelmella. pedestris, Podisma.

Pediculoides ventricosus, predacious Ephestia kuehniella on Germany, 400.

Pediculopsis graminum, causing white ear of grasses and cereals in Russian Union, 151, 680; bionomics and control of, 151.

pelargonii, Macrosiphum (Aulacorthum).

Pelargonium inquinana, Coccid on, in Japan, 617.

pellicia, Cyrtorhinus.

pellio, Attagenus.

pellionella, Tinea. pellucida, Camnula.

pellucidator, Leptocryptus.

Pelobates fuscus, observations on, in Russia, 407.

pelochra, Draecenura.

Pempheres, parasites of, in India, 505, 506.

Pempheres affinis, parasites of, in India, 506; on cotton in Siam, 49.

pempheridis, Entedon. pempheriphila, Euderus,

Pemphigus, measures against, on Poinsettia in Oklahoma, 257; use of species of, as food for Hippodamia convergens, 81.

Pemphigus protospirae, in Italy, 512. penetrans, Macroglenes.

Penicillium, Necrobia rufipes feed-

ing on, 584.

Penicillium brevicompactum, experiments with, infesting Apion spp. in Sweden, 118.

penitalis, Pyrausta.

Pennsylvania, miscellaneous pests in, 79, 597, 599, 600; orchard pests in, 52, 535; Acantholyda erythrocephala on pines in, 651; beneficial insects and biological control in, 377, 599, 655.

pennsylvanicus, Camponotus herculeanus: Gryllulus (Gryllus)

assimilis:

pentagona, Aulacaspis (Pseudaula-

caspis).

Pentalonia nigronervosa (transmitting virus diseases of banana and Manila hemp), in Australia, 289; measures against, in Fiji, 213; in Haiti, 289; in Philippines, 265, 667.

Pentatoma rufipes, in Germany, 218, 631; ecology of, attacking cereals, 631; predacious on Lymantria monacha, 218.

Penthaleus major (haematopus), on peas in California, 600; on wheat

in Japan, 556.

Pentodon idiota, beet protecting Scorzonera against, in Russian Union, 679, 680.

Peppers (see Capsicum).

peregrina, Psylla.

Peregrinator biannulipes, predacious on Dinoderus minutus in Porto Rico, 423.

perforans, Xyleborus. pergandei, Phenacoccus.

Periclitena vigorsi, on Cordia myxa in Burma, 693.

Peridroma saucia, measures against, in Wisconsin, 32.

Perilitus coccinellae, in Formosa, 618; lists of hosts of, 618.

Perilitus epitricis, parasite of Phyllotreta in Louisiana, 373.

Perilla, Pyrausta phoenicealis on, in Japan, 210; Opatrum sabulosum, on, in Ukraine, 321.

Periplaneta americana, toxicity of Bacillus cereus to, 81.

Perisierola. parasite of Cydia molesta in Australia, 640; parasite of Platyedra gossypiella in Porto Rico, 503.

Perisierola nigrifemur, $oldsymbol{503.}$

Perkinsiella vastatrix, vector Fiji disease of sugar-cane in Philippines, 667.

perniciosi, Prospaltella.

perniciosus, Aspidiotus (Aonidiella, Comstockiella); Pseudococcus.

pernyi, Antheraea.

Peronea variana, on spruce in Quebec, 464.

Perrisia (see Dasyneura).

Persea gratissima (americana) (see Avocado).

perseae, Saissetia.

Persectania ewingi, measures against, on cereals and grasses in Victoria, 389.

persegnis, Cephalobus.

Persia, locusts in, 482.

persicae, Lecanium; Myzus (Myzodes, Phorodon).

persicae-niger, Anuraphis.

persimilis, Saissetia. Persimmon, Monema flavescens on, in Japan, 207; relation of Dacus spp. to, in Formosa, 206; relation of Ceratitis capitata on, in France, 195; Carpophilus hemipterus in rotting fruits of, 167.

persona, Xyleutes.

personata, Euxoa detersa.

personatus, Chrysomphalus (Aspidiotus).

persuasoria, Rhyssa.

persulcatus, Biosteres (Opius).

Peru, Coccids on Citrus in, 165, 511; Lymantriid on coca in, 165; cotton pests in, 164, 385; beneficial insects in, 164; parasites introduced into U.S.A. from, against Diatraea, 658.

peruella, Mescinia.

peruviana, Acaulona; Paraphorantha.

peruvianus, Dermestes. petiolaris, Mesochorus.

Petrol, uses of, against insects, 128, 233, 295, 577; in mixture for impregnating wood against Coleoptera, 182.

Petrusina marginata, fungus infesting, on Jasminum pubescens in

Porto Rico, 501.

Petunia, not attacked experimentally by Phthorimaea operculella, 289.

Peucetia viridans, predacious on insects in U.S.A., 251, 500.

Pexopsis pyrrhaspis, parasite of Hypopholis in S. Africa, 611; introduced into Mauritius against Lachnosterna smithi, 611. peyerimhoffi, Dioryctria.

Pezomachus (see Gelis).

Phaedon armoraciae, on horse-radish in Germany, 110.

Phaedon brassicae, measures against, in China, 450.

Phaedon cochleariae (on crucifers), in Germany, 110; in Russian Union, 439.

Phaenocarpa leveri, sp. n., parasite of Dacus in Fiji, 213, 605.

Phaeogenes nigridens, introduced into U.S.A. against Pyrausta nubilalis, 381.

phaeorrhoea, Nygmia. phaeosoma, Arctia caja.

Phalacrus fimetarius (coruscus), on Taraxacum kok-saghyz in Kazakstan, 45.

Phalaenoides glycinae, on vines in Australia, 294, 365; parasite of, 294.

Phalera flavescens, bionomics of, in Formosa, 206; on cherry in Japan, 211.

Phalonia posterana, on Taraxacum kok-saghyz in Kazakstan, 45.

Phanerotoma planifrons, introduced into U.S.A. and Porto Rico against Etiella zinckenella, 381, 382.

phantoma, Sirex.

Phanurus (see Telenomus).

pharaonis, Monomorium.

phaseoli, Agromyza; Bruchus (Callosobruchus); Trifidaphis.

Phaseolus lunatus (see Lima Beans). Phaseolus radiatus (with var. aureus), Bruchids reared on, in Japan, 361, 557, 617.

Phaseolus vulgaris (see Beans). Phassus, on Altingia excelsa in Java, 227.

Phassus malabaricus, measures against, on teak in India, 612.

Phaulacridium vittatum, in New South Wales, 130.

Pheasants, protection of Formica rufa against, 564.

Pheidole, attacking Diparopsis castanea in S. Africa, 489.

Pheidole javana, in houses in Formosa, 446.

Pheidole punctulata, associated with Pseudococcus kenyae in Kenya, 665.

Pheletes agonus (see P. ectypus). Pheletes californicus, in U.S.A., 410,

418; damage to potatoes by, 418; trapping adults of, 410.

Pheletes canus, in U.S.A., 28, 418; damage to potatoes by, 418; reactions of, to arsenicals, 28.

Pheletes ectypus, measures against, on tobacco in Connecticut, 177.

Phenacaspis prunorum, sp. n., on plum in Caucasus, **680**.

phenacocci, Leptomastix.

Phenacoccus aceris (on fruit trees), measures against, in Nova Scotia, 662; in Uzbekistan, 474.

Phenacoccus agropyri, sp. n., on Agropyrum in Caucasus, 680.

Phenacoccus caucasicus, sp. n., in Caucasus, 680.

Phenacoccus gossypii, resistance of, to parasitism by Coccophagus gurneyi, 598; parasitised by Leptomastix dactylopii, 598.

Phenacoccus hirsutus, new introduced parasite of, in Egypt, 506.

Phenacoccus parietaricola, sp. n., in France and Corsica, 48.

Phenacoccus pergandei, susceptibility of varieties of elm to, in Japan, 443.

Phenacoccus solani, parasites of, 598.

Phenothiazine (see Thiodiphenylamine).

Phenyl Phenol (Sulphonated), uses of, in dusts and sprays, 177, 179.

Phigalia sinuosaria, bionomics of, on Pyrus sinensis in China, 668; in Japan, 668.

Philaenus leucophthalmus, not transmitting potato viruses in Oregon, 84.

philanthus, Hoplia.

philippinensis, Spalangia.

Philippines, cotton pests in, 183, 184; Nymphula depunctalis on rice in, 364; locusts in, 451, 478, 485, 486; miscellaneous pests in, 451, 667; insects and virus diseases of plants in, 365, 667; beneficial insects and biological control in, 184, 364, 452; beneficial insects introduced into other countries from, 367, 665; Leucopholis irrorata intercepted in Hawaii in aircraft from, 368.

Phloeomyzus passerinii, in Italy, 512.

Phloeosinus armatus, bionomics of, on cypresses in Turkey, 219.

Phloeosinus bicolor (on conifers), in France, 225; in Turkey, 219; maternal galleries of, 225.

Phloeosinus cedri, on cedar in

Algeria, 226.

Phloeosinus cristatus, new parasite of, on Cupressus macrocarpa in U.S.A., 324.

Phloeosinus prostratus, on Juniperus in Algeria, 226.

Phloeosinus thujae, on Juniperus in France, 226.

Phloeosinus turkestanicus, bionomics of, on junipers in Central Asia, 153.

Phloeothrips kinugasai, sp. n., intercepted in Japan from Japanese Mandated Islands, 50, 616.

Phloeotribus liminaris, measures against, on peach in Illinois, 527.

Phloeotribus scarabaeoides, on olive in Italy, 473, 568; bionomics and control of, 473.

phloxiphaga, Heliothis.

Phlugiola dahlemica, sp. n., bionomics of, in hot-houses in Germany, 109.

Phlugiola redtenbacheri, P. dahlemica recorded as, 110.

phoenicealis, Pyrausta.

phoenicis, Mackiella; Rhynchophorus; Tenuipalpus.

Phoenix, new mite on, in greenhouse in Holland, 301.

Phoenix dactylifera (see Date Palm). Pholus spp., parasitised by Apanteles congregatus in U.S.A., 253.

Phomopsis, insects associated with, on pistachio in Greece, 452, 453.

Phonoctonus lutescens, predacious on Dysdercus in Sudan, 574.

Phoranthella mendesi (see Hyalo-mvia).

Phorbia, doubtful identity of, 114. (See Hylemyia.)

Phorocera agilis, considered correct name for P. silvestris (q. v.), 581.

Phorocera silvestris, parasite of Lymantria monacha in Germany, 335, 580, 581; bionomics of, 580, 581.

Phorodon humuli, on plum and hops in Britain, 332, 333; tests of ovicides against, 332.

Phorodon persicae (see Myzus).

Phosphides, danger of using, is

baits, 102.
Phostria piasusalis, on Coleus parvi-

flora in Madras, 129.

Photoptera, parasite of Setora nitens in Java, 347.

Phryxe vulgaris (see Zenillia).

Phthia, vector of Nematospora, 389. Phthia picta, on tomato in Jamaica, 239.

phthisicus, Mischocyttarus.

Phthorimaea, not considered distinct from Gnorimoschema, 289.

Phthorimaea gudmannella, new parasite of, in Virgin Islands, 133.

Phthorimaea heliopa (on tobacco), in S. Africa, 360; measures against, in Mysore, 625; in S. Rhodesia, 125.

Phthorimaea lycopersicella, on tomato in California, 133, 379, 432; new parasite of, 133;

measures against, 379.

Phthorimaea operculella, in S. Africa, 360; bionomics of, in Brazil, 289; development of, in imported potato tubers in Germany, 583; establishment of Apanteles scutellaris against, in Hawaii, 367; in S. Rhodesia, 125; factors affecting incidence of, in Victoria, 90, 389; on potato, 90, 289, 367, 389; on tobacco, 125, 289, 360; on other solanaceous plants, 289, 389; cultural measures against, 90, 389.

Phthorophloeus liminaris (see Phloeotribus).

Phycitinae, cactus-feeding species of, in America, 511.

Phygadeuon latipetiolator, parasite of Rhogas dendrolimi in Japan, 211.

Phylacter calcarator (see Zele).

Phyllaphis fagi, measures against, on beech in Germany, 353.

Phyllobius oblongus, on apple in Britain, 333.

Phyllobius pyri, on apple in Britain, 333.

Phyllocoptes, measures against, on vines in Queensland, 365.

Phyllocoptes cornutus, on peach in California, 33.

Phyllocoptes lycopersici, measures against, on tomato in Queensland, 187.

Phyllocoptes oleivorus (see Phyllocoptruta).

Phyllocoptinae, key to genera of, 34.

Phyllocoptruta oleivorus, measures against, on Citrus in Florida, 256, 257, 259.

Phyllopertha (see Anomala).

Phyllophaga (see Lachnosterna).

Phyllotoma nemorata, in Maine, 275; outbreak of, on birch in Quebec, 275; number of instars of, 275;

weight of foliage in relation to, 143; nicotine sulphate against,

275.

Phyllotreta, on beet in Britain, 674; on Taraxacum and crucifers in Russian Union, 45, 153, 439; methods of counting hibernating, in soil, 153.

Phyllotreta armoraciae, on horseradish in Germany, 110.

Phyllotreta atra, on horse-radish in

Germany, 110.

Phyllotreta cruciferae, on horseradish in Germany, 110; on Taraxacum kok-saghyz in Kazakstan, 45.

Phyllotreta nemorum, on horse-radish in Germany, 110.

Phyllotreta undulata, on horse-

radish in Germany, 110.

Phyllotreta vittata, on crucifers in Japan, 624; measures against, in China, 450.

Phyllotreta vittata discedens, bionomics and control of, on crucifers in Louisiana, 373.

Phyllotreta vittula, on wheat in Central Asia, 453.

Phylloxera, 392.

Phylloxera vitifoliae (vastatrix) (on vines), in Brazil, 34; in Germany, 688; in Poland, 403; in Queensland, 365; in Russian Union, 403; factors affecting injury by, in Switzerland, 46; legislative and other measures against, in Tunisia, 10, 11; forms or races of, 46, 403, 688; condition of European vines not affecting infestation by, 112.

Phymateus viridipes, in Belgian

Congo, 476.

Physalis francheti, Heliothis assulta transmitting virus disease in, in Japan, 444.

Physalis mollis, Paratrioza cockerelli on, in Texas, 651.

Physalis peruviana, Epilachna vigintioctopunctata on, in Fiji, 641. Physiology (of insects), text-book

on, **436.**Physokermes, new Lecanium recorded as, on rose in Bulgaria,

402.

Phytalus (see Lachnosterna).

Phytodietus ornatus, parasite of Tortricids in Transcaucasia, 587.

Phytoecia coerulea (rufimana), bionomics and control of, on crucifers in Azerbaijan, 320.

phytolymus, Maleuterpes (see M. spinipes).

Phytometra (see Plusia).

Phytomyza affinis (nigricornis), P. atricornis recorded as, on vegetables in Japan, 207, 443, 444; synonymy of, 443, 444.

Phytomyza aquifolii (see P. ilicis).
Phytomyza atricornis, on Cynara scolymus in California, 600; bionomics of, recorded as P. affinis (nigricornis) on vegetables in Japan, 207, 443, 444.

Phytomyza ilicicola (on Ilex spp.), confined to N. America, 669.

Phytomyza ilicis (on Ilex aquifolium), in Br. Columbia, 669; parasites of, in Br. Isles, 669; synonymy of, 669.

Phytomyza nigra, on cereals and

grasses in Japan, 208.

Phytomyza nigricornis (see P. affinis).

Phytomyza rufipes (ruficornis), on horse-radish in Germany, 110.

Phytonomus (see Hypera). piasusalis, Phostria.

Picea (see Spruce).

Picea abies, pests of: in Canada, 464, 647; in Czechoslovakia, 469; in U.S.A., 529, 540, 647.

Picea canadensis, Chermes ishiharai on, in Japan, 91.

Picea excelsa (see P. abies).

Picea glauca, pests of: in Canada, 368, 464, 647; in U.S.A., 540, 647.

Picea glehni, pests of, in Japan, 91, 211, 339.

Picea jezoensis, pests of, in Japan and Sakhalin, 19, 91, 211, 339, 615.

Picea mariana, Diprion polytomum on, in Canada and U.S.A., 368, 647; not attacked by Pissodes strobi in Quebec, 464.

Picea pungens, Diprion polytomum on, in Canada and U.S.A., 647; insecticides causing injury to,529.

Picea rubra, Diprion polytomum on, in Canada, 647; not attacked by Pissodes strobi in Quebec, 464; pests of, in U.S.A., 369, 540, 647; Pikonema alaskensis reared on, 464.

Picea sitchensis, Diprion polytomum on, in Canada and U.S.A., 647; Chermes spp. on, in Japan and Sakhalin, 91.

piceae, Chermes (Adelges). piceaella, Recurvaria. piceaperda, Dendroctonus.

piceus, Attagenus.

Picromerus bidens, predacious on Lymantria monacha in Germany, 218.

picta, Phthia; Scyllina; Zenillia (Exorista).

pictipes, Aegeria (Synanthedon). pictus, Poecilocerus.

Pieris, on crucifers in Russian Union. 439.

Pieris brassicae (on crucifers), in British Isles, 608; in Canada, 459; in Germany, 110, 232, 354; in Russian Union, 109, 306, 307, **310, 455, 589**; migrations of, in Europe, 608; parasitised by Apanteles plutellae, 505; relation races, of Trichogramma evanescens to, 109, 589; Troilus luridus attacking, 354; experiments with fungi and bacteria against, 306, 307, 455; attempted infection of, with Nosema cacto-blastis, 459; influence of crowding on larvae of, 232; tests of insecticides against, 250, 310.

Pieris napi, on horse-radish in

Germany, 110.

Pieris rapae (on crucifers), legislation against introduction of, into Australia, 344; in China, 450; in Germany, 110; sometimes migrating to tobacco in Japan, 210; in Manchuria, 361; in Russia, 109, 307, 455; in U.S.A., 249, 250, 370; parasites of, 109, 505; experiments with bacteria against, 307, 455; measures against, 249, 370, 450.

Piesma, characters of Polish species

of, 352.

Piesma capitatum, forms of, not transmitting crinkle disease of beet in Poland, 163, 352; on Atriplex, 352.

Piesma maculatum (and var. viride), not transmitting crinkle disease of beet in Poland, 163, 352; on

Atriplex, 352.

Piesma quadratum (on beet), in Germany, 688; in Poland, 105, 163, 350, 351, 352; other foodplants of, 163, 351, 352; types of crinkle disease transmitted by, 163, 351, 688; bionomics and migrations of, 163, 351; experiments with Beauveria against, 105; trap-strips of beet against, 163, 351, 352; conferences on status and control of, 352.

Piesma quadratum var. dilatatum, 352.

Piestopleura thomsoni, parasite of Macroglenes penetrans in Holland,

Piezotrachelus (see Apion).

Pigeon Peas (see Cajanus cajan). Pigeons, grain mites spread by, 552. piger, Cleonus; Microbracon.

pigra, Pygaera.

Pikonema alaskensis (on spruce), in Canada, 464, 540; bionomics and control of, in Maine, 540.

Pikonema dimmocki, on spruce in Quebec, **464**.

pilleriana, Sparganothis (Oenophthira).

pilosus, Paratetranychus.

Pimelephila ghesquierei, bionomics and control of, on oil palms in Dahomey, 348.

pimpinellae, Anthrenus.

Pimpla, parasite of Cydia nigricana in Britain, 38; parasite of Pissodes notatus in Finland, 8; parasite of Pyrausta and Sesamia in Italy, 296; parasite of Neosyrista similis in Japan, 211.

Pimpla capulifera, parasite Lymantria monacha in Germany,

581.

Pimpla disparis, parasite of Dendrolimus spectabilis in Japan, 211.

Pimpla examinator, parasite of Lymantria monacha in Germany,

Pimpla instigator, parasite of Lymantria monacha in Germany, 581.

Pimpla maculator, parasite of Vanessa cardui in Transcaucasia,

Pimpla nucum, parasite of Aegeria typhiaeformis in Italy, 630; parasitised by Dibrachys cavus, 630.

Pimpla pluto, parasite of Dendrolimus spectabilis in Japan, 211.

Pimpla roborator, parasite Etiella zinckenella in Ibe in Iberian Peninsula, 495.

Pimpla ruficollis, olfactory responses of, to food-plant of host in

Britain, 391. Pimpla spectabilis, parasite of

Rhythmonotus takagii in Japan,

pinastri, Sphinx.

Pine (Pinus), Chermes boerneri mistaken for C. pini on, in Australia, 262, 546; Chermes spp. on, in Britain, 262; restrictions on importation of, into Britain, 356; pests of, in Canada, 53, 464; pests of, in China, 449,

451, 614: pests of, in Czechoslovakia, 469; Diprion pini on, in Danzig Territory, 105, 397; new Coccids on, in Dominican Republic, 392; *Pissodes* spp. on, in Finland, 8, 9; pests of, in Germany, 65, 105, 231, 354, 397, 400, 401, 521, 579; new Coccid on, in India, 224; Thaumetopoea pityocampa on, in Italy, 516; pests of, in Japan, 209, 211, 339, 441; Hylastes ater on, in New Zealand, 504; pests of, in Poland, 401, 670, 676; pests of, in Rumania, 355; pests of, in Sweden, 123; pests of, Turkey, 219; pests of, in U.S.A., 53, 324, 329, 392, 461, 530, 568, 599, 651; effect of feeding Lymantria monacha on, 399; symbiosis of Siricids with fungi in, 401; insecticides causing injury to, 530; (timber), Hylotrupes bajulus in, 204, 335; Micromalthus debilis in gold mines in, 60; tests of preservatives against termites in, 180.

Pine, Jack (see Pinus banksiana). Pine, Japanese Red (see Pinus densiflora).

Pine, Loblolly (see Pinus taeda). Pine, Mugho (see Pinus mugo). Pine, Pitch (see Pinus rigida).

Pine, Ponderosa (see Pinus pondevosa).

Pine, Red (see Pinus resinosa).

Pine, Scots (see Pinus sylvestris). Pine, White (see Pinus strobus).

Pine Beetle, Southern (see Dendroctonus frontalis).

Pine Geometrid (see Bupalus piniarius).

Pine Sawflies (see Acantholyda erythrocephala and Diprion pini). Pine Shoot Moth (see Rhyacionia

buoliana).

Pine Sphingid (see Sphinx pinastri).

Pine-tar Oil (see Oil, Pine-tar).

Pineapple, Anomala orientalis on, in Hawaii, 367; Coccids on, in Mauritius, 611; Pseudococcus brevipes on, in Porto Rico, 502. pineti, Brachonyx.

pinetum, Diprion (Neodiprion).

Pineus (see Chermes).

pini, Chermes (Pineus); Crypto-parlatorea; Dendrolimus; Diprion (Pteronus); Lestodiplosis; Lygaeonematus (see Pristiphora abietina); Pissodes.

piniarius, Bupalus.

pinifolias, Chionaspis. Myelophilus biniberda. phagus).

piniphilus, Pissodes.

Pinnaspis buxi, introduction of Coccinellids against, on coconut in Hawaii, 366, 367.

pinnulifer, Chrysomphalus.

Pinthaeus sanguinipes, feeding habits of, in Germany, 354.

Pinus (see Pine).

Pinus banksiana, Diprion lecontei on, in Michigan, 461.

Pinus caribaea, Diprion lecontei on, in Canada and U.S.A., 53.

densiflora, Acantholyda Pinus erythrocephala on, in U.S.A., 651. Pinus echinata, Diprion lecontei on,

in Canada and U.S.A., 53. Pinus halepensis, Aesiotes leucurus

on, in New South Wales, 602; Chrysobothris igniventris on, in Turkey, 219.

Pinus koraiensis, Saperda laterimaculata on, in Russian Far East,

Pinus longifolia, treatment of timber of, against Platypus biformis in India, 612.

massoniana, Dendrolimus Pinus punctatus on, in China, 449.

Pinus mercusi, pests of, in Netherlands Indies, 577, 578.

Pinus monophylla, Pityophthorus on, in U.S.A., 324.

Pinus mugo, Diprion sertifer on, in Czechoslovakia, 469: Acantholyda erythrocephala on, in U.S.A., 279, 651.

Pinus nigra, Chrysobothris igniventris on, in Turkey, 219; (and var. austriaca), Acantholyda erythrocephala on, in U.S.A., 279, 651.

Pinus pinaster, Chermes boerneri recorded as C. pini on, in Australia, 262, 546.

Pinus ponderosa, pests of, in U.S.A., 140, 377, 378, 599; type of injury to, by Pityophthorus spp., 140.

Pinus radiata, Chermes boerneri recorded as *C. pini* on, in Australia, **262**, **546**; *C. boerneri* restricted to, in U.S.A., **546**; Aesiotes leucurus on, in New South Wales, 602.

Pinus resinosa, pests of, in U.S.A., 73, 177, 279, 461, 529, 600, 651; insecticides causing injury to, 529.

Pinus rigida, Diprion lecontei on,

in Canada, **53**; pests of, in U.S.A., **53**, **378**.

Pinus strobus, Pissodes strobi on, in Quebec, 464; pests of, in U.S.A., 279, 461, 651; weight of foliage of, in relation to insects, 143.

Pinus sylvestris, olfactory responses of parasites of Rhyacionia buoliana to, in Britain, 391; pests of, in Czechoslovakia, 469; pests of, in U.S.A., 461, 529, 651; insecticides causing injury to, 529.

Pinus sylvestris var. hamata, Brachonyx pineti on, in Rumania, 355.

Pinus sylvestris var. turfosa, Anthonomus varians on, in Rumania, 355.

Pinus taeda, Dendroctonus frontalis on, in U.S.A., 377, 599.

Piophila casei (Cheese Skipper), Necrobia rufipes predacious on, 584; cage for rearing, 78.

pipiens, Culex.

Pipunculus, new species of, in U.S.A. 596.

Pipunculus confraternus var. melanis, n., parasite of Eutettix tenellus in Utah, 596.

Pipunculus trochanteratus var. tenellus, n., parasite of Eutettix tenellus in Utah, 596.

piri, Anthonomus.

piricola, Diaspis (see Epidiaspis leperii).

pirifoliae, Epitrimerus.

Piripiri (see Acaena).
pirivorella, Numonia (Nephopteryx).
pisi, Contarinia; Macrosiphum
(Acyrthosiphon, Illinoia) (see M.
onobrychis).

pisivorus, Kakothrips.

pisorius, Ichneumon.

pisorum, Bruchus.
Pissodes gyllenhali, bionomics of, on conifers in Finland, 8.

Pissodes harcyniae, natural enemies of, on spruce in Finland, 9.

Pissodes notatus, bionomics of, on conifers in Finland, 8.

Pissodes pini, parasite of, on pine in Finland, 8.

Pissodes piniphilus, on pine in Finland, 9.

Pissodes strobi, on conifers in Quebec, 464.

Pissodes validirostris, bionomics of, on conifers in Finland, 8.

Pistacia vera (Pistachio), pests of, in Greece, 452, 453; Megastigmus ballestrerii on, in Sicily, 452;

insects associated with *Phomopsis* on, **452**, **453**.

pistaciae, Eurytoma; Tinea. Pisum sativum (see Pea). Pitch Pine (see Pinus rigida).

Pith Helmets, damaged by Rhizopertha dominica in India, 612.

pityocampa, Thaumetopoea.

Pityophthorus, new parasite of, on Pinus monophylla in U.S.A., **324**; unusual injury to P. ponderosa by, **140**.

Pityophthorus confertus, on Pinus ponderosa in California, 140. Pityophthorus confinis, on Pinus

ponderosa in California, 140.

Plaesius javanus (predacious on Cosmopolites sordidus), introduced into Cook Islands, 212; in Fiji, 212, 238, 365; introduced into Formosa, 205; introduced into Jamaica, 212, 238; in Java, 205; introduced into Tahiti, 212, 365.

plagiata, Anaitis; Tiracola.

plagiatus, Cneorrhinus; Prosoestus (Derelomus).

planifrons, Phanerotoma.

Plant Diseases (see Bacteria, Fungi and Virus Diseases).

Plant Pest Legislation, in Australia, 130, 274, 344, 592; criticism of, in Australia, 184; against introduction of Anthonomus grandis into Barbados, 661; in Br. Isles, 355, 356; against fruit-flies in Hawaii, 344; in Jugoslavia, 592, 690; in Kenya, 274, 666; in Tunisia, 11, 344; list of Coccids subject to, in Russian Union, 58; in U.S.A., 176, 274, 275, 344, 591; summaries of, in other countries, 176, 274, 344, 592.

Plantago lanceolata, Anuraphis roseus on, in Connecticut, 428.

Plants, injuries to, caused by insect toxins, 624; relation between condition of, and insect attack, 507; papers on breeding and selection of, for resistance to insects, 285, 456; bibliography on insecticides derived from, 392.

planus, Smerinthus.

platani, Lithocolletis.

Platanus occidentalis, Lithocolletis platani on, in Turkey, 219.

Platanus orientalis, Parasa sinica on, in China, 449; Lithocolletis platani on, in Turkey, 219.

Platyacus ruralis, on cacao in New Guinea Territory, **544.**

Platycampus laricis, on larch in Idaho, 599.

Platycampus laricivorus, on larch in Idaho, 599.

Platycampus ovatus, bionomics of, on larch in Germany, 216, 217, 218.

Platycampus pectoralis, bionomics of, on larch in Germany, 216, 217, 218.

Platychirus scutatus (predacious on Aphids), bionomics of, in Britain, 628, 629; morphology and

anatomy of, 628.

Platyedra gossypiella (Pink Cotton Bollworm), in Brazil, 22, 86; in Belgian Congo, 572; in Cyprus, 270; in India, 575, 576; in Korea, 381; in Manchuria, 340; in Nyasaland, 126; in Philippines, 183; in Siam, 50; compulsory fumigation of cottonseed against, in Sicily, 174; in Sudan, 572; in Tanganyika, 492; in U.S.A., 53-55, 381; organisation of control of, in U.S.A., 242; in W. Indies, 221, 222, 223, 271, 503, 660; factors affecting penetration of bolls by, 223; survival of, in cotton residues, 223; alternative food-plants of, 572, 577, 661; resting stage in larvae of, 184, 223, 272, 572, 573; other bionomics of, 184, 572; parasites and biological control of, **86**, **184**, **381**, **503**, **575**; suitable laboratory host of *Micro*bracon greeni, 129; other measures against, 53-55, 174, 184, 223, 272, 381, 576, 661; X-rays for detecting, in cotton seeds, 147: effect of soil treatment with sodium selenate on infestation by, 221.

Platygaster (parasitising Cecidomyiids), in Holland, 269; in India, 577; in Italy, 296.

Platygaster oscus, bionomics of, in Italy, 5.

Platygaster tuberosula, parasite of wheat gall-midges in Holland, 269.

Platygenia barbata, bionomics of, on oil palm in Dahomey, 348.

Platygenia 'quadrimaculata, bionomics of, on oil palm in Dahomey, 348.

platyhypenae, Euplectrus.

Platylabus, parasite of Erannis in Crete, 172.

Platynychus formosanus (see Cardio-phorus).

Platyptilia acanthodactyla, new parasite of, in California, 133.

Platyptilia pusillidactyla, of no value in checking Lantana in India, 612. Platypus biformis, treatment of pine timber against, in India, 612. Platysystrophus (see Arrhenodes). Platytelenomus hylas, parasite of Sesamia cretica in Sudan, 574. plebeja, Atreides.

Plecoptera reflexa, parasites of, on Dalbergia sissoo in India, 612.

Plesiocoris rugicollis, tests of new winter washes against, on apple in Britain, 236, 237.

pleurostigma, Ceuthorrhynchus.

Pleurotropis, parasite of Galerucella nymphaeae in Italy, 3; parasite of Apanteles congregatus in Tennessee, 254.

Pleurotropis amyntas, parasite and hyperparasite of Phytomyza ilicis in Britain, 669.

Pleurotropis parvulus (parasite of Promecotheca spp.), in Java, 366; distribution and utilisation of, in Pacific Islands, 366, 641; methods of distinguishing sexes of, 366.

plicatrix, Dasyneura. plicatus, Rhytidoderes.

Plodia interpunctella, in Britain, 154; in Germany, 230; in Italy, 172, 198; in Japan, 360; in Korea, 225; in dried fruit and nuts, 154, 225, 230; in other stored products, 154, 172, 198, 225, 360; bionomics of, 173, 198, 225, 230; measures against, 154, 173, 225, 360, 361; summary of data on, 198.

Plum, Timocratica haywardi on, in Argentina, 188; pests of, in Australia, 21, 292; pests of, in Baluchistan, 51, 577; pests of, in Britain, 236, 332, 333, 396; pests of, in Cyprus, 270, 271; pests of, in Czechoslovakia, 468; Hoplocampa minuta on, Denmark, 117; pests of, in France, 622, 689; pests of, in Germany, 46, 114, 297, 398, 633; Aphids on, in Hungary, 458; pests of, in India, 503, 577; pests of, in Italy, 4, 97, 98; Monema flavescens on, in Japan, 207; Monosteira unicostata on, Morocco, 359; Parasa latistriga on, in S. Rhodesia, 125; Coccids on, in Russian Union, 316, 474, 552, 680; Cydia funebrana on, in Switzerland, 336; pests of, in

U.S.A., 288, 429; reduced importance of Lecanium corni on, in Central Europe, 690; spray programme for, 396; oil sprays causing injury to, 6, 430; quassia sprays harmless to, 98.

Plum Curculio (see Conotrachelus

nenuphar).

Plum Sawflies (see Hoplocampa flava and H. minuta).

Plum Tortricid (see Cydia fune-

Plusia brassicae, on cabbage in U.S.A., 250, 370, 379; measures against, 370, 379, 380.

Plusia californica, on tomato in California, 431: confused with Heliothis armigera, 431.

Plusia chalcites, on cabbage and tomato in Fiji, 641.

Plusia orichalcea, measures against, on flax in Bengal, 577.

Plusia signata, tests of insecticides against, on tobacco in Sumatra, 17.

Plutella cruciferarum (see

maculipennis).

Plutella maculipennis (on crucifers), in Britain, 262; in Fiji, 641; in India, 129, 575, 576; in Jamaica, 239; in Louisiana, 250, 370; in New South Wales, 545; in New Zealand, 262, 545; in S. Rhodesia, 125; in Russian Union, 307, 439; bionomics of, 262; Microbracon greeni reared on, 129; parasites and biological control of, 262, 505, 545, 575, 576; experiments with bacilli against, 307; dusting against, 370. plutellae, Apanteles.

pluto, Pimpla.

plutus, Pantorhytes.

Poa, Pediculopsis graminum on, in

Russian Union, 151.

Poa pratensis, relation of Pentatomid cereal pests to, in Germany, 631; Pyrausta nubilalis experimentally fed on, 229.

podana, Tortrix (Cacoecia).

Podisma mikado (see Miramella). Podisma pedestris, in Siberia, 320. Podisma sapporensis, in Japan, 618. Poecilocerus pictus, plants attacked by, in India, 577.

Poecilonota festiva, bionomics of, on cypresses in Turkey, 218, 219.

Poeciloscytus cognatus, on Taraxacum kok-saghyz in Kazakstan, 45. Poecilus (see Pterostichus).

Poinciana, pests of, in W. Indies, 181, 464.

Poinsettia, Pemphigus on, in greenhouse in Oklahoma, 257.

Poison Hemlock (see Conium macu-

latum).

Poland, Piesma and leaf-crinkle of beet in, 105, 163, 350-352; utilisation of Beauveria against Piesma in, 105; forest pests in, 232, 350, 398, 401, 438, 670, 676; orchard pests in, 162, 163, 350, 675, 694; pests of stored grain in, 162, 220, 350; miscellaneous pests in, 162, 350, 403; locusts in, 9, 403; Melolontha spp. in, 9, 676; beneficial insects in, 232, 438, 671, 677; utilisation of Aphelinus mali in, 162.

Polia legitima, on tobacco in Con-

necticut, 178.

Polistes, predacious on Monema flavescens in Japan, 207; predacious on Crocidolomia binotalis in Seychelles, 336.

Polistes crinitus var. americanus, predacious on Laphygma frugiperda in Porto Rico, 503.

Polistes major, predacious on Laphygma frugiperda in Porto Rico, 503.

Polistes metricus, predacious Protoparce sexta in Florida, 251.

politana, Eulia; Paraphorantha. politus, Chilocorus.

Pollard (see Bran).

Polyblastus stenocentrus, parasite of spruce sawflies in Czechoslovakia, 518.

Polyblastus tener, parasite of spruce sawflies in Czechoslovakia, 518.

Polycaon stouti, in furniture in Ú.S.A., **499.**

against Polychloropentane, mites, 422.

Polychrosis botrana (Vine Moth), in Cyprus, 270; in France, 155; in Germany, 102, 111, 458, 465, 496, 582, 689; in Jugoslavia, 494; in Palestine, 191; in Russian Union, 403; in Sicily, 516; susceptibility of varieties of vines to, 191; on plums, 270; bionomics of, 111, 155, 191, 466, 494, 496, 516; breeding of, for testing insecticides, 111; sprays against pupae of, 103, 689; dusts and sprays against larvae of, 191, 516; timing of sprays against, 155; baits for, 465, 494, 496, 582; banding against, 458; use of females for trapping males of, 466, 582.

Polychrosis viteana (Grape Berry Moth), measures against, in U.S.A., 377.

polygonata, Pulvinaria.

polygoni-yonai, Myzus.

Polygonum, new Aphid on, in Japan, 104; Tanymecus palliatus on, in Ukraine, 322; insecticides tested on, 425, 532.

Polygraphus gracilis, Nitidulid predacious on, on Picea glehni in Japan, 339.

Nitidulid Polygraphus proximus, predacious on, on Abies sachali-

nensis in Japan, 339.

Polyhedral Disease, affecting parasitism of Lymantria monacha by Phorocera silvestris in Germany, 582; in wheat gall-midges in Holland, 268.

Polyodaspis endogena, sp. n., parasite of Cvdia leucostoma in Java,

Polyphylla fullo, measures against, in forests in Germany, 521, 522; on vines in Russian Union, 403.

Polyphylla olivieri, on vines in Russian Union, 403.

polytomum, Diprion.

Pomegranate, pests of, in Baluchistan, 51, 577; pests of, in India, 128, 385.

pometaria, Alsophila.

pomi, Aphis.

pomonella, Cydia(Carpocapsa, Laspeyresia); Rhagoletis.

pomonellae, Stomatoceras (Anthrocephalus).

pomorum, Anthonomus.

Poncirus trifoliata, Papilio xuthus on, in China, 360.

ponderosae, Dendroctonus.

Pontania (see Nematus).

Popillia japonica (Japanese Beetle), risk of introduction of, into Bermuda, 85, 86; in U.S.A., 84, 104, 138, 139, 165, 180, 377, 532; legislation against, in U.S.A., 274, 591; on fruit trees and elm, 532, 533; damaging lawns, 533; growth of larvae of, in different media, 336; parasites and biological control of, 84, 165, 377, 533; influence of instars of, on Tiphia popilliavora, 84; intestinal protozoa of larvae of, 392; infection of, with milky disease, 377; other measures against, 104, 138, 139, 274, 532, 533, 591.

popilliavora, Tiphia.

Poplar (Populus), pests of, in Britain, 563, 627; Alsophila pometaria on, in Canada, 663; Ceresa bubalus on, in France, 689; pests of, in Germany, 398, 401; pests of, in Morocco, 196, 359: Aspidiotus slavonicus on, in Uzbekistan, 474; Lachnosterna implicita on, in Wisconsin,

Poplar, Black (see Populus nigra). Poppy, Ceuthorrhynchus maculaalba on, in Czechoslovakia, 468, 469, 470.

Poppy Weevil (see Ceuthorrhynchus macula-alba).

Popular Names, of Arthropods, lists of, 437, 512.

Population Studies, of insects, 27, 80, 84, 98, 105, 131, 138, 140, 153, 161, 175, 258, 320, 330, 387, 388, 393, 489, 492, 531, 537; theoretical basis of, 145-147.

populi, Smerinthus. populicaulis, Pemphigus. populitransversus, Pemphigus. populnea, Saperda. Populus (see Poplar).

Populus nigra, Lepidoptera on, in Morocco, 196.

Populus tremula, pests of, Germany, 400; dinitro-orthocresol causing injury to, 400.

Populus tremuloides, pests of, in Ū.S.A., **528**, **599**.

Porotermes, in Australia, 545. portentosus, Brachytrypes. Porthesia dispar (see Lymantria). Porthesia similis (see Arctornis

chrysorrhoea).

Porthetria (see Lymantria). Porto Rico, pests of bamboos in, 423, 501; maize pests in, 328, 501, 503; miscellaneous pests in, 434-436, 501-503; sugar-cane pests in, 382, 424, 435, 502, 505, 657; termites in, 464, 465; annual insect population records in, 141; beneficial insects and biological control in, 85, 382, 423, 424, 434, 435, 502, 503, 505, 657; introduction of beneficial insects into other countries from, 24, 658, 660, 670; Bufo marinus introduced into Florida from, 657; pests from, intercepted in

U.S.A., 462. Portugal, locusts in, 479; pests of lupins and Ornithopus in, 495. Portulaca grandiflora, Rhizoecus kondonis on, in Japan, 617.

posterana, Phalonia.

postfasciatus, Euscepes. postvittana, Tortrix.

Potassium Ammonium Seleno-Sulphide, inadvisable for use against mites on Citrus, 460.

Potassium Antimony Tartrate (see

Tartar Emetic).

Potassium Cyanide, spraying with, against locusts, 477; soil treatment with, against Naupactus leucoloma, 260.

Potassium Fluoride, tests of, in baits for grasshoppers, 134.

Potassium Fluosilicate, spraying with, against Criocerid on rice,

Potassium Metabisulphite, reducing fermentation in baits for Dacus,

Potassium Thiocarbonate, causing

injury to azalea, 605.

Potato, pests of, in Australia, 90, 186, 226, 389; Leptinotarsa decemlineata on, in Belgium, 106, 583; Phthorimaea operculella on, in Brazil, 289; pests of, in Britain, 96, 97, 122, 236, 332, 562, 563; pests of, in Canada, 67, 176; pests of, in Czechoslovakia, 469; L. decemlineata on, 333, 398, 512, 583, 633, 634, 636, 686, 687; pests of, in Hawaii, 367; pests of, in Holland, 62, 106, 157, 269, 563, 578, 583; Megymenum gracilicorne on, in Japan, 441; L. decemlineata on, Luxemburg, 106, Epilachna vigintioctomaculata on, in Manchuria and Korea, 340, 362; Myzus persicae on, in S. Rhodesia, 125; pests of, in Russian Union, 302, 303, 311, 679; L. decemlineata on, in Switzerland, 160, 350, 583; pests of, in U.S.A., 53, 70, 71, 83, 84, 134, 177, 342, 372, 378, 379, 408, 418, 421, 500, 563, 651; Ceresa bubalus reared on, 390; experiments with strains and hybrids of, in relation to L. decemlineata, 107, 158, 193, 357, 358; susceptibility of varieties of, to other pests, 303, 389; types of injury to, by Psyllids, 176, 636, 651; insects and virus diseases in, 83, 125, 157, 236, 267, 301, 302, 303, 333, 546, 563, 578, 633, 634, 636, 686, 687; sprays to

destroy, when infected with virus diseases, 333; calcium cyanamide injuring wet foliage of, 159; (tubers), restrictions on importa-tion of, into Britain, 355, 356; Coleopterous fauna in stores of, in Finland, 392; Phthorimaea operculella in imported, in Germany, 583; legislation dealing fumigation of, against with Popillia japonica in U.S.A., 274; as bait for wireworms, 656, 683.

Potato Aphis (see Macrosiphum

solanifolii).

Potato Beetle, Colorado (see Leptinotarsa decemlineata).

Potato Leafhopper (see Empoasca

fabae). Potato Moth (see Phthorimaea operculella).

praeceps, Scolytus. praecox, Lycophotia.

praelonga, Orthezia. praetiosa, Bryobia.

praetor, Rhogas.

Praon simulans, parasite of Macrosiphum onobrychis in U.S.A., 30. prasina, Palomena.

pratensis, Lygus.

Prays citri, on Citrus in Philippines, 452.

Prays oleellus (Olive Moth), baits for, in Italy, 98.

Prenolepis longicornis (see Paratrechina).

Prepodes similis var. amabilis, natural enemies of, on Citrus in Jamaica, 239.

Prepodes vittatus, natural enemies of, on Citrus in Jamaica, 239.

priapus, Microphanurus.

Prickly-Pear (see Opuntia).

prima, Heteroschema (see H. rugosopunctata).

Primula, Jassid on, in greenhouses in Britain, 115.

Priobium carpini (see Trypopitys). Prionus californicus, in bridge timber in U.S.A., 499.

Priophorus pallipes (padi, auct.), bionomics of, on cherry, etc., in

Germany, 114.

Pristiphora abietina, on spruce in Czechoslovakia, 469, 517; Syrphid experimentally predacious on, in Germany, 334; in Switzerland, 518; parasites of, 518.

Pristiphora erichsoni (on larch), bionomics of, in Germany, 216 in Michigan, 461; utilisation of Mesoleius tenthredinis against, in Quebec, 464; maps showing distribution of, in Europe and N. America, 216; birds and mice destroying, 217; measures against, 217, 461.

Pristiphora laricis (on larch), in Czechoslovakia, 469; in Germany, 7, 216, 217, 218; biono-

mics of, **217**.

Pristiphora pallida (stecki), characters and synonymy of, 518.

Pristiphora subarctica, status and characters of, 518.

Pristiphora wesmaeli, on larch in Germany, 7, 216, 217; bionomics and control of, 217.

Pristomerus, introduced into U.S.A. from Korea against Platyedra gossypiella, 381.

Pristomerus ocellatus, parasite of Cydia molesta in U.S.A., 377.

Pristomerus testaceicollis, of little value against Holcocera pulverea in India, 129.

Pristomerus vulnerator, parasite of Cydia pomonella in Holland, 579. Probits, for expressing toxicity of insectioides at 26, 202, 200

insecticides, etc., 66, 243, 309. processionea, Thaumetopoea.

Prochiloneurus, key to species of, 40.

Prochiloneurus clavatus, sp. n., parasite of Pseudococcus spp. in E. Africa and Natal, 40.

Prociphilus konoi, migrating to Picea from Lonicera in Japan, 211.

Prociphilus oriens, migrating to Abies from Fraxinus and Syringa in Japan, 211.

proclia, Aphytis.

Procris ampelophaga (see Theresimima).

Prodenia eridania (see Xylomyges).
Prodenia litura, on cabbage and tomato in Fiji, 641; on tobacco in Japan, 210; on cacao in New Guinea Territory, 544; on cotton in Philippines, 184; on tobacco in Sumatra, 17; derris dust ineffective against, 17, 184.

Prodenia ornithogalli, baits for, on lucerne in Illinois, 67; on peas in Porto Rico, 501.

profana, Mictis.

prolongata, Xiphydria.

Promargarodes sinensis, gen. et sp. n., in China, 48.

Promecotheca (on coconut), distribution of Plurotropis parvulus against species of, in Pacific Islands, 366, 641; new parasites

of, in New Guinea Territory, 294, 506.

Promecotheca cumingi, on coconut in Philippines, **451**.

Promecotheca opacicollis, on coconut in New Hebrides, 366.

Promecotheca papuana, on coconut in New Britain, 366, 506; new parasite of, 506.

Promecotheca reichei, value of Pleurotropis parvulus against, on coconut and Metroxylon in Fiji, 366; in Tonga, 366.

promecothecae, Eurytoma.

Prontaspis citri, measures against, on Citrus in Argentina, 23.

pronuba, Agrotis. pronubana, Tortrix.

Prophanurus alecto (see Telenomus). Provhinotermes inopinatus, in damp wood in Fiji, 641.

Prorops nasuta, utilisation of, against Stephanoderes hampei in Brazil, 238; in Uganda, 238.

Prosoestus spp., bionomics of, on oil palm in Dahomey, 348.

prosopidis, Heterospilus. Prosopis spicigera, **505**.

Prospaltella, parasite of Aspidiotus chinensis in Japan, 617.

Prospattella aurantii, parasite of Lepidosaphes gloveri in Transcaucasia, 149.

Prospaltella berlesei, establishment of, against Aulacaspis pentagona in Brazil, 34, 87.

Prospaltella perniciosi, parasite of Aspidiotus perniciosus in Argentina, 24.

prostratus, Phloeosinus.

Protargionia larreae, bionomics and control of, in Argentina, 23, 24.

Proteose Protein, and sucrose, Dacus ferrugineus tryoni experimentally fed on, **345**.

Protoparce, measures against, on tobacco and tomato in U.S.A., 252, 380, 431, 432.

Protoparce quinquemaculata, on tobacco, etc., in U.S.A., 178, 252, 253, 431; bionomics of, 252; natural enemies and disease of, 178, 253; characters of, 252.

Protoparce sexta, on tobacco and tomato in U.S.A., 178, 251, 252, 253, 431; bionomics of, 251, 252; natural enemies and disease of, 178, 251, 253; characters of, 252.

Protoparce sexta paphus, parasite of, 253.

protoparcis, Sturmia. protospirae, Pemphigus.

protuberans, Dendrosoter. proximum, Trypodendron. proximus, Nematus (Pontania). pruinosum, Lecanium. prunastri, (Sphaero-Lecanium

lecanium).

Prune, Bryobia praetiosa on, in New South Wales, 57; Taeniothrips inconsequens on, in Oregon, 377, 498.

Prune Thrips (see Taeniothrips inconsequens).

pruni, Illiberis.

pruniella, Coleophora.

prunifoliae, Rhopalosiphum.

prunivora, Enarmonia.

prunorum, Aspidiotus; Diptilomiopus; Phenacaspis.

Prunus avium, pests of, in Germany, 156, 297.

Prunus emarginata, Hadrobregmus gibbicollis on, in Oregon, 499. Prunus mume, weevil on, in Japan,

Prunus spinosa, Ceresa bubalus on, in France, 689; Tibicen haematodes on, in Germany, 46; Coccids on, in Uzbekistan, 474.

psallidis, Erythmelus.

Psallus seriatus (Cotton Fleahopper), in U.S.A., 244, 245, 380, 538; overwintering on Croton, 245, 538: relation of wind to spread of, 245; Mymarid parasites of, 381, 538; dusting against, 244, 245, 380; methods of trapping, 245.

psammaula, Decadarchis.

psarae, Microgaster.

pentagona Pseudaulacaspis (see Aulacaspis).

Pseudiastata brasiliensis, predacious Pseudococcus brevipes Brazil, 23.

pseudobrassicae, Rhopalosiphum. pseudococcina, Hambletonia.

Pseudococcus, distribution and utilisation of parasites of, in Africa, 40, 503, 665; measures against, on apple in Br. Columbia, 66; ant associated with, on coconut in Mauritius, 611; classification of, 224.

Pseudococcus adonidum, possibly on vines in S. Australia, 164; in greenhouses in Japan, 616; on Citrus in New Zealand, 277; parasites of, 598; P. citriculus mistaken for, 616; measures against, 164, 278.

Pseudococcus boninsis, measures against, on sugar-cane in Florida,

Pseudococcus brevipes, in Brazil, 23; in Hawaii, 502; in Mauritius, 611; in Porto Rico, 502; on date palm and Cyperus, 23; on pineapple, 502, 611; natural enemies and biological control of, 23, 502.

Pseudococcus citri, in S. Africa, 40, 323, 503; in Egypt, 509; in Florida, 256, 259; in France, 266; altitude affecting primary food-plant of, in Java, 161; in Kenya, 666; in Russian Union, 403; on Citrus, 256, 259, 266, 323; on coffee, 161, 666; other food-plants of, 161, 259, 403, 509; natural enemies and biological control of, 40, 266, 323, 503, 598; ants hindering biological control of, 323; partial immunity from Coccophagus gurneyi in, 598; other measures against, 161, 509; tests of thiocyanate spray against,

Pseudococcus citriculus, on Citrus in Japan, 616.

Pseudococcus filamentosus, new parasite of, in Natal, 40.

Pseudococcus furcatispinus, sp. n., on graminaceous plants in Caucasus, 680.

Pseudococcus gahani, in Russian Union, 150, 589; on Citrus, 150; utilisation of Cryptolaemus montrouzieri against, 589; reactions of, to parasites, 598; fumigation against, 150.

Pseudococcus kazanskyi, sp. n., on Agropyrum in Caucasus, 680.

Pseudococcus kenyae, on coffee in Kenya, 39, 665; in Uganda, 503; natural enemies and biological control of, 39, 503, 665; differen-

ces between P. lilacinus and, 665. Pseudococcus lilacinus, natural of, introduced enemies Kenya from other countries against P. kenyae, 665; differences between P. kenyae and, 665.

Pseudococcus longispinus (see P. adonidum).

Pseudococcus maritimus, measures against, on Citrus in New Zealand, 277, 278; resistance to certain parasites in, 598.

Pseudococcus perniciosus, new parasites of, on coffee in Tanganyika,

Pseudococcus virgatus (see Ferrisiana).

J

pseudomagnoliarum, Coccus.

Pseudomicromelus australia, synonymy of, 434.

Pseudoparlatoria ostreata, measures against, on papaya in Porto Rico, 435.

Pseudophonus griseus (see Harpa-lus).

Pseudophonus pubescens (see Harpalus rufipes).

pseudosolani, Aulacorthum (Myzus) (see Macrosiphum solani).

pseudospretella, Borkhausenia. Pseudotsuga, restrictions on importation of, into Britain, **356.**

Pseudotsuga taxifolia (mucronata), pests of, in Germany, 230, 522; pests of, in U.S.A., 67, 324, 499, 599; insect fauna of, in northern Rocky Mountains, 67; insecticides causing injury to, 530; Coleoptera in timber of, 499.

pseudotsugae, Dendroctonus. psidii, Dacus (Chaetodacus).

Psidium, new Tineid on, in Argentina. 188.

Psidium guayava (see Guava).

Psila rosae (on carrot), in Czechoslovakia, 468; bionomics and control of, in Massachusetts, 496.

Psilocephala rufiventris, predacious on Horistonotus uhleri in S. Carolina, 500.

Psilura monacha (see Lymantria). Psocids, measures against, damaging papers and foodstuffs in Germany, 102; infesting house insulation material in U.S.A., 640.

Psylla costalis, bionomics and control of, on apple in Switzerland, 232; characters of, 232.

Psylla mali, on potato in Germany, 636; on apple in Nova Scotia, 662; on apple in Switzerland, 232; measures and tests of insecticides against, 232, 332, 662; characters of, 232.

Psylla peregrina, on potato in Germany, 636.

Psylla pyri, on pear in Rumania, 357.

Psylla pyricola (on pear), bionomics of, in Italy, 115; in New York, 535; in Rumania, 357; measures against, 115, 535.

Psylla pyrisuga, on pear in Rumania, 357.

Psyllia (see Psylla).

Psyllid Yellows, caused by Paratrioza cockerelli on potato and tomato, 72, 176, 651.

Psyllids, of Alberta, 175; trapped by leaves of tomato in Germany, 113.

Psylliodes angusticollis, on tobacco in Japan; 210.

Psylliodes attenuata, on hops in Britain, 333; measures against, on hemp in Ukraine, 322.

Psylliodes chrysocephala, measures against, on rape in Holland, 269.

Pterocarpus angolensis, Bostrychid in timber of, in S. Africa, 60.

Pterochlorus tropicalis (see Lachnus).
Pteromalus alboannulatus (see Dirhicnus).

Pteromalus gelechiae (see Dibrachys cavus).

Pteromalus klugi (nematicida) (see Tritneptis).

Pteronidea ribesii (see Nematus). Pteronus pini (see Diprion).

Pterostichus chalcites, predacious on Lachnosterna in Wisconsin, 423.

Pterostichus cupreus, measures against, on strawberry in Germany, 457.

Pterostichus vulgaris, measures against, on strawberry in Germany, 457.

Ptinus japonicus, bionomics of, in Japan, 445.

Ptinus tectus, damaging building timber in Germany, 7.

Ptychomyia remota, parasite of Levuana iridescens in Fiji, 212.

Ptychomyia selecta, parasite of Pristiphora wesmaeli in Germany, 217; parasite of Pikonema alaskensis in Maine, 540.

pubescens, Ophonus (Pseudophonus) (see Harpalus rufipes).

pudibunda, Dasychira. pudicus, Carpocoris.

puera, Hyblaea. pugnax, Solubea.

pulchricornis, Meteorus.

pullus, Anagyrus.

pulsatorium, Trogium. pulverea, Holcocera.

pulverulenta, Monima (see M. cruda). Pulvinaria, utilisation of Crypto-laemus montrouzieri against certain species of, in Russian Union, 454, 455, 589.

Pulvinaria aurantii, on Citrus in Transcaucasia, 150, 454; fumigation against, 150.

Pulvinaria bétulae, measures against, on vines in Germany, 103.

Pulvinaria floccifera, on tea in Abkhazia, 455.

Pulvinaria polygonata, bionomics of, on Citrus and Murraya in Formosa, 557.

Pumpkin, pests of, in Fiji, 641; Megymenum gracilicorne on, in Japan, 441.

punctatum, Anobium. punctatus, Dendrolimus.

punctiferalis, Dichocrocis.

punctiger, Ceuthorrhynchus; Isostasius.

punctillum, Stethorus.

punctiventris, Cleonus (Bothynoderes).

punctulata, Pheidole.

punctulatus, Cratosomus. punicae, Paratetranychus.

punicaeella, Euzophera.

Punjab, Bemisia gossypiperda on cotton in, 560; fruit pests in, 503; new Pyralid on sugar-cane in, 456.

pupivora, Trichospilus. purchasi, Icerya. purgatus, Enicospilus. purpurascens, Bothriophryne. purpureae, Nematus (Pontania).

pusilla, Oscinella frit. pusillidactyla, Platyptilia. pusillus, Corynetes.

putaminana, Cydia pomonella. putrescentiae, Tyrophagus.

Pycnarmon cribata, bionomics of, on Coleus parviflora in Madras, 129. Pygaera pigra, on poplar and willow in Morocco, 196.

pygmaea, Leptispa. pygmaeus, Cephus; Microbracon.

pyloalis, Margaronia. Pyrameis (see Vanessa).

pyrastri, Lasiophthicus (Catabomba, Scaeva).

Pyrausta aurata meridionalis, bionomics and control of, on mint in Morocco, 193.

Pyrausta insequalis, new parasite of, in California, 133.

Pyrausta nubilalis (European Corn Borer), in Britain, 116, 295; in Italy, 174, 198, 199, 295; in Japan, 556; in Manchuria, 340, 361; in Ontario, 25–27; in Philippines, 452; in Russian Union, 109, 146, 148, 152, 228-230, 587, 589; in U.S.A., 80, 138, 179, 180, 381, 414, 456, 645, 656; geographical distribu-tion of, 229; on Artemisia vulgaris, 116, 295; on hemp, 229; on range, 587; on sorghum, 198, 199, 295, 340; on wheat, 361, 556; on willow, 152; (1875)

bionomics and physiology of, 25-27, 80, 146, 148, 228, 230, 296, 327, 414; parasites and biological control of, 80, 109, 174, 296, 381, 589, 656; fumigation of broom corn under winter conditions against, in Canada, 416; other measures against. 138, 179 199, 296.

Pyrausta penitalis, adaptation of Chelonus annulipes to, 80.

Pyrausta phoenicealis, bionomics of, on Perilla in Japan, 210.

Pyrausta subsequalis (see P.

sequalis).

Pyrethrins, content of, in pyrethrum dusts and sprays, 249, 281, 380, 412, 417, 418, 537, 593, 620, 663; contents and evaluation of, in pyrethrum flowers, 134, 135, 392; effect of aqueous extraction on content of, in pyrethrum, 135.

Pyrethrum (including extracts), against Acridids, 618; against Aphids, 395, 450, 536; against Jassids, 53, 115, 281, 412, 501, 537: against other Rhynchota, 24, 417, 449, 463, 471, 543, 562, 667; against Coleoptera, 155, 156, 158, 188, 247, 249, 298, 342, 373, 380, 450, 623, 634; against Diptera, 297, 380, 497; against Lepidoptera, 28, 191, 154, 199, 207, 213, 310, 370, 376, 380, 400, 418, 431, 435, 449, 450, 521, 592, 614, 620, 663, 664, 689; against mites, 310, 536; against sawflies, 400; against thrips, 71, 525; uses of, against pests of stored products, 154, 188, 380, 521; tests of toxicity of, to insects, 144, 249, 277, 373, 620, 689; action of, on honey bees, 231, 402; effect of, as a stomach poison, 231, 310; inactivation of, after ingestion by Xylomyges, 524; dusting with, 24, 53, 155, 156, 158, 213, 247, 249, 281, 298, 370, 373, 376, 380, 400, 417, 418, 431, 449, 463, 471, 537, 543, 634, 662, 667; carriers for, 213, 249, 380, 449, 463, 537, 634; formulae for, in sprays, 310, 412, 663; preparation of extracts of, 135, 136, 310, 450; effect of aqueous extraction on, 135; effect of addition of chloropicrin to dusts of, 471; and derris, 155, 156, 158, 298, 471, 634; and lauryl thiocyanate, 413; and lead arsenate, 28, 536, 663; and limesulphur, 536; and nicotine,

449; and oil emulsions, 28, 376, 450, 689; atomised sprays of oils containing, 412; and soap, 71, 199, 207, 310, 501, 618, 620; and sodium carbonate, 449; and sulphur, 213, 247, 417, 418, 431, 463, 537; and tar distillates, 298, 689; and tobacco, 667; effect of storage on, 310; evaluation of, 134, 135, 136, 392; survey of data on, 104; other insecticides compared with, 155, 298, 380, 451, 536, 663.

Pyrethrum Plants, Thrips tabaci

on, in Kenya, 666.

pyri, Anthonomus (see A. piri); Dasyneura (Perrisia); Eriophyes; Phyllobius; Psylla; Stephanitis (Tingis).

pyricola, Psylla.

Pyridine, spray containing, against Contarinia nasturtii, 514.

Pyrilla, measures against, on sugarcane in India, 576.

pyrina, Zeuzera.
pyrisuga, Psylla.

pyrivora, Contarinia.
Pyrolene M.P., composition and use of, against whiteflies, 530;

and derris, 530.

Pyrophorus luminosus, introduced into Barbados and Mauritius from Porto Rico against Lachnosterna smithi, 660, 670; predacious on Prepodes spp. in Jamaica, 239; technique for transporting, 670.

Pyrosis idiota, on oaks in China,

360.

Collops.

pyrrhaspis, Pexopsis.
Pyrrhia umbra, on soy beans in
Manchuria, 340.

Pyrus sinensis (Sand Pear), Lepidoptera on, in China, etc., 667-669.

Q.

quadratum, Piesma.
quadratus, Stenocranophilus.
quadridentata, Ascogaster.
quadrigibbus, Tachypterellus.
quadriguttatus, Luperodes.
quadrilineatus, Opsimus.
quadrimaculata, Ceratia (Aulacophora); Platygenia (Clastocnemis);
Temnoschoita (see T. quadripustulata).
quadrimaculatus, Bruchus (Callosobruchus) (see B. maculatus);

quadripes, Xylotrechus.

quadripunctatus, Thanatophilus (Xylodrepa).

quadripustulata, Winthemia; Exochomus.

Quarantine, pests intercepted in: in Germany, 298; in aeroplanes in Hawaii, 368; in Japan, 50, 208, 568; in Russian Union, 439; in U.S.A., 462.

Quassia, spraying with, against Hoplocampa spp., 46, 97, 117, 197, 297; against Hyalopterus arundinis, 98; effect of, on Lepidoptera, 98, 197, 298; harmless to foliage and bees, 98; preparation of sprays of, 97, 298; keeping quality of, 46.

Quebec, forest pests in, 275, 276, 463; orchard pests in, 664; fumigation of broom corn against *Pyrausta nubilalis* in, 416; beneficial insects and biological

control in, 276, 464, 601.

Queensland, cotton pests in, 281, 602; Scirtothrips signipennis in, 329; sugar-cane pests in, 87, 88, 185, 186, 187; tomato pests in, 187, 281, 603; vine pests in, 89, 365, 602; miscellaneous pests in, 89, 186, 602; book on pests and their control in, 87; insects and virus diseases in, 87, 186, 448; beneficial insects and biological control of insects and plants in, 57, 185, 186, 263, 294, 345, 546, 602, 603.

Queensland Fruit Fly (see Dacus

ferrugineus tryoni).

Quercus (see Oak). Quercus ilex, new Coccid on, in France and Corsica, 235.

Quercus mongolica, Eutetrapha metallescens on, in Russian Far East, 42.

quercus, Kuwania.

Quince, Cydia pomonella on, in Algeria, 192; Typhlocyba froggatti on, in S. Australia, 292; Lepidoptera on, in Baluchistan, 51; Cantharis obscura destroying flowers of, in France, 622; Lepidoptera on, in Italy, 4, 629, 630; Monosteira unicostata on, in Morocco, 359; Coccids on, in Uzbekistan, 474.

quindecimpunctata, Anatis.

quinquefasciatus, auct. Culex (see C. fatigans).

quinquemaculata, Protoparce. quinquesignata, Hippodamia.

R.

radicum, Hylemyia (Paregle).

Radish, Phytoecia coerulea on, in Azerbaijan, 320: Athalia rosae on, in Denmark, 117; Hylemyia vadicum on, in Morocco, 605: (dried), Plodia interpunctella in, in Korea, 225.

raffrayi, Coptotermes.

Ragwort (see Senecio jacobaea). Ragwort Seed-fly (see Hylemyia

seneciella).

Rainfall, relations of locusts and grasshoppers to, 12, 13, 14, 92, 93, 94, 178, 214, 291, 388, 393, 394, 427, 481, 484, 485, 487, 545, **553, 557, 559, 602**; relations of other insects to, 16, 39, 85, 136, 182, 215, 216, 222, 226, 229, 238, 281, 327, 351, 386, 428, 435, 436, 447, 454, 457, 502, 550, 551, 588, 613, 664, 665, 677; effects of, on deposits and toxicity of sprays, 241, 249, 272.

Raisins, as food for Pleurotropis parvulus, 641. (See Fruit, Dried.) Rajputana, locusts in, 559, 560.

Ramie (see Boehmeria nivea). ramulorum, Myrmelachista ambigua.

rapae, Pieris (Ascia).

hirta Epicometis on, in Rape, Czechoslovakia, 468; Meligethes aeneus on, in Germany, 511; Psylliodes chrysocephala on, in Holland, 269; Athalia rosae on, in Siberia, 313. Rape Beetle (see Meligethes aeneus).

Rape Sawfly (see Athalia rosae).

Raspberry, Amphorophora rubi on, in Britain, 333; Carposina adreptella on, in New Zealand, 37; Carposina pests of, in U.S.A., 139, 178, 380; Parlatoria oleae on, in Uzbekistan, variety of, resistant to A. rubi, 280; Dasyneura plicatrix on, 673.

Raspberry Fruitworm (see Byturus

unicolor). réaumuri, Patchiella.

rectivostris, Anthonomus.

recurvalis, Hymenia. piceaella, measures Recurvaria against, on Picea abies in New

York, 529. Red Bud (see Cercis canadensis). Red-legged Earth Mite (see Halotydeus destructor).

Red-top Grass (see Agrostis vulgaris).

redtenbacheri, Phlugiola.

Redwood-bark Flour, as carrier for

dinitro-o-cyclohexylphenol dusts. 261, 650.

reflexa, Plecoptera.

regalis, Paranthrene (Sciapteron).

veichei. Promecotheca.

remota, Ptychomyia.

remotus, Hill, Coptotermes.

remotus, Silv., Coptotermes (see C. formosanus).

remulus, Trichacis.

renipustulatus, Chilocorus.

repandus, Calotermes.

Resin, and sodium carbonate in sprays, 278, 279, 560; in repellent paint against pests of books,

Resin Residue, as adhesive and spreader for sprays, 28, 241, 532.

Resin Soap (see Soap, Resin). Reticulitermes hesperus, in building

timber in U.S.A., 499. reuteri, Drepanothrips.

Revena nitida, infesting seeds of

Cocos spp. in Brazil, 661.

reversa, Sarcophaga.

Reviews:—Appel (O.) Ed. (in Sorauer, P.), Plant Protection,
290; Bondar (G.), Insect Pests and Parasites of Cacao in Bahia, 662; Borkhsenius (N. Coccidae of Quarantine Value for U.S.S.R. and their Allied Species, Burr (M.), The Insect Legion, 455; Buzin (N.), Printz (Ya.), Lazarevskii (M.), Negrul' (A.) & Katz (Ya.), Viticulture, 403; Couch (J. N.), The Genus Septobasidium, 242; Esaki (T.), Hori (H.) & Yasumatsu (K.), Insectorum Japonicorum illustratio iconographia coloribus ad naturam depicta, 446; Fishkis (I. S.), Mateev (V. A.) and Morozov (A. A.), Naphthalene and its Application in the Control of Granary Mites in Beet-growing Farms, 686; Frickhinger (H. W.), Elements of Pest Control, 606; Graham (S. A.), Principles of Forest Entomology, 2nd edn., 527; Hartnack (H.), 202 Common Household pests of North America, 382: Hopkins (A. D.), Bioclimatics, 325; Hunter (L.), Domestic Pests..., 237; Kemper (H.), The Pests of Stored Products... and their Control, 513; Metcalf (C. L.) & Flint (W. P.), Destructive and Useful Insects, 2nd edn., 655; Osborn (H.), Meadow and Pasture Insects, 527; Pape (H.), The Practice

of Control of Diseases and Pests of Ornamental Plants, 3rd edn., 437; Takahashi (T.) & Tsumagari (H.), Manual of Tobacco Insects of Japan, 210; Thompson (W. R.), Some Beneficial Insects, 4th edn., 436; Veitch (R.), Insect Pests and their Control, 87; Weber (H.), An Outline of Entomology, 170; Whelan (D. B.), Field Crop Entomology, 600: Wigglesworth (V. B.), The Principles of Insect Physiology, 436; Pests and Diseases of Sugar-beet, 439; A Handbook of Philippine Agriculture, 667.

Rhabdocnemis obscura (on sugarcane), in Hawaii, 90, 367; effect of trashing against, in Queensland, 185, 187; susceptibility of varieties of sugar-cane to, 186, 367; migrations of, 367; parasite of, 90, 367; tests of traps

for, 90.

Rhabdophaga saliciperda, bionomics and morphology of, on willow and poplar in Britain, 627, 628; measures against, 628.

Rhaconotus, parasite of Scirpophaga spp. in Mysore, 291; key to Indian species of, 506.

Rhaconotus cleanthes, sp. n., parasite of Pempheres affinis in India,

Rhaconotus menippus, sp. n., hosts of, in India, 506.

Rhaconotus menippus var. africanus,

n., in S. Africa, 506.

Rhagoletis cerasi (Cherry Fruit-fly), measures against, in Germany, intercepted in Russian Union, 439.

Rhagoletis pomonella (in apples), fumigation against, in Canada, intercepted in Russian Union, 439.

rhamni, Aphis (Doralis).

Rhamnus spp., Aphids on, Holland and Germany, 578, 686. Rhaphidopalpa similis (see Ceratia). vhinoceros, Oryctes.

Rhinocola aceris, on potato in Germany, 636.

Rhinotermes, in Australia, 545.

Rhizoecus kondonis, bionomics and control of, on Citrus, etc., in Japan, 617.

Rhizoglyphus echinopus, heat treatment of lily bulbs against, in Japan, 617.

Rhizoglyphus hyacinthi, measures

against, on Lilium longiflorum in U.S.A., 419; in Japan, 419.

Rhizopertha dominica, damaging pith helmets in India, 612; in 360: stored rice in Japan, measures against, 360, 612.

Rhizophagus spp., predacious on bark-beetles in Japan and

Sakhalin, 339.

Rhizophora, timber of, practically immune from termites in Porto Rico. 465.

Rhizopus, infesting chestnuts in France, 358.

Rhizotrogus solstitialis (see Amphimallus).

Rhodanates (see Thiocyanates). Rhode Island, work of Division of

Entomology in, 180.

Rhodesia, Northern, cotton pests in, 222, 492; locusts in, 483, 484.

Rhodesia, Southern, Citrus pests in, 14, 15, 16; cotton pests in, 222, 491, 492; pests of Dolichos lablab in, 124, 488, 489, 491; maize pests in, 124, 491, 492; miscellaneous pests in, 125; locusts and grasshoppers in, 124, 125, 637: insects transmitting virus diseases of tobacco in, 124, 125, 226; beneficial insects in, 16, 567.

rhodesiaensis, Bemisia.

Rhodesina parasitica, gen. et sp. n., parasite of Busseola fusca in S. Rhodesia, 567.

rhododendri, Giardomyia; Stephani-

Rhododendron, pests of, in Britain, 295; susceptibility of species of, to Arthrochlamys spilota in Japan, 212; new Cecidomyiid on, in New York, 499. (See Azalea.)

Rhododendron sinense, preparation of insecticide from, 450.

rhodophaga, Neocerata.

Rhogas, parasite of Malacosoma fragilis in Colorado, 528.

Rhogas dendrolimi (parasite Dendrolimus spectabilis), parasites of, in Japan, 211.

Rhogas praetor, parasite of Smerinthus planus in Japan, 624.

Rhogas testaceus, parasite of Lepidoptera in Poland, 438.

Rhopalosiphoninus latysiphon, on potato in Holland, 578.

Rhopalosiphoninus staphyleae (tulipaella), not transmitting disease of tulips in Oregon, 369.

Rhopalosiphum graminum (see

Toxoptera).

Rhopalosiphum maidis (see Aphis). Rhopalosiphum melliferum (see

Hyadaphis sii).

Rhopalosiphum prunifoliae, measures against, on apple in Britain, 333; causing white ear in wheat in Russian Union, 680; possible race of, on wheat in U.S.A., 542.

Rhopalosiphum pseudobrassicae, on crucifers in Louisiana, 254, 371; application of mathematics to study of, 254; measures against, 371.

Rhopalosiphum rufomaculatum, not experimentally transmitting lettuce mosaic in Britain, 671.

Rhopalosiphum splendens, on wheat in Kansas, 542.

Rhotonon, 592.

Rhubarb, Anthomyiid on, in Holland, 269.

Rhus javanica var. roxburghi, Aphids producing tannin galls on, in

Formosa, 205.

Rhyacionia buoliana (Pine Shoot Moth), olfactory responses of parasites of, to Pinus sylvestris in Britain, 391; parasites and biological control of, in Connecticut, 73, 74, 176, 177; in Czechoslovakia, 469; bionomics of, 73, 74; sprays against, 177.

Rhyacionia turionana, on Pinus sylvestris in Czechoslovakia, 469.

Rhyncaphytoptus ficifoliae, gen. et sp. n., on Ficus in California, 288. Rhynchaenus, synonymy of, 595.

Rhynchaenus pallicornis, nomenclature of, **595.**

Rhynchites aequatus (on fruit trees), in Britain, 333; bionomics and control of, in Germany, 156.

Rhynchites awatus, bionomics and control of, on fruit trees in Germany, 156.

Rhynchites auratus ferganensis, tests of insecticides on, in Uzbekistan, 627.

Rhynchites bacchus, bionomics and control of, on apple in Germany, 156.

Rhynchites betulae, bionomics and control of, on vines and fruit trees in Germany, 156.

Rhynchites bicolor, systematic position of, **595**.

Rhynchites coeruleus, bionomics and control of, on fruit trees, etc., in Germany, 155, 156.

Rhynchites cupreus, on fruit trees and vines in Germany, 155; bionomics and control of weevil

identified as, on *Prunus mume* in Japan, **556.**

Rhynchites heros, measures against, on pear in China, 450.

Rhynchites pauxillus, on fruit trees and vines in Germany, 155.

Rhynchocoris longirostris, on Citrus in Philippines, 452.

Rhynchophorus palmarum, bionomics of, on coconut in Brazil, 22.
Rhynchophorus phoenicis, bionomics

and control of, on oil palms in Dahomey, **347**, **348**.

Rhyncolus culinaris, investigations on, in building timber in Russia,

321.

Rhyparida spp., on cacao in New

Guinea Territory, 544.

Rhyssa persuasoria (parasite of Sirex), ecology of, in Britain, 504; establishment of, in New Zealand, 504; destroying Ibalia leucospoides, 504.

Rhythmonotus takagii, parasite of Dendrolimus spectabilis in Japan, 211; parasites of, 211.

Rhytidodera simulans, bionomics of, on mango in Netherlands Indies,

Rhytidoderes plicatus, bionomics and control of, on crucifers in France, 458, 459, 621; in Corsica and N. Africa, 459.

Ribes, species of Aphis infesting, 596. (See Currant and Goose-

berry.)

Ribes alpinum, new Aphid on, in Germany, 100.

Ribes gracile, Aphis ribis on, in Missouri, 597.

ribesii, Nematus (Pteronidea); Syrphus.

ribicola, Dasyneura; Macrosiphum. ribiensis, Aphis.

ribi-gillettei, Aphis.

ribis, Aphis; Capitophorus (Cryptomyzus); Eriophyes.

Rice, pests of, in China, 451, 613; Cirphis unipuncta on, in Fiji, 213,641; Spodoptera mauritia on, in Formosa, 363; Chilo simplex on, in Hawaii, 367; pests of, in

India, 291, 577; pests of, in Japan, 207, 209, 225, 362, 440, 441, 555, 556, 619, 620; pests of, in Korea, 557, 618; Hispids on, in Madagascar, 664; pests of, in Manchuria, 340; pests of, in Philippines, 451, 452; Aethemenes chloris on, in Sierra Leone, 271;

Solubea pugnax on, in U.S.A., 537; susceptibility of varieties

of. to Chlorops orvzae, 555; Nephotettix transmitting dwarf

disease of, 362.

Rice (Stored), pests of, and their control, 120, 225, 248, 339, 360, 378; nutritive value of forms of, for Ephestia and Calandra, 64; (milled), types of bags for protecting, from pests, 248.

Rice Borer (see Chilo simplex).

Rice Caseworm (see Nymphula depunctalis).

Rice Stem Maggot (see Chlorops

oryzae).

Rice Stink Bug (see Solubea pugnax). Rice Weevil (see Calandra oryzae). Rice Weevil, Dicky (see Maleuterpes spinipes).

Ricinus communis (see Castor).

rimac, Ipobracon.

rimosalis, Evergestis.

riparius, Hypnoidus (Cryptohyphus). ripperti, Eutermes (Nasutitermes). Robinia, pests of, in Russian Union,

152, 216, 474.

roborator, Pimpla. robusta, Cantheconidea; Hypsipyla. robustus, Kakothrips (see K. pisivorus).

Rock Phosphate, and arsenic, dustting with, against Laphygma exempta, 368.

Rocky Mountain Locust (see Melanoplus mexicanus).

Rodolia cardinalis (predacious on Icerya purchasi), in Argentina, 23; in Formosa, 446; utilisation of, in Japan, 445; introduced into Mauritius from S. Africa against I. seychellarum, 611; liberation of, in Porto Rico, 434; establishment of, in Russian Union, 589: attacking Palaeococcus, 434.

vodriguezensis, Campsomeris.

Romalea microptera, ecology of, damaging narcissus in U.S.A., 654.

rosa, Ceratitis.

rosaceana, Tortrix (Cacoecia).

rosae, Aphidius; Athalia; Macrosiphum; Palaeococcus (Crypti-cerya); Psila.

Rose (Rosa), new Coccid on, in Bulgaria, 402; Mictis profana on, in Fiji, 641; Cantharis obscura on, in France, 623; Aphids on, in Hungary, 458; Neosyrista similis on, in Japan, 210; Macrosiphum rosae on, in Tasmania, 21; pests of, in U.S.A., 141, 142, 429, 600; species of Macrosiphum overwintering on, 472, 563, 686; insecticides causing injury to, 272, 458.

Rose Midge (see Neocerata rhodo-

phaga).

Rose Sawfly (see Neosyrista similis). rosea, Chondacris; Monolepta. Rosette Disease (see Virus Diseases).

roseus, Anuraphis. rossi, Chrysomphalus.

rostrata, Aelia.

Rotenone, 593; dusting with, 32, 194, 250, 325, 376, 380, 461, 623; carriers for, 32, 250, 325, 623; spraying with, 32, 55, 71, 194, 380, 460, 592, 623; and oil emulsions, 55, 460; and soap, 71, 194; action of, on honey bees, 402; effect of ingestion of, by insects, 241; contents of, in derris (q. v.), 17, 30, 69, 70, 179, 184, 249, 250, 251, 269, 289, 294, 365, 370, 371, 373, 380, 402, 417, 426, 493, 497, 498, 523, 530, 537, 642; contents of, in cubé, 30, 71, 177, 179, 250, 286, 426, 498, 530; in Tephrosia, 250, 251, 502; methods of isolating and determining, 224, 294, 347, 392, 493, 502; review of literature on, 224; other insecticides compared with, 78, 250, 380.

Rottboellia exaltata, Aethemenes chloris not observed on, Sierra Leone, 271.

rotundatum, Gymnosoma.

rubea, Scolia.

rubeo, Cremastus.

rubi, Amphorophora; Anthonomus. rubiellum, Macrosiphum.

rubiginosa, Lachnosterna (Phyllophaga).

rubra, Kuwania.

rubrocinctus, Selenothrips.

Rubus, Dasyneura plicatrix confined to, 673; primary food-plant of Macrosiphum avenae, 472. (See Blackberry, Dewberry and Rasp-

Rubus australis, probably original food-plant of Carposina adrep-

tella in New Zealand, 37.

rufa, Formica.

Agrilus ; ruficollis, Dysdercus;

Necrobia; Pimpla.

ruficornis, Aphanistes; Harpalus (see H. rufipes); Phytomyza (see P. rufipes).

rufifemur, Eulimneria.

rufimana, Phytoecia (see P. coerulea).

rufimanus, Bruchus.

rufipes. Angitia: Harpalus; Hylastes (Hylurgopinus); Necrobia; Pentatoma; Phytomyza. rufiscutellaris, Glypta.

rufiventris, Psilocephala.

rufocuprea, Anomala.

rufomaculatum, Rhopalosiphum. ruforum, Achrysocharella

(Tetracampe).

rufus, Aptinothrips; Lophyrus (see Diprion sertifer).

rugiceps, Euetheola.

rugicollis, Plesiocoris.

rugosa, Lachnosterna (Phyllophaga); Oxyglypta.

rugosopunctata, Heteroschema. rugostriatus, Otiorrhynchus.

rugulosum, Lecanium. rugulosus, Scolytus.

Rum Puncheons, Xyleborus badius damaging, in transit to England from Br. Guiana, 328, 524.

Rumania, orchard pests in, 357, 690; pests of pines in, 355; list of Tachinids in, 608; summary of plant quarantines in, 592.

Rumex, Aphids on, in Britain, 673, 674.

rumicis, Aphis.

ruralis, Platyacus; Voria. Russia (U.S.S.R.), beet pests in, 306, 321, 322, 407, 439, 677, 678, 684; cereal pests in, 109, 145, 148, 152, 229, 311, 315, 322, 453, 454, 554, 587; Coccids on *Citrus* in, 148, 149, 150, 454; cotton pests in, 43, 147, 150, 215, 304, 313, 314, 552, 677, 680; 206, 307 crucifers in, 109, 152, 306, 307, 310, 316, 320, 439, 455, 589; pests of cucurbits in, 152, 310; Pediculopsis graminum on grasses in, 151; flea-beetles on hemp and flax in, 322, 684; pests of leguminous plants in, 108, 150, 152, 305, 322, 454, 474, 475, 568, 681, 683: Lymantria dispar on oak in, 456; orchard pests in, 43, 109, 303, 304, 305, 315, 316, 455, 588, 589, 680, 681, 683; pests of ramie in, 587; pests of rubberproducing plants in, 43, 44, 45, 679; pests of vines in, 42, 43, 403, 683; Pyrausta nubilalis on willow in, 152; pests of stored cereals and seeds in, 147, 304, 312, 318, 319, 440, 551, 585, 685, 686; Coleoptera in building timber in, 321, 322; new Coccids in, 680; book on Coccids subject to quarantines in, 58; key to

fruit-flies intercepted in, 439; locusts and grasshoppers 109, 146, 147, 160, 321, 386, 553; Opatrum spp. in, 41, 321; (Crimea), injurious Thysanoptera of, 43; factors affecting distribution and outbreaks of important pests in, 145-147, 228, 230, 454; insects and virus diseases of potato and beet in, 302, 303, 439; beneficial insects and biological control in, 109, 148, 149, 215, 303, 304, 305, 316, 319, 320, 453, 454, 455, 551, 555, 587, 588-590; experiments with bacteria and fungi against pests in, 306, 307, 308, 455, 684; difficulty of establishing Antheraea pernyi in, 403, 586; food of toads, etc., in, 407; review of literature on economic entomology in, 214. (See Siberia and Asia, Central.)

Russian Far East, Aspidiotus perniciosus in, 315; cereal pests in, 230, 454; Lamiids and Scolytids in, 42, 456; Coccinellid introduced into Caucasus from, against

A. perniciosus, 303. rustica, Cantharis; Dexia.

rutella, Setomorpha.

Rye, pests of, in Czechoslovakia, 468, 469; Oscinella frit on, in Denmark, 117; pests of, in Germany, 63, 352, 353, 519, 631; Blissus leucopterus on, Illinois, 284; Calliptamus italicus damaging, in Poland, 9; Chloropisca notata on, in Japan, 556; pests of, in Russian Union, 311, 312, 554, 680; Mayetiola destructor not developing on, 691; types of injury to, by Pentatomids, 352, 353; insects and white ear in, 680.

Rye (Stored), pests in, and their control, 64, 119, 173, 220, 305, 318, 319, 355; (bran), Attagenus pellio in, in Germany, 224.

Rye Beer, as bait for Theresimima,

Rye Grass (see Lolium).

S.

sabulosum, Opatrum. saccharalis, Diatraea. sacchari, Aphis. Saccharine, in baits for locusts, 478. saccharina, Lepisma; Tomaspis. Saccharosydne saccharivora, sugar-cane, etc., in Florida and Porto Rico, 657; susceptibility of varieties of sugar-cane to, 657; parasites and biological control of, 657.

sachalinensis, Cephaleia. saeuberlichi, Systates.

Sainfoin (see Onobrychis sativa).

Sainfoin Weevil (see Tanymecus palliatus).

St. Croix, new parasite of *Phthorimaea gudmannella* in, **133.**

St. John's Wort (see Hypericum perforatum).

St. Kitts, biological control of *Diatraea saccharalis* on sugarcane in, **659**, **660**.

St. Lucia, Leucoptera coffeella on coffee in, 435; biological control of Diatraea spp. on sugar-cane in, 182, 505, 659; D. lineolata erroneously recorded in, 182.

St. Vincent, miscellaneous pests in, 271, 272.

Saipan (see Japanese Mandated Islands).

Saissetia coffeae, bionomics and control of, on Citrus in New Zealand, 524.

Saissetia cuneiformis, new parasite of, in Eritrea, 503.

Saissetia nigra, food-plants of, in Japanese Empire, 445; on grape-vines in Queensland, 365.

Saissetia oleae (on Citrus), in California, 342, 377, 424, 433, 460, 649, 653; new parasites of, in Natal, 40, 503; bionomics of, in New Zealand, 524; on apricot, 649; proposed utilisation of Diversinervus against, 424; other measures against, 377, 433, 460, 524, 649; resistance of strains of, to HCN, 342; method for distinguishing dead, 653.

Saissetia perseae, new parasite of, in Natal, 503.

Saissetia persimilis, new parasites of, in S. Africa, 40.

saliciperda, Rhabdophaga.

saliciperdae, Eurytoma.

salicis, Stilpnotia.

salignae, Aphidius. salignus, Lachnus (Tuberolachnus).

Salt (Sodium Chloride), ineffective against Cantharis obscura, 623; in baits for cutworms, 390; as repellent for Melolontha, 299; and sand, use of, against Oryctes, 347; tea bushes injured by wash containing, 340; effects of, on removal of spray residues from apples, 189, 648, 649.

Sakhalin, Coccids of, **456**; forest pests in, **19**, **91**, **339**, **615**; beneficial insects in, **91**, **339**.

Salix (see Willow).

Salix alba var. coerulea, pests and diseases of, in Britain and Holland, 562, 563.

Salsola pestifer, relation of Eutettix tenellus and curly-top to, in Idaho, 286.

saltatrix, Meromyza.

Saltbush (see Atriplex).

salvalis, Pagyda.

Salvia sclarea, thrips on, in Crimea, 43.

Samoa, Dacus spp. in, 212, 365; Oryctes rhinoceros in, 641; beneficial insects introduced into, 212, 365, 641.

samsonovi, Eurytoma.

San José Scale (see Aspidiotus perniciosus).

sanborni, Aphis.

sanctus, Microdus (Bassus).

Sand Pear (see Pyrus sinensis).

Sand Wireworm (see Horistonotus uhleri).

Sandal, Dacus passiflorae on seeds of, in Fiji, 366.

Sandwich Method, for testing toxicity of insecticides, 241, 243, 524. sanguinipes, Pinthaeus.

Santo Domingo (see Dominican Republic).

Santobane A Emulsifier, 141.

Saperda spp., trees attacked by, in Russian Far East, 42, 43.

Sappocallis ulmicola (see Myzocallis).

sapporensis, Podisma.

Sarcocephalus cordatus, Coleoptera in timber of, in Burma, 693.

Sarcophaga, Dirhinus reared on, 212; larvae of Helicobia australis compared with, 57.

Sarcophaga aldrichi (parasitic on Malacosoma disstria), bionomics of, in Minnesota, 646.

Sarcophaga barbata, parasite of Vanessa cardui in Transcaucasia, 587.

Sarcophaga depressa, parasite of grasshoppers in Australia, 545.

Sarcophaga lambens, parasite of Protoparce sexta in Florida, 251.

Sarcophaga reversa, parasite o grasshoppers in Illinois, 75.

Sarcophaga sinuata, parasite of grasshoppers in Illinois, 75.

Sarcophagulopsis trigonophymi, parasite of Trigonophymus arrogans in Argentina, 214.

sarcostega, Tortrix (Cacoecia).

sarothamni, Bryobia.

Sarothamnus vulgaris, new mite on, in Holland, 301.

sasakii, Calandra.

Saskatchewan, Loxostege sticticalis on beet in, 27; miscellaneous pests in, 600; grasshoppers in, 543, 600.

satellitia, Pholus.

Satinwood, West Indian (see Zanthoxylum).

satyriniformis, Melittia.

saucia, Peridroma (Lycophotia).

savagei, Neochryopus. Sawfly, Black Wheat-stem (see

Trachelus tabidus).

Sawfly, Pine (see Acantholyda erythrocephala and Diprion pini). Sawfly, Plum (see Hoplocampa flava and H. minuta).

Sawfly, Rape (see Athalia rosae). Sawfly, Rose (see Neosyrista

similis).

Sawfly, Spruce (see Diprion poly-

Sawfly, Wheat Stem (see Cephus cinctus).

Sawfly, Yellow-headed Spruce (see Pikonema alaskensis).

Sawdust, in baits for grasshoppers, 240.

sayi, Chlorochroa.

scaber. Dendrosoter.

Scaeva pyrastri (see Lasiophthicus). scalaris, Saperda.

Scale, Black (see Saissetia oleae). Scale, Citricola (see Coccus pseudo-

magnoliarum).

Scale, Citrus Red (see Aonidiella aurantii).

Scale, Cottony Cushion (see Icerya purchasi).

Scale, Juniper (see Diaspis visci). Mussel (see Lepidosaphes ulmi).

Scale, Obscure (see Chrysomphalus obscurus).

Scale, Purple (see Lepidosaphes beckii).

Scale, San José (see Aspidiotus perniciosus).

Scale, Violet (see Parlatoria oleae). Scale Insects, list of introduced, in Argentina, 87; of Central Asia, **336**: of Br. Columbia, **511**; of France, 48, 103; records of, in Germany, 223; of Italy, 48, 511; of Madeira 392; book on,

subject to quarantines in Russian Union, 58; of S. Sakhalin, 456; physiological factors affecting infestation by, 112, 113; natural enemies of, 15, 19, 21, 23, 24, 25, 34, 40, 87, 126, 128, 129, 212, 266, 270, 290, 294, 303, 304, 308, 316, 323, 335, 345, 349, 366, 367, 385, 424, 434, 437, 446, 454, 455, 471, 501, 502, 503, 506, 546, 558, 575, 589, 611, 617, 665 ants associated with, 323, 363, 611, 665; Septobasidium associated with, 242; microtechnique for testing reactions of, to oil insecticides, 143; classification and new species of, 48, 103, 224, 235, 290, 392, 402, 456, 623, 624, 680.

scamander, Papilio.

Scapteriscus vicinus, Larra americana introduced into Porto Rico from Brazil against, 85, 435; in other countries, 85.

Scaptomyza flaveola, on horse-radish in Germany, 110.

scarabaeoides, Phloeotribus.

Scelionids, records of parasitic, in Germany, 352.

Sceliphron caementarium, possibly predacious on Laphygma frugiperda in Porto Rico, 503.

Scepticus insularis, on tobacco in Japan, 210.

schineri, Carpomyia.

schioedtei, Chilocorus. schistaceana, Eucosma.

Schistocerca gregaria, in E. Africa, 478, 482, 570; in N. Africa, 477, 478, 481; in S. and S. W. Africa, 93; in Fr. W. Africa, 481, 487; in Arabia, 480, 481, 482; in Belgian Congo, 476; in India, 291, 484, 485, 599, 560; in Persia, 482; in Spain, 477; in Anglo-Egyptian Sudan, 479, 480; phases of, 48, 477, 481, 482, 484, 486, 487, 603; morphology and pigmentation of, 48, 486; outbreak centres and ecology of, **480**, **481**, **482**, **487**; migrations of, 478, 487; effects of temperature, humidity and rainfall on, 291, 480, 481, 487, 603; correlation of outbreaks of, with sun spots, 485; reactions of, to light, 603; effects deficient diets on, 604; respiration of, 104; micro-organisms infesting, 486; measures against, 477, 478, 481.

Schistocerca paranensis, in Argentina, 476, 543; in Paraguay, 543; measures against, 476.

Schizobremia coffeae, introduced into Kenya from Tanganyika against Pseudococcus kenyae, 665.

Schizoneura lanigera (see Eriosoma). Schizotetranychus hindustanicus, on Citrus and Melia in India, 128. schmutzi, Limothrips.

schneideri, Dasyneura (see D. alpes-

Schoenobius bipunctifer (incertellus) (on rice), measures against hibernating, in China, 613; in India, Philippines, in parasites of, 577, 620.

schwartzi, Anuraphis (Appelia).

Sciapteron regalis (see Paranthrene). Sciara, transmitting Phomopsis on pistachio in Greece, 453; measures against, on mushrooms in Maryland, 380.

sciogama, Stenoma.

Scirpophaga innotata, on rice in

Philippines, **451.**

Scirpophaga monostigma, on sugarcane in India, 291, 558; damage caused by, to sugar-cane varieties, 558; parasites of, 291.

Scirpophaga nivella, on sugar-cane in India; in India; 291, 558, 575; damage caused by, to sugarcane varieties, 558, 575; parasites

Scirtothrips aurantii, factors affecting damage to Citrus by, in S. Rhodesia, 14; spray against, 14.

Scirtothrips citri (Citrus Thrips), measures and tests of insecticides against, in U.S.A., 377, 425, 432-434, 460, 649; bionomics of, 432; characters distinguishing Frankliniella moultoni and, 433.

Scirtothrips dorsalis, type of injury to chillies by, in India, 494.

Scirtothrips signipennis (Banana Rust Thrips), bionomics of, in Queensland, 329; on Citrus, 330; factors affecting outbreaks of, **329, 330**; measures against, **330**. scitella, Leucoptera (Cemiostoma).

Sclerodermus, parasite of Catorama

herbarium in Fiji, 366. Sclerotinia pseudo-tuberosa, warm water treatment of chestnuts

favouring infestation by, 359. Scolia rubea, establishment

against Lachnosterna smithi in Mauritius, 611.

Scoliopteryx libatrix, on poplar and willow in Morocco, 196.

Scolothrips longicornis, bionomics of, on soy beans in Crimea, 43.

Scolothrips sexmaculatus, bionomics of. in U.S.A., 410.

scolyticida, Coeliodes.

Scolytids, classification and new species of, 512, 615. (See Barkbeetles.)

scolytivorus, Dendrosoter.

Scolytus destructor (see S. scolytus). Scolytus multistriatus (on elm), in Britain, 65; in Holland, 396; in U.S.A., 68, 190, 378, 529, 599, 644; factors determining outbreaks of, 397; effect of sunlight and location on infestation of logs by, 190: associated with Dutch elm disease, 65, 68, 378, 396, 397, 529; crotch injury to elm-twigs by, 77, 644; bionomics of, 65, 396, 397; measures against, 66, 397, 529.

Scolytus praeceps, on Abies concolor in U.S.A., 284, 324; associated with Spicaria anomala, 284; new Braconid parasite of, 324.

Scolytus rugulosus, measures against, on peach in Illinois, 527.

Scolytus scolytus (on elm), 77; bionomics of, associated with Dutch elm disease in Britain and Holland, 65, 396, 397; measures against, 66, 397.

Scolytus subscaber, associated with Spicaria anomala on Abies concolor in California, 284.

Scolytus sulcatus, bionomics of, on elm, apple, etc., in U.S.A., 77.

Scolytus ventralis, associated with Trichosporium symbioticum Abies concolor in California, 284. scolytus, Scolytus.

Scorodite Ore, tests of, as an insecticide, 313.

Scorpion Fly (see Harpobittacus australis).

Scorpions, destroying locusts in S. Africa, 394.

Scorzonera tau-saghyz, pests of, in Russian Union, 44, 679.

scorzonerae, Xerophilaphis.

Scots Pine (see Pinus sylvestris). scripta, Sphaerophoria.

scrophulariae, Anthrenus.

scrupulosa, Teleonemia.

sculptilus, Prosoestus.

scutatus, Platychirus. scutellaris, Apanteles.

scutellata, Tritneptis (Coelopisthia). scutellatus, Gonipterus; Pachy-

nematus.

Scutigerella immaculata, movements of, in soil in California, 411. Scyllina picta, in Argentina, 476.

Scymnodes lividigaster, introduced into Porto Rico against Sipha flava, 382.

Scymnus, introduced into Kenya against Pseudococcus kenyae, 665; predacious on Tetranychus telarius in Queensland, 603.

Sea-gulls, destroying Cleonus punctiventris in Czechoslovakia,

470.

Seal Oil, effect of spraying with, on bud development of pear, 94.

secalella, Tinea.

secalis, Trachea (Apamea).

sedecimpunctata, Eutetrapha (Saperda).

Seed-corn Beetle (see Agonoderus lecontei).

Seeds, X-rays for detecting pests in, 147, 440.

segetum, Agrotis (Euxoa, Feltia). segregata, Tiphia.

segregata, auct. Parasetigena (see Phorocera silvestris).

Seius, predacious on Paratetranychus pilosus in Connecticut, 429.

Selandria sixi, reaction of Trichogramma to eggs of, 35.

Selatosomus (see Corymbites). selecta, Ptychomyia (Bessa).

selenaria, Boarmia.

Selenium, use of sprays containing, inadvisable on *Citrus*, **460.** (See Sodium Selenate.)

Selenothrips rubrocinctus, on cacao and mango in W. Indies, 24, 25, 239; attempted utilisation of Dasyscapus parvipennis against, 25, 239; dusting against, 24.

sellatus, Xylopsocus.

Selocide, 460.

Semasia diniana (see Enarmonia). semblidis, Trichogramma.

seminarius, Bruchus (Bruchidius). semipunctata, Lissopimpla.

semistriatus, Microphanurus (Telenomus).

semivelutina, Hexamitodera.

seneciella, Hylemyia.

Senecio jacobaea, failure to establish Tyria jacobaeae against, in Australia, 282; insects introduced into New Zealand against, 449.

Senecio vulgaris, lettuce mosaic in, in Britain, 671.

senegalensis, Hyperaspis.

seniculum, Apion.

Senna (Stored), Tortilia viatrix in, in New Jersey, 29.

separata, Chrysopa oculata.

sepia, Hylemyia.

septemfasciata, Nomadacris.

septendecim, Magicicada.

septentrionis, Arrhenodes (see A. minutus).

Septobasidium, monograph on, associated with Coccids, 242.

Sequoia, restrictions on importation of, into Britain, 356.

Sequoia gigantea, Phloeosinus bicolor

on, in France, 225.
seriatus, Dasus (Gonocephalum);

Psallus. sericariae, Sturmia (Crossocosmia).

sericea, Trichispa. sericeus, Agriotes; Metamasius.

serinopa, Nephantis. Serradella (see Ornithopus).

Serratia marcescens, disease in Zootermopsis caused by, in California, 597.

serricorne, Lasioderma.

sertifer, Diprion (Neodiprion).

Sesamia, on sugar sorghum in Italy, 198.

Sesamia cretica, bionomics of, on maize and sugar sorghum in Italy, 174, 197, 198, 295, 296; on sorghum in Sudan, 574; parasites of, 296, 574; measures against, 296.

Sesamia inferens, measures against,

on rice in China, 613.

Sesbania, pests of, in Formosa, **363.** Setaria italica, Chilo simplex on, in Manchuria, **340.**

setariae, Carolinaia (Hysteroneura). setarius, Calomycterus.

setiger, Lacon.

setipennis, Bigonicheta.

Setomorpha, nomenclature of species of, in stored tobacco in Netherlands Indies, 50.

Setora nitens, parasites of, on tea in Java, 347.

Sexava, parasites and biological control of, on coconut in New Guinea Territory, 22, 277; tests of insecticides against, 277.

sexdens, Atta. sexdentatus, Ips.

sexmaculatus, Scolothrips; Tetranychus.

sexnotata, Macrosteles (Cicadula).

sexta, Protoparce.

sexualis, Besserioides.

seychellarum, Icerya.

sevchellensis, Microphanurus.

Seychelles, miscellaneous pests and biological control in, 335, 336.

Shallot, Thrips tabaci on, in Louisiana, 372.

Sheep, *Helicobia australis* reared on, in Australia, **57.**

sheldoni, Eriophyes.

Shellac, in repellent paint against pests of books, 366.

shiraii, Nurudea.

Shorea robusta, list of Longicorns

attacking, 612.

Shortland Islands, attempted introduction of *Liothrips urichi* into, 366.

Sialis lutaria, in Britain, 390; unsuitable host for Trichogramma evanescens, 36; natural host of T. semblidis, 36, 390.

Siam, cotton pests in, 49.

Siberia, new Anthomyiid on cereals in, 311; grain mites in, 312; locusts in 145, 146, 147, 320; fungus infesting Loxostege sticticalis in, 307; wireworms in, 678; key to Acridids of western, 694.

sibirica, Aelia. sibiricus, Aeropus (Gomphocerus).

sicarius, Apanteles.

Sicily, Geometrid on Citrus in, 630; Bruchid on lupins in, 201, 495; miscellaneous pests in, 5, 452, 522; Polychrosis botrana on vines in, 516; compulsory fumigation of cottonseed against Platyedra gossypiella in, 174; occurrence of males of Icerya purchasi in, 5.

Sida, Platyedra gossypiella on, in Sudan, 572.

sidae, Dysdercus.

Sierra Leone, miscellaneous pests in, 271; parasite of Syrphids in, 40.

signata, Pagria; Plusia. signatus, Anthonomus.

signifera, Chirida; Theresia (Paratheresia) (see T. claripalpis).

signipennis, Scirtothrips.

Signiphora, parasite of Protargionia larreae in Argentina, 24.

sii, Hyadaphis.

Silica, toxicity of forms of, to Bruchus obtectus, 531.

Siliceous Earths, reducing volatilisation of nicotine, **133.** (See Bentonite and Lloyd's Reagent.)

Silicic Acid (dust of), effect of, against Calandra granaria, 120, 582.

Silk, review of Arthropods injurious to, 224.

Silkworm (see Bombyx mori).

Silkworm, Chinese Oak (see Antheraea pernyi).

Silvanus fasciatus, measures against, in oil-palm kernels in Dahomey, **349**.

Silvanus surinamensis (see Oryzaephilus).

silvaticus, Tetrastichus.

silveirai, Neolecanium.

silvestris, Phorocera.

similis, Ceratia (Rhaphidopalpa); Coccophagoides (Diaspiniphagus); Diprion; Microcryptus; Neosyrista; Porthesia (see Arctornis chrysorrhoea); Prepodes; Tachinomyia.

simplex, Anabrus; Chilo; Dasus (Gonocephalum); Erromenus; Rhizophagus; Taeniothrips.

simulans, Praon; Rhytidodera.

Sind, locusts in, 559, 560.

sinensis, Ceroplastes; Lachnosterna (Holotrichia); Promargarodes.

singularis, Otiorrhynchus.

sinica, Parasa.

Sinoxylon, new parasite of, in India, 505.

sinuosaria, Phigalia.

Sipha flava (on sugar-cane), in Florida, 657; parasites introduced into Porto Rico against, 382.

Sipha maydis, on wheat in Central Asia, 453.

Siphonaphis padi, auct. (see Rhopalosiphum prunifoliae).

Sirex, parasites and ecology of, in timber in Britain, 504; symbiosis of species of, with fungi in Germany, 401.

Sirex gigas, 401.

Sirex juvencus, 401.

Sirex noctilio, 401; introduction of parasites of, into New Zealand from Britain, 504.

Sirex phantoma, 401.

Sirex spectrum, 401.

Sisymbrium altissimum (see Norta). Sitka Spruce (see Picea sitchensis). sitkaënsis, Lasius niger.

Sitodiplosis mosellana (on wheat), increase of, in Br. Columbia, 66; bionomics and control of, in Holland, 268, 269.

Sitodrepa panicea, Catorama herbarium recorded as, in Fiji, 366; infesting stuffed lizard in Germany, 8; natural enemies of, 8,584.

Sitona, effect of damage to leguminous plants by, 454; measures against, 108.

Sitona callosa, on lucerne in Russia,

Sitona crinita, on leguminous crops in Russia, 151, 454.

Sitona cylindricollis, on lucerne in Russia, **108.**

Sitona inops, on lucerne in Russia, 108.

Sitona lineata, on peas and vetches in Russia, 454.

Sitophilus (see Calandra).

cerealella, Sitotroga measures against, in stored products in U.S.A., 172, 341, 451; use of eggs of, for rearing Trichogramma, 35, 36, 306, 391, 589, 590, 660; factors concerned in finding of eggs of, by T. evanescens, 166.

Six-spotted Leafhopper (see Macrosteles divisus).

sixi, Selandria.

Skunks, destroying Protoparce spp. in U.S.A., 251, 253.

Slag (Porous), tests of, as an insecticide, **313.**

slavonicus, Aspidiotus.

Slippery Elm (see Ulmus fulva). Sloe (see Prunus spinosa).

Smartweed (see Polygonum).

Smerinthus ocellatus var. atlanticus, poplar and willow Morocco, 196.

Smerinthus planus, Rhogas praetor parasitic on, in Japan, 624.
Smerinthus populi, suitability of,

as host for Trichogramma evanescens, 35, 36.

Smerinthus populi var. austauti, on poplar and willow in Morocco, 196.

smithi, Lachnosterna (Phytalus). Smoke, use of, against Sexava and

Melolontha, 277, 299.

Smynthurus viridis (on clover and lucerne), in W. Australia, 131, 292, 551; in New Zealand, 550; utilisation of Biscirus lapidarius against, 131, 292, 551; other measures against, 550.

Soap, spraying with, 89, 451; tests of penetration of solutions of, into tracheae of insects, 579; in mixed sprays, 115, 151, 177, 194, 199, 207, 231, 233, 235, 278, 300, 310, 332, 359, 365, 395, 446, 448, 450, 458, 501, 514, 519, 530, 540, 541, 570, 618, 620; affecting keeping quality of quassia sprays, 46; for emulsifying oils, 45, 333,

353, 525, 530, 535, 566, 626, 642 : formulae containing, 45, 89, 115, 194, 231, 233, 235, 278, 310, 331, 333, 359, 365, 395, 448, 458, 514, 530, 540, 541, 570, 618, 620.

Soap, Coconut-oil, in sprays, 71. Soap, Fish-oil, in sprays, 128, 498, 560; in repellent paint

Chrysobothris, 420.

Soap, Resin, in sprays, 128, 560. Scapstone, as carrier for fluorine dusts, 431.

Society Islands, Dacus psidii in,

294. (See Tahiti.)

Sodium Aluminium Fluoride (see Cryolite).

Sodium Arsenate, against termites, 422; reactions of wireworms to, 28; penetration of leaves by, **309.** (See Sodium Pyroarsenate.)

Sodium Arsenite, in baits, 67, 68, 92, 98, 130, 134, 240, 260, 291, 320, 390, 476, 478, 537; dusting and spraying with, against locusts, 92, 479; soil treatment unsatisfactory against Naupactus leucoloma, against termites, 422; reactions of wireworms to, 28; tests of toxicity of, to insects, 134, 537, 627; formulae containing, 67, 92, 98, 260, 320, 390, 476, 478; causing injury to plants, 68, 627.

Sodium Benzoate, in baits for grasshoppers, 134.

Sodium Caseinate, dusting sulphur made wettable by, 460.

Sodium Carbonate, uses of, in sprays, 278, 279, 420, 449, 451, of, in 460; interactions of, with other sprays on Citrus, 278, 279; formulae containing, 278, 449, 560.

Sodium Chlorate, spraying with, to destroy plants, 333, 375; and nicotine, 333.

Sodium Chloride (see Salt).

Sodium Cyanide, used for fumigating with hydrocyanic acid (q. v.), 29, 174, 260, 357, 440, 508, 637; factors affecting reaction between sulphuric acid and, 508.

Sodium Dinitro-ortho-cresylate, in dormant sprays against orchard

pests, 360, 533, 534.

Sodium Fluoaluminate (see Cryo-

lite).

Sodium Fluoride, in baits, 32, 249, 321, 322, 510, 537; in baitfor Argyresthia, 232; spray against Blattids, 367; tests of toxicity of, to insects, 537, 627.

Sodium Fluosilicate, in baits, 67, 131, 520, 537, 609, 652; dusting with, 17, 82, 108, 191, 239, 283, 322, 417, 431, 497; carriers for, 191, 283, 431; effect of electric charges and carriers on adhesiveness of dusts of, 309, 310; use of, against Rhizopertha in pith helmets, 612; tests of toxicity of, to insects, 537, 627; formulae containing, 17, 67, 131, 283, 431, 520, 609, 652; and calcium arsenate, 283; and injury to plants, 191, 309; other insecticides compared with, 17, 431, 537, 627.

Sodium Hydroxide, in formula for "instantaneous" oil emulsions,

Sodium Lauryl Sulphate, as a spreader and wetting agent, 33, 377, 642.

Sodium Oleate, as a spreader for nicotine, 332; affecting deposits of oil sprays, 411. Sodium Oleyl Sulphate, as an

emulsifier for oils, 642.

Sodium Pentachlorphenate, against termites, 422.

Sodium Pyroarsenate, in bait for cutworms and grasshoppers, 67. Sodium Selenate, effect of soil

treatment with, on cotton and

its pests, 220, 221, 222.

Sodium Silicate, for removing lead residues from apples, 189, 190. Sodium Tetraborate (see Borax).

Sogata furcifera (on rice), in Japan, 362; in Manchuria, 340;

position of, 362.

Soil, sampling of, for pests, 153, 317, 318, 411, 656, 679; measures for insect control causing depletion of, 342; regulations dealing with treatment of potting, against pests in U.S.A., 591, 592; effects of character of: on wireworms, 96, 97, 206, 418, 501, 679, 683; on other pests, 46, 107, 113, 131, 177, 206, 222, 351, 384, 411, 430, 431, 454, 457, 507, 519, 574, 578, 592, 593, 651, 663.

soja, Trionymus.

sokolowskii, Tetrastichus. solanella, Lita (see Phthorimaea operculella).

solani, Macrosiphum (Aulacorthum); Phenacoccus.

solanifolii, Macrosiphum (Illinoia). Solanum, reactions of Leptinotarsa decemlineata to species and hybrids with potato of, 107, 158, 169,

357, 358; Phthorimaea operculella not attacking certain species of, 289.

Solanum aethiopicum, Helopeltis bergrothi on, in Sudan, 574.

Solanum auriculatum, Dacus ferrugineus on, in Madras, 577.

Solanum chacoense, **107.**

Solanum demissum, 107, 169, 357, 358.

Solanum dulcamara, possible reservoir of potato virus diseases in Britain, 236; preferred foodplant of Leptinotarsa decemlineata, 107.

Solanum henryi, **107.**

Solanum jamesii, 107.

Solanum melongena (see Egg-plant). Solanum nigrum, Epilachna vigintioctomaculata on, in Korea, 362; Myzus persicae on, in S. Rhodesia, destruction of, against 125; Phthorimaea operculella in Victoria, 389; possible reservoir of diseases of tobacco and potato, 125, 236.

Solanum polyadenium, 107.

Solanum triquetrum, Paratrioza cockerelli on, in Texas, 651. Solanum tuberosum (see Potato).

Solar Heat, reactions of insects to, 603, 604, 692; timber, cottonseed, etc., exposed to, against pests, 181, 190, 573, 574, 576, 612.

Solar Oil, 627.

Solenopsis geminata, attacking vegetables in Jamaica and Br. attacking Guiana, 239, 524; predacious on Nymphula depunctalis in Philippines, 364.

solitarius, Apanteles.

Solomon Islands, attempted introduction of Liothrips urichi into. 366.

solstitialis, Amphimallus (Rhizotrogus).

Solubea pugnax, bionomics of, on rice and grasses in U.S.A., 537.

sommeri, Hypopholis.

somnulentella, Bedellia.

sonchi, Macrosiphum.

Sonchus, destruction of, against Hercothrips fasciatus in Cali-fornia, 240; Phytomyza atricornis on, in Japan, 444; pests on, in Russian Union, 44, 677.

Sonchus asper, experimentally infected with lettuce mosaic, 671.

Sonchus oleraceus, insects on, in Britain, 61, 672.

Soot, repellent to adults of Psila

rosae, 497. Sooty Moulds, insects associated with, **365**, **524**, **657**; possibly encouraged by sprays containing sugar, **425.**

Sophia spp., relation of Eutettix tenellus and curly-top virus to,

in Idaho, 286.

Sophora japonica, Protargionia larreae on, in Argentina, 23.

sophorae, Brassolis.

Sorbus americana, Scolytus sulcatus on, in U.S.A., 77.

Sorbus aucuparia, Erannis spp. on, in Poland, 438.

sordidus, Acanthocoris; Cosmopolites.

sorditaria, Erannis leucophaearia.

Sorghum, Nomadacris septemfas-ciata associated with, in Africa, 62, 570; pests of, in India, 128, 560; Pyrausta nubilalis on, in Manchuria, 340; relation of Aphis maidis and, to sugar-cane mosaic in Queensland, 88; Loxostege sticticalis on, in Russia, 307; pests of, in Sudan, 572, 574; termites damaging cotton in rotation with, in Tanganyika, 222; pests of, in U.S.A., 248, 261, 274, 341, 374, 376, 381; resistance of varieties of, to pests, 274, 341; use of, in baits for locusts, 478; (broom corn), HCN fumigation of, under winter conditions against Pyrausta nubilalis in Canada, 416.

Sorghum vulgare var. saccharatum (Sugar Sorghum), pests of, in Italy, 197, 198, 199, 295, 472; virus disease of, 472.

soror, Diabrotica.

Soxhlet Method, 142.

Soy Bean, Scolothrips longicornis on, in Crimea, 43; pests of, in Japan, 19, 103, 210, 338, 361, 556, 624; Cydia glycinivorella on, in Korea, 19; Aphids transmitting virus disease of, in Germany, 686; pests of, in Manchuria, 19, 340; Ootheca mutabilis on, in S. Rhodesia, 125; pests of, in U.S.A., 67, 179, 358, 370, 371, 373, 422; susceptibility varieties of, to Anomala rufocuprea, 210; Bruchus chinensis seldom developing on, "safened" calcium arsenates injuring, 258; (stored), Aphomia gularis in crushed residue of (1875)

imported, in Germany, 520; Plodia interpunctella in, 225.

Soy-bean Caterpillar (see Anticarsia

gemmatalis).

Soy-bean Flour, effect on Plodia interpunctella of feeding on, 173; uses of, in sprays and dusts, 70, 208, 373.

Soy-bean Oil (see Oil, Soy-bean). Soy-bean Pod Borer (see Cydia

glycinivorella).

Spain, pests of fruit trees in, 6; pests of wild lupins and serradella in, 495; locusts in, 477; danger of spread of Leptinotarsa decemlineata into, 583; beneficial insects in, 477, 495; Aphelinus mali introduced into Poland from, 162.

Spalangia cameroni, parasite of Dacus and Musca in Fiji, 366. Spalangia philippinensis, parasite

of Dacus ferrugineus in Madras,

Sparganothis pilleriana (on vines), measures against, in Germany, 103; in Russian Union, 403.

Sparrows, grain mites spread by, 552.

Spartium junceum, males of Icerya purchasi on, in Sicily, 5.

Spathius critolaus, sp. n., parasite of Coleoptera in India, 505.

Spathius labdacus, sp. n., parasite of Pempheres affinis in India, 505. spectabilis, Dendrolimus; Pimpla

(Exeristesoides).

spectrum, Sirex (Xeris). Spelt (see Triticum spelta).

Spermophagus cisti (see Bruchus).

spermotrophus, Megastigmus. Sphaerakis mayri, parasite of Bruchus brachialis in U.S.A., 599.

Sphaerolecanium (see Lecanium). Sphaerophoria flavicauda, bionomics and morphology of, in

Britain, 628, 629. Sphaerophoria scripta, bionomics and morphology of, in Britain, 628, 629.

Sphecodina abboti, parasitised by Apanteles congregatus in U.S.A., 253.

Sphedanolestes impressicollis, predacious on Rhynchites cupreus in Japan, 556.

Sphegigaster flavicornis, parasite of Phytomyza ilicis in Britain, 669; by Pleurotropis parasitised amyntas, 670.

sphenophori, Ceromasia. sphenotoma, Aeolarchis.

Sphinx chersis, parasite of, in U.S.A., 253.

Sphinx kalmiae, parasite of, in U.S.A., 253.

Sphinx pinastri (Pine Sphingid), bionomics of, in Poland, 676.

Spicaria anomala, Scolytus spp. associated with, on Abies concolor in California, 284.

Spicaria fumosorosea, penetration of chitin of insects by, 307.

Spicaria heliothis, sp. n., infesting Heliothis armigera in U.S.A.,

Spiders, destroying insects, 251, 282, 288, 338, 374, 500.

Spilochalcis albifrons, parasite of Bedellia somnulentella in Kansas, 375.

Spilocryptus adustus, parasite of Diprion similis in Poland, 671.

Spilonota (Eucosma) ocellana (on fruit trees), in Baluchistan, 51, 577; in Canada, 662, 664; in China, 668; in Czechoslovakia, 468; in New York, 535; bionomics of, 51, 668; measures against, 535, 662, 664, 668.

Spilosoma lubricipeda (see Diacrisia). Spilostethus equestris (see Lygaeus). spilota, Arthrochlamys.

spilotus, Anthonomus.

Spinach, virus disease in, in Britain and Germany, 672, 688; Hymenia recurvalis on, in Japan, 441; Piesma quadratum and crinkle disease of, in Poland, 163, 351, 352: Eutettix tenellus on, in Texas, 594.

Spinaria bicolor, parasite of Setora nitens in Java, 347.

spinarum, Athalia (see A. rosae). spinicornis, Hoplocerambyx.

spinipes, Maleuterpes. spinolae, Lygus.

spinosus, Dectes.

spiraecola, Aphis.

splendana, Cydia (Carpocapsa, Laspeyresia).

splendens, Rhopalosiphum.

Spodoptera exempta (see Laphygma). Spodoptera mauritia, measures against, on rice in Assam, 577; parasite of, on sugar-cane, etc., in Formosa, 363.

sponsor, Cryptus.

Sprays, types of equipment for applying, 199, 247, 280, 302, 383, 438, 536, 561; method of applying concentrated, 383; use of aircraft for applying, 74, 645; factors affecting deposits of, 52, 143, 272, 411, 412; compared with, 177, 179, 249, 266, 495, 497, 498, 514; aeroplane dusting compared with, 610; programmes for: for orchards in N. America, 104, 534, 623, 662, 664; for Citrus in Argentina, 23; for orchard and small fruits in Britain, 396.

spretus, Melanoplus (see M. mexi-

canus).

Spruce (Picea), restrictions on importation of, into Britain, 356; pests of, in Canada, 275, 276, 368, 369, 382, 463, 464, 540, 647; pests of, in Czechoslovakia, 469, 517, 518, 564; pests of, in Germany, 230, 334, 398, 400, 401, 563, 580, 581; Pissodes spp. on, in Finland, 8, 9; Lymantria monacha on, in Poland, 398; pests of, in Sweden, 122, 123; pests of, in U.S.A., 368, 369, 382, 529, 540, 647; Acantholyda erythrocephala not observed on, in U.S.A., 651; symbiosis of Siricids with fungi on, 401; effect of feeding L. monacha on, 399; effects of insecticides on, 400, 529, 530.

Spruce, Black (see Picea mariana). Spruce, Blue (see Picea pungens).

Spruce, Norway (see Picea abies). Spruce, Red (see Picea rubra).

Spruce, Sitka (see Picea sitchensis). Spruce, White (see Picea glauca). Spruce Budworm (see Harmologa

fumiferana). Spruce Sawfly (see Diprion polytomum).

Spruce Sawfly, Yellow-headed (see

Pikonema alaskensis). spuria, Gossyparia.

spurius, Apanteles.

sputator, Agriotes. Squash, Melittia satyriniformis on, in Wisconsin, 33.

Stahlia, timber of, practically immune from termites in Porto Rico, 465.

Stantex R and Dispersing Oil, composition and uses of, as insecticides, 530.

staphyleae, Rhopalosiphoninus.

Staphylinus caesareus, predacious on Pentatomids in Germany, 632. Staphylinus olens, predacious on Pentatomids in Germany, 64.

Staple Fibre, insects damaging, in Japan, **440.**

Starch, Aleurodes brassicae fed on media containing, 61.

Statistical Methods, application of, to entomological problems, 240,

Stauroderus scalaris, in Siberia, 320. Steam, use of, against hibernating larvae of Cydia in orchards, 383; regulations for using, against Naupactus in U.S.A., 592.

stecki, Pristiphora (Lygaeonematus)

(see P. pallida).

Stegobium panicea (see Sitodrepa). Steirastoma depressum, on cacao in Grenada, 25.

Stellaria media, Erythroneura pallidifrons overwintering on, in greenhouses in Britain, 115; Rhizoecus kondonis on, in Japan, 617; Micromyzus oliveri on, in Washington, 541.

Stemtox A, 537.

Stenaraeoides octocinctus, parasite of Dendrolimus spectabilis in Japan, 211.

Stenobothrus, in Jugoslavia, 480.

Stenobothrus apricarius (see Chorthippus).

Stenobothrus haemorrhoidalis Omocestus).

stenocentrus, Polyblastus.

Stenocranophilus quadratus, parasite of Saccharosydne saccharivora in Porto Rico, 657; suggested introduction of, into Florida, 657. Stenocranus, on coconut in Sey-

chelles, 335.

Stenoma sciogama, on Anonaceae in Brazil, 566.

Stenomalus micans, parasite Chloropids in Britain, 562.

Stenomalus muscarum, mass occurrence of, in houses, 201.

Stenothrips graminum, on wheat in Sweden, 121.

Stenotus binotatus, type of injury to wheat by, in New Zealand, 548.

Stephanitis pyri, on fruit trees in Czechoslovakia, 468; on chestnut in Greece, 452.

Stephanitis rhododendri, on rhododendron in Britain, 295.

Stephanoderes hampei (Coffee Berry Borer), absence of, in Italian E. Africa, 175; utilisation Prorops nasuta against, in Brazil, 188, 238; treatment of coffee seed against, in Kenya, 667; in Uganda, 188.

Sterculia, relation of Dysdercus spp. to, in Africa, 489, 490, 573.

sternale, Trogoderma.

Sternochetus (see Cryptorrhynchus). Stethophyma fusca (see Arcyptera).

Stethorus, predacious on Paratetranychus pilosus in Connecticut,

Stethorus punctillum, predacious on Tetranychus turkestani in Central Asia, **626.**

sticticalis, Loxostege.

sticticraspis, Diatraea (Argyria). Stictocephala festina, bionomics and control of, on leguminous crops in Louisiana, 370, 371.

stigma, Hemerobius.

stigmaterus, Microdus (Bassus). stigmatias, Euxesta.

stigmaticus, Coeloides.

Stilpnotia salicis, parasites and biological control of, in U.S.A.,

Stizolobium, Aphis on, in Australia, 448.

stokesi, Gambrus.

Stomatoceras pomonellae, parasite of Cydia spp. in Australia, 293, 640.

Stomatodexia diadema, parasite of Diatraea spp. in S. America and Trinidad, 183; proposed introduction of, into St. Lucia, 183.

Stored Products, pests of, in Egypt, 219; book on pests of, and their control in Germany, 513.

stouti, Polycaon.

stramineus, Diversinervus.

Strawberry, pests of, in Britain, 97, 395; spray programme for, in Britain, 396; Harpalus rufipes on, in Denmark, 117; pests of, in Germany, 155, 457, 466; Galerucella distincta on, in Korea, Cerosipha forbesi on, in Ontario, 276; Tarsonemus pallidus on, in Switzerland, 280; pests of, in U.S.A., 31, 165, 178, 179, 372, 374, 376, 378, 380, 425, 498; wheat intersown with, against wireworms, 97; effects of feedingpunctures of C. forbesi on, 276; Aphids transmitting virus diseases of, 395; susceptibility of varieties of, to virus diseases, 395.

Strawberry Blossom Weevil (see

Anthonomus rubi).

Strawberry Leaf-roller (see Ancylis

comptana). Strawberry Mite (see Tarsonemus

pallidus). Strawberry Root Aphis (see Cerosipha forbesi).

Strawberry Weevil (see Anthonomus signatus).

striatellus, Delphacodes.

striatum, Anobium (see A. punctatum).

striatus, Laevicephalus (Deltocephalus).

strigator, Calyptus.

String, made from wood fibre, insects damaging, in Japan, **440**. Striped Morning Sphinx (see *Gelerio lineata*).

strobi, Chermes (Pineus); Pissodes. strobilobius, Chermes (Cnaphalodes). Stromatium, in timber, **493**.

Stromatium barbatum, distribution of, in timber, 465, 611, 612.

Stromatium fulvum, in imported furniture and building timber in U.S.A., 160, 600; longevity of, 160.

Stromatium longicorne, in furniture in Netherlands Indies, 493.

Strophostyles umbellata, Chalcodermus aeneus on, in U.S.A., 77.

Strumeta (see Dacus).

Sturmia, parasite of Protoparce spp. in U.S.A., 253.

Sturmia bella, parasite of Vanessa cardui in Transcaucasia, 587.

Sturmia halli, sp. n., parasite of Heliothis armigera in S. Rhodesia, 567.

Sturmia inconspicua (parasite of Diprion spp.), in Germany, 398; in Poland, 671.

Sturmia inconspicuoides, parasite of Cirphis loreyi in Queensland, 186.

Sturmia protoparcis, parasite of Protoparce sexta in Florida, 251.

Sturmia sericariae (parasite of Bombyx mori in Japan), galvanometer for detecting, in host pupae, 442; parasitised by Mormoniella vitripennis, 443.

Stylocryptus brevis, parasite of spruce sawflies in Czechoslovakia, 518.

Stylocryptus erythrogaster, parasite of spruce sawflies in Czechoslovakia, 518.

subarctica, Pristiphora (Lygaeonematus).

Subcidine, 605.

subcompleta, Microgaster.

subcrinita, Epitrix.

subfumatus, Eutelus.

subguttatus, Microcryptus.

subnigrescens, Mermis.

subnudus, Anacentrinus.

subproximus, Anagyrus.

subscaber, Scolytus.

subsequalis, Pyrausta (see P. insequalis).

subsidiaria, Tortrix (Cacoecia). subterranea, Symphylella.

Subterranean Clover (see Trifolium subterraneum).

subvittatus, Prosoestus (Derelomus).

succincta, Patanga.

Sudan, Anglo-Egyptian, cotton pests in, 488, 572, 573, 574; pests of Sorghum in, 572, 574; Ceroplastes africanus in, 40; locusts in, 479, 480, 481; beneficial insects in, 40, 574.

Sudan Cotton Bollworm (see Dipa-

ropsis castanea).

Suet, Necrobia spp. in, 584.

Sugar, types of, used for feeding insects, 61, 128, 345, 426, 682; in baits, 92, 102, 131, 194, 291, 466, 496, 582; in bait-sprays for *Cydia*, 300; use of, in sprays and dusts against thrips, 425, 526; possibly encouraging sooty moulds on *Citrus*, 425. (See

Syrup.)

Sugar-cane, Diatraea on, in Br. Guiana, 182, 183, 505, 524; pests of, in Formosa, 206, 339, 363, 364, **444, 446, 447**; pests of, in Hawaii, 90, 367, 368; pests of, in India, 128, 291, 385, 456, 558, 575, 576; pests of, in Mauritius, 611; pests of, in Queensland, 87, 88, 185, 186, 187; little attacked by pests in Peru, 165; pests of, in Philippines, 368, 451, 667; pests of, in U.S.A., 248, 371, 372, 373, 378, 382, 656-659; pests of, in W. Indies, 24, 25, 182, 183, 238, 272, 382, 384, 424, 435, 502, 543, 657, 658, 659, 660; food-plant of Euxesta stigmatias, 328; susceptibility of varieties of, to pests, 24, 87, 88, 186, 238, 367, 372, 378, 384, 444, 543, 558, 575, 657; effect of borers on weight and sucrose content of, 558; insects and virus diseases of, 87, 658, 667; insecticides causing injury to, 88, 368.

Sugar-cane Aphis, Yellow (see Sipha

flava).

Sugar-cane Beetle Borer (see Rhab-docnemis obscura).

Sugar-cane Moth Borer (see Diatraea saccharalis).

Sugar-cane Rootstock Weevil (see Anacentrinus subnudus).

Sugar Maple (see Acer saccharum).

Sugar Sorghum (see Sorghum vulgare var. saccharatum).

sulcatus, Dendrosoter; Otiorrhynchus; Scolytus.

sulmo, Microphanurus.

Sulphite Lye, dust mixture containing, 431; as an emulsifier, 332.

Sulphur, against Aphis spiraecola, 461; effect of, on Coccid populations, 257, 377, 433; against Rhynchota on cotton, 244, 245, 246, 282, 380, 381, 417; against other Rhynchota, 72, 421, 463, 537, 595, 651; against mites, 55, 57, 127, 150, 164, 187, 256, 257, 308, 410, 429, 433, 460, 461, 595, destroying predacious of Paratetranychus enemies pilosus, 429; against Psila rosae, 497; against thrips, 71, 72, 377, 433, 460; against weevils, 249, 380; ineffective against Cantharis obscura, 623; dusting with, 71, 72, 150, 164, 187, 244, 245, 246, 282, 308, 377, 380, 381, 417, 429, 433, 461, 463, 623, 637, 651; carriers for, 187, 380; adsorbed on carriers, 150, 308; on carriers, 130, 308; on-impregnated dusts of, 308; in mixed dusts, 71, 179, 191, 213, 244, 245, 246, 247, 250, 371, 380, 381, 383, 417, 418, 421, 431, 461, 463, 537, 595; spraying with, 71, 72, 187, 256, 257, 429; wetting agents for, 70, 71, 80, 257, 460; aluminium sulphate with lime as an adhesive for, 256, 257; in mixed sprays, 52, 57, 71, 72, 139, 186, 410, 433, 460, 534, 595, fumigation tests with, 662; against ants, 44, 287; types of, 72, 80, 187; formulae containing, 52, 57, 72, 179, 186, 187, 244, 245, 246, 247, 250, 256, 257, 308, 380, 381, 383, 410, 417, 418, 460, 537, 595; basic copper arsenate compatible with, 421; effect of, on lead arsenate spray residues, 139; thiuram sulphides compatible with, 532; addition of zinc compounds to dusts of, preventing injury to Citrus, 433; interactions of, with other sprays on Citrus, 278, 425; causing injury to onion, 71, 72; unsuitable for use on hemp for smoking, 127; methods for determining particlesize distribution of, 541.

Sulphur Nitride, as an insecticide,

78, 79.

Sulphuric Acid, and sodium cyanide, for generating hydrocyanic-acid gas (q.v.), 260, 357, 440, 508.

sulphurifera, Lissonota. sumatranus, Dacryphalus.

sumatrensis, Compsilura concinnata.

Sun Spots, correlation of locust outbreaks with, 485.

Sunflower, pests of, in Russian Union, 321, 682.

Sunoco Oil, 435.

Supella supellectilium, paint for protecting books from, in Fiji, 366.

Superphosphate, ineffective against *Halotydeus destructor*, **545**; effect of, mixed with wheat before sowing against wireworms, **97**.

superstitiosus, Dysdercus.

suppressaria, Buzura.

surinamensis, Oryzaephilus (Silvanus).

suspensa, Anastrepha.

suturellus, Dysdercus.

Swede, pests of, in Britain, 96, 332, 333; pests of, in Denmark, 117; Meligethes aeneus on, in Germany, 511.

Sweden, Thysanoptera on cereals and grasses in, 121; Apion spp. on clover in, 117; forest pests in, 122, 123; miscellaneous pests in, 355, 673; Calandra spp. in stored grain in, 118–120; Aphids of, 160; Leskia aurea in, 629; migrations of Pieris brassicae in, 609.

Sweet Cassava (see *Manihot dulcis* var ai pi).

Sweet Pea, experimentally infected with lettuce mosaic, 671.

Sweet Potato, locusts damaging, in Fr. W. Africa, 571; pests of, in Japan, 444, 445, 446, 620; Brachmia macroscopa on, in Korea, 362; pests of, in Loochoo Islands, 444, 616; pests of, in U.S.A., 285, 373, 379, 539, 643; pests of, in W. Indies, 239, 434; calcium chloroacetate injuring foliage of, 239; (tubers), treatment of, against Cylas formicarius, 379, 539; as baits for Cylas and wireworms, 373, 656.

Sweet Potato Weevil (see Cylas formicarius).

Switzerland, forest pests in, 169, 517; Leptinotarsa decembineata in, 106, 350, 583; miscellaneous pests in, 270, 280; orchard pests in, 169, 232, 336, 438; Phylloxera on vines in, 46; Acarapis woodi infesting honey bees in, 169; Stenomalus muscarum in houses in, 201; summary of plant quarantines in,

344; legislation against introduction of L. decemlineata into Britain from, 356.

Svagrus morio, on cotton in Nyasaland, 126.

sycophanta, Calosoma.

Sylepta derogata, Eulophus parasitic on, in Japan, 557; in Fiji, 641; in Seychelles, 335; in Siam, 49, 50; on cotton, 49, 50, 557; on Hibiscus, 335, 641.

syma, Chrysocharis.

Symitha mangifera, on mango in Seychelles, 335.

Symphylella subterranea, movements of, in soil in California, 411.

Sympiesis turcicus, sp. n., parasite of Lithocolletis platani in Turkey, 219.

Synanthedon (see Aegeria).

Syngenaspis oleae (see Parlatoria). Synopeas, parasite of Diplosis mori in Korea, 338.

Syntomosphyrum indicum (parasite of Dacus), introduced into Cook Islands, 365; establishment of, in Fiji, 213; introduced into Hawaii, 366; introduced into South Queensland from New Wales, 602; introduced into Samoa, 365; synonymy of, 434.

Syria, in dates in Egypt, 506.

Syringa (see Lilac).

Syringopais temperatella, effect of meteorological conditions on cereals in Palestine, 127.

Syrphids, of Illinois, immature stages of, 82.

Syrphus balteatus, bionomics and morphology of, in Britain, 628,

Syrphus luniger, bionomics and morphology of, in Britain, 628, 629.

ribesii, bionomics Syrphus and morphology of, in Britain, 628, 629.

Syrphus torvus, predacious Anuraphis roseus in Connecticut,

Syrphus tricinctus, insects destroyed by, in Germany, 334.

Syrup, in baits for Cydia, 31, 293; in mopping mixture and spray against Anthonomus, 247.

Systates exaptus, on maize and Dolichos in S. Rhodesia, 124.

Systates saeuberlichi, on cotton in Kenya, **666**.

Systena hudsonias, measures against, on potato in U.S.A., 342.

Systena taeniata blanda, measures against, on potato in U.S.A., 342.

T.

tabaci, Thrips.

tabacum, Hypopteromalus.

Tabanus, predacious on Nymphula depunctalis in Philippines, 364.

tabatae, Eutelus. tabidus, Trachelus.

tachardiae, Apanteles.

Tachinids, list of, in Rumania, 608; classification of, 624.

Tachinomyia similis, parasite of Stilpnotia salicis in U.S.A., 499.

Tachycines asynamorus, bionomics and control of, in greenhouses in U.S.A., 651, 652.

Tachypterellus quadrigibbus (Apple Curculio), in New York, 533; measures against, in Quebec, 664. taeniata, Systena.

taeniopus, Chlorops.

Taeniothrips fedorovi, bionomics of, on Salvia sclarea in Crimea, 43.

Taeniothrips inconsequens, Crimea, 43; in U.S.A., 377, 410, 498; on fruit trees, 43, 377, 410, 498; on maple, 498; bionomics of, 498; measures against, 410, 498.

Taeniothrips simplex, tests of greenhouse fumigant against, 272.

Tahiti, Plaesius javanus introduced from Fiji into, against Cosmo-polites sordidus, 212, 365.

takagii, Rhythmonotus.

Talc, toxicity of dusts of, to Bruchus obtectus, 531; repellent to Psila rosae, 497; as a carrier for dusts, 32, 139, 179, 191, 247, 258, 261, 286, 313, 325, 371, 379, 380, 431, 449, 461, 523, 525, 537, 595, 623, 634; affecting toxicity of certain insecticides, 139, 258, 261: preventing detonation of sulphur nitride, 79.

Tamarix, Icerya purchasi on, in Morocco, 691.

tamsuyensis, Melanotus.

tanaceti, Galeruca. tanajoa, Tetranychus.

Tanganyika Territory, Aleurodids and cassava mosaic in, 94, 153; coffee pests in, 40, 126; cotton pests in, 222, 492; locusts in, 387, 480, 484; beneficial insects in, 40, 127, 387, 503; Schizobremia coffeae introduced into

Kenya from, 665.

Tank-mixture Method, for preparing sprays, 432.

Tanymecus destructor, on maize in S. Rhodesia, 124.

Tanymecus palliatus, food-plants and control of, in Russian Union, 45, 322, 407; bionomics of, 322. Tapinoma melanocephalum, in

houses in Formosa, 446.

Tar Distillates, as dormant sprays against orchard pests, 20, 115, 232, 236, 237, 333, 360, 428, 429, 467, 534, 535, 578, 662, 690; interfering with natural control of Paratetranychus pilosus, 429, 662; against pests of conifers, 517, 529; against vine pests, 45, 689; against Coccids on other plants, 402, 509; soil treatment with, against *Rhagoletis cerasi*, **297**; as a repellent, **299**, **420**; tests of toxicity and action of, on insects, **467**; fish killed by, 514; types of, 45, 65, 297, 298, 428, 467, 535; formulae for, 20, 45, 428, 517, 529, 534, 535, 689; preparation of emulsions of, 420; and mineral oils, 236, 534, 535; as a carrier for paradichloro-benzene against borers, 527; other insecticides combined with, 298, 689; and injury to plants, 45, 65, 527, 535; in spray for destroying potato plants and Aphids, 333; other insecticides compared with, 236, 237, 429, 467, 509, 529, 534.

taraxaci, Macrosiphum.

Taraxacum dens-leonis (see T. officinale).

Taraxacum kok-saghyz, pests of, in Russian Union, 44, 45.

Taraxacum megalorrhizon, pests of, in Russian Union, 44.

Taraxacum officinale (Dandelion), Aleurodes brassicae on, in Britain, 61; pests breeding on, in Russian Union, 44, 45.

Targionia vitis, natural enemies of, on vines in Italy, 471.

Tarnished Plant Bug (see Lygus pratensis).

Taro (see Colocasia).

Tarred Disks, against Hylemyia brassicae, 116.

tarsale, Trogoderma.

Tarsoneminae, of America and Hawaii, 224.

Tarsonemus, on mushrooms in Pennsylvania, 79.

Tarsonemus bancrofti, measures against, on sugar-cane in U.S.A., 657.

Tarsonemus floricolus, on mushrooms and iris in U.S.A., 79.

Tarsonemus pallidus (fragariae) (on strawberry), in Britain, 395; ecology of, in Switzerland, 280; measures against, 280, 396.

measures against, 280, 396.
Tartar Emetic, in bait-spray for Trypetids, 256; dusts and sprays containing, against thrips, 425, 525; Citrus not injured by small quantities of, 257; papers treated with solution of, against Ctenolepisma, 510.

Tarucus theophrastus, parasites of,

in India, 129.

Tasmania, pests of ornamental plants in, 21, 226; orchard pests in, 20, 21, 159, 226; vegetable pests in, 20, 21, 510; beneficial insects and biological control in, 20, 21, 294; Habrolepis dalmanni introduced into Australia from, against Asterolecanium variolosum, 546.

tau, Aglia.

Tau-saghuiz (see Scorzonera tau-saghyz).

taurella, Ochsenheimeria.

tavaresi, Aphis. taveuniensis, Calotermes.

Tea, Homona coffearia on, in Ceylon, 19; Monema flavescens on, in Japan, 207; pests of, in Java, 94, 347; pests of, in Formosa, 339, 340, 447; mites on, in India, 128; Helopettis bergrothi a potential pest of, in Nyasaland, 126; Pulvinaria floccifera on, in Transcaucasia, 455.

Tea Bark-borer (see Linoclostis gonatias).

Teak, pests of, in Burma, 692, 693; pests of, in India, 612.

Teak Borer (see Phassus malabaricus).

Technomyrmex, hindering biological control of Coccids in Seychelles, 335.

Technomyrmex detorquens (albipes), fostering Coccids on coconut in Mauritius, 611; measures against, 611.

Tectona grandis (see Teak).

tectus, Ptinus.

tegalensis, Chionaspis.

telarius, Tetranychus.

Telenomus, parasite of sugar-cane moth borers in Formosa, 447.

Telenomus alecto, parasite of Diatraea saccharalis in Porto Rico, 424.

Telenomus beneficiens, parasite of sugar-cane moth borers in Formosa, 364, 447.

Telenomus beneficiens var. elongatus, parasite of sugar-cane borers in Formosa, 447.

Telenomus clisiocampae, parasite of Malacosoma disstria in Minnesota,

Telenomus mitsukurii, parasite of Megymenum gracilicorne in Japan,

Telenomus semistriatus (see Microphanurus).

Telenomus tischleri, sp. n., parasite of Dolycorus baccarum in Germany, **632.**

Telenomus ullyetti (parasite of Heliothis armigera), bionomics of, in Rhodesia, 16; proposed utilisation of, 17.

Telenomus vassilievi, parasite of Eurygaster integriceps in Russia, 555; suggested utilisation of, 555.

Teleonemia lantanae (destroying Lantana), establishment of, in Australia, 546, 603; shipped to New Caledonia and New Hebrides from Fiji, 212; introduced into Tonga, 212, 366; considered a synonym of T. scrupulosa, 366.

Teleonemia morio, on Anonaceae in Brazil, **566.**

Teleonemia scrupulosa, T. lantanae considered a synonym of, 366.

Temnoschoita quadripustulata (quadrimaculata), bionomics and control of, on oil palms and banana in Dahomey, 348.

temperatella, Syringopais.

Temperature, relations of locusts and grasshoppers to, 11, 12, 41, 92, 93, 146, 178, 345, 388, 393, 427, 480, 483, 484, 485, 553, 557, 569, 603, 604, 655; relations of the ripsects to 45, 46, 47, 26, 57 other insects to, 15, 16, 17, 26, 27, 40, 41, 42, 47, 59, 60, 61, 63, 64, 99, 102, 118, 119, 120, 121, 146, 147, 149, 200, 216, 217, 218, 224, 225, 228, 229, 230, 234, 235, 254, 255, 262, 265, 266, 267, 299, 300, 305, 313, 314, 315, 316, 325-327, 337, 351, 357, 360, 361, 362, 374, 399, 403, 404, 405, 406, 408, 428, 441, 442, 451, 454, 455, 489, 491, 550, 553, 557, 564, 572, 586, 587, 588, 589, 590, 606, 607, 613, 615, 617, 622, 647, 658, 662, 663, 678, 681, 682, 687, 692; relations of mites to, 131, 147, 151, 209, 312, 313, 314, 318, 319, 552, 585, 625, 626, 672; control of, in small cabinets, 568. (See Bioclimatics.)

Tenebrio molitor, measures against, in flour mills in Russian Union, 685; suitability of, as a host for Trichogramma evanescens, nutritional studies on, 624; tests of tar distillates on 467.

tenebrioides, Zabrus.

Tenebroides mauritanicus, in stored products in China, 451; associated with Lasioderma serricorne in stored tobacco and cumin seed Cyprus, 270; in flour in Germany, 400; in Italy, 173; damaging string made from wood fibre in Japan, 440; in flour mills in Russian Union, 685; in stored grain in U.S.A., 341; predacious on other pests, 173, 400; toxicity of HCN to, 451.

Pipunculus tenellus, Eutettix; trochanteratus.

tener, Polyblastus.

Tennessee, tobacco pests in, 252, 380, 652; beneficial insects in, 253, 254; parasites of Apanteles congregatus in, 254.

tenthredinis, Mesoleius.

tenuicornis, Bothriophryne (Trichomasthus); Frankliniella. Tenuipalpus, measures against, on

vines in Queensland, 365.

Tenuipalpus oudemansi, sp. n., on apple and beech in Holland, 301.

Tenuipalpus phoenicis, sp. n., on Phoenix in greenhouse in Holland,

tenuis, Heterotermes (Leucotermes). tenuiscapa, Xylocopa.

Tephritis, on chrysanthemums in Britain, 333; in Canary Islands, 333.

Tephrosia, cultivation of, for insecticides in Belgian Congo, 294; survey of data on, as an insecticide, 104.

Tephrosia toxicaria, pests of, in Porto Rico, 502.

Tephrosia virginiana, toxicity of dusts of, to Pieris brassicae, 250, 251.

Tephrosia vogeli, possibilities of, as a source of rotenone, 502.

Termes assmuthi, damaging sugarcane in India, 558, 559.

Termes badius, in building timber in S. Rhodesia, 125.

Termes pauperans, in building timber in S. Rhodesia, 125.

Termes transvaalensis, in building timber in S. Rhodesia, 125.

Terminalia, catabba, Lasiccampid

Terminalia catappa, Lasiocampid on, in India, 385.

terminaliae, Eupelmus. terminalis, Eutane.

terminifera, Chortoicetes.

Termites, important genera of, in Australia, 545; of Italian E. Africa, 174; of W. Indies, 55, attacking living plants, 89, 128, 222, 271, 464, 465, 523, 528, 544, 546, 558, 559; attacking timber, 55, 56, 125, 180, 181, 212, 239, 434, 464, 465, 498, 499, 545, 641, 693; susceptibility of timbers to, 239, 434, 465, 545; international exposure test of wood preservatives against, 180; measures against, 55, 89, 128, 180, 181, 222, 271, 465, 544, 693, 694; method of evaluating soil poisons against, 422; bacterial diseases of, 597; gaseous environment of, 282; classification and new species of, 384.

terrae-reginae, Empoasca.

terrenus, Bruchus (Bruchidius).

Tessaratoma javanica, attacking lac host-trees in India, 128.

testacea, Luperina (Apamea); Palpostoma.

testaceicollis, Pristomerus.

testaceimanus, Lygocerus.

testaceus, Coptotermes; Laemophloeus; Rhogas.

testudinea, Hoplocampa.

testulalis, Maruca.

tetensi, Dasyneura.

Tetracampe, parasite of Phytomyza ilicis in Britain, 669.

Tetracampe diprioni, parasite of Diprion similis in Poland, 671. Tetracampe ruforum (see Achryso-

Tetracampe ruforum (see Achrysocharella).

Tetrachlorethane, tests of, against Rhagoletis cerasi, 297.

Tetracyphus odontomus, on cotton in Kenya, 666.

Tetrahydronaphthalene, ineffective against Cylas formicarius, 539.

Tetramethyl Thiuram Bisulphide (see Thiuram Sulphides).

Tetramorium guineense, in Formosa, 363, 446; infesting houses, 446; fostering Aphids and Coccids, 363; utilisation of, against sugar-cane moth-borers, 364.

Tetraneura fusiformis (see Byrso-

crypta).

Tetraneura ulmi (see Byrsocrypta gallarum).

Tetranychus, on vines in Germany, 103; on Hibiscus in India, 128; bionomics of, on apple in Japan, 209, 616; food-plants of, in Russian Union, 150, 308, 310, 313, 314, 403; effects of temperature and humidity on, 209, 313, 314; measures against, 103, 150, 308, 310.

Tetranychus bioculatus (see Paratetranychus).

Tetranychus bituberculatus, on pear in Japan, 225.

Tetranychus fici, on figs in India, 128. Tetranychus hindustanicus (see

Schizotetranychus).

Tetranychus pacificus, hibernation of, on apple in Washington, 525. Tetranychus sexmaculatus, on Citrus in U.S.A., 256, 261; measures against, 261.

Tetranychus tanajoa, sp. n., measures against, on cassava in

Brazil, 290.

Tetranychus telarius, on hops and mangels in Britain, 1, 332; on Hydrangea in Germany, 583; food-plants of, in India, 127; Scymnus predacious on, on papaya in Queensland, 603; food-plants of, in U.S.A., 79, 141, 261, 432, 583, 595, 649, 650; disseminating Uromyces caryophyllinus on carnations, 79; measures and tests of insecticides against, 1, 45, 127, 141, 142, 261, 272, 432, 583, 595, 649, 650.

Tetranychus turkestani, factors affecting infestation of cotton by, in Central Asia, 147, 625, 626; Stethorus punctillum predacious on, 626.

Tetrastichus, hosts of, in India, 576, 577; parasite of Crioceris asparagi in Italy, 472; hosts of, in Japan, 144, 212; introduced into U.S.A. against Galerucella luteola, 144, 324; other hosts of, in U.S.A., 422, 528, 569; bionomics of, 324, 472; effect of cold storage on reproduction of, 144.

Tetrastichus brevistigma (parasite of Galerucella luteola), bionomics and value of, in U.S.A., 132.

Tetrastichus coccinellae, parasite of Chilocorus spp. in Caucasus, 304. Tetrastichus coerulescens, parasite of Gelechia cercerisella in Kansas,

Tetrastichus giffardi, utilisation of, against Ceratitis capitata in Brazil, 34.

Tetrastichus giffardianus (parasite of Trypetids), introduced into Cook Islands and Samoa, 365; introduced into Fiji, 212, 213; bionomics of, 212, 213.

Tetrastichus haitiensis, utilisation of, against Prepodes spp. in

Jamaica, 239.

Tetrastichus silvaticus, parasite of Malacosoma disstria in Minnesota, 597.

Tetrastichus sokolowskii, parasite of Plutella maculipennis in India, 575, 576; bionomics of, 576.

Tetrastichus xanthomelaenae (parasite of Galerucella luteola), introduced into U.S.A., 324; bionomics of, 324.

Tettigoniella viridis (see Cicadella). texana, Anomis (see A. luridula);

Atta.

texanus, Microdus (Bassus).

Texas, Atta texana in, 287; cotton pests in, 53, 244, 245, 380, 381, 413, 417, 538; Eutettix tenellus in, 593, 594; Paratrioza spp. in, 422, 651; miscellaneous pests in, 379, 537, 599; Mymarid parasites of Psallus seriatus in, 538.

Textiles, pests of, and their control, 47, 140, 187, 188, 196, 237, 513,

584, 635.

Thalassodes, parasite of, of India, 129.

Thallous Sulphate, tests of, in baits for grasshoppers, 134.

Thanatophilus quadripunctatus, predacious on Erannis spp. in Poland, 438.

Thaneroclerus buqueti, predacious on Lasioderma serricorne in Mauritius, 611.

Thaumetopoea pityocampa, bionomics and control of, on pine in Italy, 516.

Thaumetopoea processionea, bionomics and control of, on oak in Italy, 516.

Thecodiplosis mosellana (see Sito-diplosis).

theobaldi, Thomasiniana.

Theophrastus, as an economic entomologist, **592**.

theophrastus, Tarucus.

Therapha hyoscyami, on tobacco in Poland, 162.

Theresia [Tachinid], Paratheresia considered distinct from, 143.

Theresia [Zygaenid] (see Theresimima).

Theresia claripalpis (parasite of Diatraea saccharalis), attempted

introduction of, into Antigua from Trinidad, 25, 660; question of introduction of, into U.S.A., 143, 656, 658, 659; identity, forms and factors affecting distribution of, 143, 658, 659.

Theresia diatraeae, considered distinct from T. claripalpis (q.v.),

143.

Theresia signifera, considered distinct from T. claripalpis (q.v.), 143.

Theresimima ampelophaga, on plums in Cyprus, 271; on vines in Russian Union, 403, 683; use of females for trapping males of, 683.

Theretra oldenlandiae, on vines in Queensland, 365.

Therioaphis ononidis (see Myzo-callis).

thermoptera, Titanoceros.

Theronia atalantae, hosts of, in Czechoslovakia, 145; parasite of Dendrolimus spectabilis in Japan, 211; parasite of Ichneumon pisorius in Poland, 677.

Thespesia rogersi, Dysdercus fasciatus on, in N. Rhodesia, 222, 492; experiment with D. nigrofasciatus

and, 490.

Thiocyanates, against Aphids, 236, 237, 259, 332, 533, 534; against Coccids and Aleurodids, 259; against Capsids, 236, 463; against Jassids, 413, 501; against Lepidoptera, 155, 236, 237, 332; against mites, 1, 259, 430; ovicidal properties of, 1, 236, 331, 332, 533, 534; spray formulae containing, 1, 236, 258, 259, 413; spreaders and emulsifiers for, 259, 331; atomised sprays of oils containing, 155, 413; dusting with, 463; and lime-sulphur, 259; and oil emulsions, 236, 237, 259, 332; and pyrethrum, 413; causing injury to plants, 236, 534.

Thiodiphenylamine, against Cydia pomonella, 66, 142, 273, 528, 653; against other Lepidoptera, 17, 377, 379, 431; ineffective against Scirtothrips citri, 425; tests of toxicity of, to insects, 142, 249, 287, 528, 537; in baits, 287, 537; formula for dusting with, 431; formulae for, in sprays, 66, 249, 379; wetting agent for, 377; and cryolite, 379; thiuram sulphides compatible with, 532; other insecticides compared with, 66, 79, 249.

Thiuram Sulphides, in spray formulae as repellents for *Popillia* japonica, 532.

Thomasiniana crataegi, sp. n., on hawthorn in Britain, 673.

Thomasiniana oculiperda, crataegi compared with, 673.

Thomasiniana theobaldi, T. crataegi compared with, 673.

thomsoni, Piestopleura.

thoracicus, Macrocentrus; Triaspis. Thosea, on coconut in Philippines, 451.

thrax, Erionota.

Thrips, measures against, on cotton in Peru, 164; on wheat in Sweden, 121.

Thrips angusticeps, on peas in Holland, 233; characters of, 233.

Thrips lini (linarius), on flax in Holland, 233, 234, 465; bionomics of and type of injury by, 233; synonymy and characters of, 233; measures and tests of insecticides against, 233, 234, 465.

Thrips tabaci, new Gamasid predacious on, in Britain, 672; bionomics of, in Crimea, 43; in Japan, 210; in Kenya, 666; in New Zealand, 546; in Porto Rico, 436; in U.S.A., 72, 141, 372, 525; on cotton, 43, 372, 672; on tobacco, 43, 210, 546; on vegetables, 43, 71, 72, 372, 436, 546; on other plants, 141, 525, 666; transmitting spotted wilt, 546; measures against, 71, 141, 142, 436, 525, 666.

Thrips, Banana Rust (see Scirto-

thrips signipennis).

Thrips, Bean (see Hercothrips fasciatus).

Thrips, Cacao (see Selenothrips rubrocinctus).

Thrips, Citrus (see Scirtothrips citri).

Thrips, Flower (see Frankliniella moultoni).

Thrips, Onion (see *Thrips tabaci*). Thrips, Pear or Prune (see *Taenio-*

thrips inconsequens).
Thrips, Wheat (see Haplothrips tritici).

thripsi, Typhlodromus.

Thuja, restrictions on importation of, into Britain, 356; Phloeosinus bicolor on, in France, 225; insecticides causing injury to, 530.

thujae, Phloeosinus.

Thurberia, Platyedra gossypiella scarce on, in Arizona, 381.

Thyanta custator, and boll rots in cotton in Arizona, 416, 417; measures against, 417.

Thymus, relation of Dociostaurus maroccanus to, in Transcaucasia,

7.....

thymus, Hyssopus.

Thysanoptera, list of, in Louisiana, 372; cellular structure of grasses as protection from, 121; possibly causing leaf-curl of chillies, 494; and plant diseases, 360, 448, 472; predactious species of, 371, 410, 429, 453; natural enemies of, 25, 239, 672; new species of, 50.

Tibicen haematodes, on vines and

sloe in Germany, 46.

Tiger Pear (Opuntia aurantiaca) (see Opuntia).

Tilia (see Lime, Tilia).

Tilia amurensis, Eutetrapha sedecimpunctata on, in Russian Far East, 43.

Tilia glabra, Alsophila pometaria on, in Canada, 663.

Timber, Coleoptera in, 6, 7, 60, 78, 110, 160, 181, 202, 203, 204, 268, 321, 322, 328, 335, 465, 469, 493, 499, 509, 520, 564, 584, 600, 611, 612, 693; age of, affecting infestation by Hylotrupes bajulus, 520; termites in, 55, 56, 125, 180, 181, 212, 239, 434, 464, 465, 498, 499, 545, 641, 693; other pests of, 61, 181, 239, 337, 342, 401, 498, 499, 504, 562, 693; analysis of, for preservatives, 280.

Timocratica haywardi, sp. n., on fruit trees in Argentina, 188.

Tinea columbariella, characters of eggs of, 49.

Tinea fuscipunctella, characters of eggs of, 49.

Tinea granella, bionomics of, in stored grain in Germany, 220; in stored products in Italy, 172; in flour mills in Russian Union, 685; measures against, 220, 685.

Tinea misella (insectella), synonymy of, **50.**

Tinea pellionella, tests of mothproofing preparations against, 140; characters of eggs of, 49.

Tinea pistaciae, bionomics of, on pistachio in Greece, 452, 453.

Tinea secalella, sp. n., bionomics of, in Germany, 220; in Poland, 220.

Tinea vastella, measures against, damaging horns in Belgian Congo, 294.

tineivorus, Typhlodromus.

Tineola biselliella, infesting house insulation in U.S.A., 640; tests of moth-proofing preparations and fumigants against, 140, 635.

Tingis pyri (see Stephanitis).

Tiphia, parasite of Lamellicorns in Florida, 657.

Tiphia hispaniolae, sp. n., bio-

nomics of, in Haiti, 84.

Tiphia popilliavora, utilisation of, against Popillia japonica U.S.A., 84, 377, 533; influence of instars of P. japonica on, 84; strains of, 377.

Tiphia segregata, parasite of Anomala orientalis in Hawaii, 367.

Tiphia vernalis, in U.S.A., 377, liberation of, against 533; Popillia japonica, 533; seminating diseases of P. japonica,

tiphiaeformis, Synanthedon (see

Aegeria typhiaeformis).

Tipula, control of, in golf greens, 95; other measures against, 519. Tipula czizeki, damaging cereals and pastures in Germany, 519.

Tipula oleracea, bionomics of, in Britain and Germany, 95, 519. Tipula paludosa, on wheat in Denmark, 117; bionomics of, in

Britain and Germany, 95, 519. Tipula vernalis, observations on, in Britain, **95, 96.**

tipulae, Leptacis.

Tiracola plagiata, on cacao in New Guinea Territory, 544.

Tirathaba trichogramma, Tineid feeding on pupae of, in Fiji, 213.

Tischeria complanella, on oak in Germany, 204.

Tischeria omissa, bionomics of, on hollyhock in California, 569. tischleri, Telenomus.

Titanoceros thermoptera, bionomics of, in Australia, 545.

Tits, destroying Cydia pomonella, 114; question of danger of arsenical insecticides to, 159.

Toads, destroying insects, 160, 185, 407, 452, 641, 656; damaging trap-ditches in beet fields, 407.

Tobacco, pests of, in S. Africa, 360; Phthorimaea operculella on, in Brazil, 289; Agrotis ypsilon on, in Fiji, 213; Aphids on, in Germany, 633; locusts damaging, in Greece, 478; pests of, in

Japan, 210, 444; new Aleurodid on, in Madagascar, 637; heliopa on, in Mysore, 625; pests of, in Netherlands Indies, 17, 493, 637; Heliothis virescens on, in Philippines, 452; Rhynchota on, in Poland, 162; pests of, in S. Rhodesia, 124, 125, 226; S. Rhodesia, 124, 125, 226; pests of, in Russia, 41, 43; pests of, in U.S.A., 84, 177, 178, 251, 252, 253, 254, 380, 599, 652; H. armigera on, in Victoria, 389, 390; incorts and circuit. 390: insects and virus diseases in, 84, 124, 125, 226, 302, 303, 360, 444, 493, 546, 547, 637, 688; effects of insecticides on, 177,

Tobacco (Stored), pests of, and their control, 50, 125, 154, 196, 210, 234, 235, 270, 380, 513, 611.

Tobacco (as an insecticide), dusting with, 177, 353, 667; as a carrier for dusts, 177, 250, 371, 667; tests of sprays containing, against Bemisia gossypiperda, 560; action of, against Psila rosae, 497.

Tobacco Beetle (see Lasioderma

serricorne).

Tobacco Hornworm (see Protoparce). Tobacco Stem Borer (see Phthorimaea heliopa).

Tobago, Coptotermes testaceus in, 56, 384.

todo, Xyleborus.

Toluene, and carbon tetrachloride, as a solvent for dinitro-o-cyclohexylphenol, 261.

Tolyphus, on Scorzonera tau-saghyz

in Kazakstan, 44.

Tomaspis, on sugar-cane in Florida, 658.

Tomaspis guppyi, bionomics and control of, on sugar-cane in Trinidad, 543.

Tomaspis saccharina, 543; soil types and varieties of sugar-cane in relation to, in Trinidad, 24, 384;

measures against, 24.

Tomato, Paratrioza cockerelli on, in greenhouses in Alberta, 176; pests of, in Australia, 187, 281, 389, 449, 603; pests of, in greenhouses in Britain, 115, 333; pests of, in Fiji, 641; Tetranychus telarius on, in India, 127; Phthia picta on, in Jamaica, 239; Acanthocoris sordidus on, in Japan, 446; new mite on, in Morocco, 636; relation of Thrips tabaci to spotted wilt in, in New Zealand, 546, 547; Myzus persicae on, in S. Rhodesia, 125; pests of, in

Asiatic Russia, 589, 609, 679; pests of, in U.S.A., 72, 73, 79, 142, 252, 253, 340, 372, 379, 408, 422, 431, 525, 599, 651, 652; effect of infestation by *Paratrioza cockerelli* on, 72, 73, 651; not attacked by *Phthorimaea operwiella* 280; approximation of the content of the culella, 289; as reservoir of virus diseases of other plants, 125, 688; insects caught by secretion and spines of leaves of, 113, 142; tops of, in baits for Listroderes obliquus, 186; lime-sulphur causing injury to, 72, 73.
Tomato (Fruit), legislation dealing

with fumigation of, against Popillia japonica in U.S.A., 274; problems of insecticide residues

on, **79, 432.**

Tomato Mite (see Phyllocoptes lyco-

persici).

Psyllid Tomato (see Paratrioza cockerelli).

Worm Tomato (see Protoparce sexta).

tomentosus, Lyctus.

Tomicobia, synonymy of, 434.

Tonga, introduction of beneficial insects from Fiji into, 212, 366, 641.

torquens, Contarinia (Dasyneura) (see C. nasturtii).

Tortilia viatrix, measures against, in stored senna in New Jersey, 29.

Tortrix dumetana, parasites of, on ramie in Transcaucasia, 587.

Tortrix heparana, on soy beans in Manchuria, **340.**

Tortrix lafauryana, parasites of, on ramie in Transcaucasia, 587.

Tortrix murinana, on Abies alba in Czechoslovakia, 469.

Tortrix podana, tests of winter washes against, on apple in Britain, 236, 237.

measures Tortrix postvittana, against, on vines in S. Australia,

Tortrix pronubana, imported into Germany with carnation cuttings from Italy, 218; on poplar and willow in Morocco, 196; study of, 218.

Tortrix rosaceana, on dewberry in

Utah, **31.**

Tortrix sarcostega, bionomics of, on fruit trees in Baluchistan, 51. Tortrix subsidiaria, on fruit trees in

Baluchistan, 577.

Tortrix viridana, on rhododendron in Britain, 295.

Tortrix xylosteana, bionomics and control of, on fruit trees in Japan, 361.

torvus, Syrphus.

Toxoptera graminum, possible vector of disease of sugar sorghum in Italy, **472.**

trabealis, Emmelia.

Trachea basilinea, on wheat in Central Asia, 453; measures against, 454.

Trachea secalis, rôle of harvesting machinery in control of, on wheat in Russia, 315; bionomics

of, in Scotland, 122.

Trachelus tabidus (Black Wheatstem Sawfly), in U.S.A., 83, 381; Collyria calcitrator introduced from Canada against, 381.

tragopoginis, Amphipyra.

transsylvanica, Hypera (Phytonomus).

transvaalensis, Termes.

trapezalis, Marasmia. trapezina, Calymnia.

Traps, for Coleoptera, 42, 90, 184, 347, 410, 457, 466, 532; mobile types of, for Meligethes aeneus, 511; for Forficula auricularia, for Lepidoptera, 466, 577, 510; 614, 683, 689; for Trypetids, 270. (See Light-traps.)

Treacle (see Molasses).

Tree Pear (Opuntia streptacantha) (see Opuntia).

Tremex spp., symbiosis of, with wood-destroying fungi, 401.

tremulae, Melasoma (Chrysomela). Trenches, against Celerio lineata, 239; against locusts, 450, 477, 618; against weevils on beet, 407, 470, 678.

Trialeurodes vaporariorum (Greenhouse Whitefly), in Britain, 333, 356, 437; in Czechoslovakia, 471: utilisation of Encarsia formosa against, 333, 437; fumigants against, 272, 356; effect of adding chloropicrin to dusts against, 471.

triangularis, Microbracon (Bracon). Triaspis thoracicus, introduced into U.S.A. against Bruchids, 381, 599.

Tribolium (in flour), in Germany, 223; fumigants against, in U.S.A., 378; Corynetes coeruleus probably predacious on, 584.

Tribolium castaneum, in imported coffee in Britain, 666; fumigation experiment against, in China, 451; in stored grain and house insulation material in U.S.A., 341, 640; predacious on Ephestia kuehniella, 400; feeding on potato flour, 65; characters of, 623.

Tribolium confusum, in imported coffee in Britain, 666; measures against, in flour mills in Russian Union, 685; in stored grain in U.S.A., 341; action of electromagnetic waves on, 137; HCN not causing protective stupe-344; studies of faction in, nutrition, physiology and populations of, 27, 80, 174, 624; method for determining movement of, in flour, 52; characters of, **623.**

Tribolium destructor, in imported Lantana seed in Britain, 623; characters and distribution of,

Tribolium madens, characters of,

Tribolium navale, auct. (see castaneum).

Trichacis remulus, introduced into U.S.A. against Mayetiola destructor, 381.

Trichispa sericae, bionomics and control of, on rice in Madagascar,

Trichlorethylene, fumigation with, 120, 539.

Trichlorobenzene, as a soil insecticide against termites, 422.

trichodactvla. Hylemyia (Chortophila).

Trichogramma (parasite of Lepidopterous eggs), in Canada, 663; in Central Asia, 609; in India, 577; introduction and utilisation of Burmese species of, in India, 612.

Trichogramma australicum, parasite of sugar-cane moth-borers in Formosa, 364, 447.

Trichogramma evanescens (parasite of Lepidopterous eggs), 588; in Italy, 4; in Minnesota, 597; in Poland, 677; local races, breeding and utilisation of, in Russian Union, 109, 148, 304, 305, 589, 590; characteristics affecting suitability of hosts for, 35-37, 390; factors concerned in hostfinding by, 166; T. semblidis distinct from, 390.

Trichogramma japonicum, parasite of Chilo simplex in Japan, 619; mass rearing of, 619.

Trichogramma luteum (parasite of Heliothis armigera), bionomics of, in S. Rhodesia, 16, 17.

Trichogramma minutum, in India, 612; introduced into Philippines, 184, 452; in U.S.A., 540, 656; in W. Indies, 424, 435, 660; parasitising Diatraea, 424, 435, 656, 660; parasitising other Lepidoptera, 184, 612; sawfly parasitised by, 540; rearing and utilisation of, 435, 452, 612, 656, 660; climatic factors affecting, 435, 660.

Trichogramma semblidis, bionomics of, in Britain, 36, 390, 391; distinct from T. evanescens, 390. trichogramma, Tirathaba.

Trichogrammatoidea nana, parasite of eggs of sugar-cane mothborers in Formosa, 447.

Trichomasthus, characters distinguishing Bothriophryne and, 504. Trichomasthus tenuicornis (see Bothriophryne).

Trichopsidea oestracea, parasite of grasshoppers in Australia, 56.

Trichospilus pupivora, liberation of, against Nephantis serinopa in Ceylon, 638.

Trichosporium symbioticum, Scolytid associated with, on Abies in California, 284.

Trichothrips liliorus (see Hoplothrips).

tricinctus, Syrphus.

tricolor, Encarsia.

Tricresylphosphate, papers treated with solution of, against Ctenolepisma, 510.

Triethanolamine Oleate, as an emulsifier for oils, 411, 642; affecting oil deposits of sprays,

trifasciatus, Closterocerus.

trifenestrata, Cricula,

Trifidaphis phaseoli, on cotton in Russia, 215.

Trifolium (see Clover).

Trifolium hybridum, Apion seniculum on, in Latvia, 47.

Trifolium subterraneum, Halotydeus destructor on, in Australia, 545; Collembola on, in New Zealand, 550, 551.

trigonophymi, Sarcophagulopsis.

Trigonophymus arrogans, bionomics and control of, in Argentina, 213, 214, 476.

trimaculatus, Macrocentrus (Leptozele).

Trinidad, cotton pests in, 220, 221; miscellaneous pests in, 24, 85; Tomaspis spp. on sugar-cane in, 24, 384, 543; parasites and biological control of Diatraea spp. in, 24, 183, 505; termites damaging timber and buildings in, 55, 56, 384; Larra americana apparently not occurring in, 85; beneficial insects introduced into other W. Indian Islands from, 25, 239, 502.

trinitatis, Azya.

Trionymus levis, sp. n., in Caucasus, 680.

Trionymus mori, sp. n., on mulberry in Japan, 103.

Trionymus soja, sp. n., on soy bean

in Japan, 103.

Trioza nigricornis, bionomics of, in Germany, 636; importance of, on potato, 636.

Triphleps (see Orius).

triseti, Dasyneura.

Trisetum flavescens, new Cecidomyiid on, in Austria, 635.

Trisopsis, predacious on Tyroglyphus farinae in Russian Union, 320.

triste, Opatrum.

tristicolor, Orius; Lachnosterna (Phyllophaga).

trisulcus, Megastigmus (Epimega-

stigmus).

Tritaxys braueri, sp. n., parasite of Cirphis unipuncta in Queensland, 294.

Tritaxys goniaeformis, apparently confined to Tasmania, 294; T. braueri recorded as, 294.

tritici, Contarinia; Frankliniella; Haplothrips; Hylemyia (Phorbia).

Triticum spelta, nutritive value of flour and groats of, for Ephestia kuehniella, 64.

Tritneptis, status and key to N. American species of, 434.

Tritneptis diprionis, sp. n., parasite of sawflies in U.S.A., 434.

Tritneptis hemerocampae, type of genus, 434.

Tritneptis klugi, synonymy of, 434. Tritneptis koebelei, sp. n., parasite of sawfly in California, 434.

Tritneptis scutellata, synonymy of, 434.

Triumfetta, Dysdercus superstitiosus on, in N. Rhodesia, 492.

trivialis, Colaspis.

trivittatus, Leptocoris.

trochanteratus, Pipunculus. Troctes divinatorius (see Liposcelis). Trogium pulsatorium, measures against, in houses in U.S.A., 648. Trogocarpus ballestrerii (see Megastigmus).

Trogoderma spp., review of feeding

habits of, 196.

Troilus luridus, in Britain, 632; in Germany, 218, 354, 632; predacious on noxious insects, 218, 354; parasite of, 632.

tropicalis, Lachnus (Pterochlorus). Tropidophryne, key to species of, 503; characters of, 504.

Tropidophryne flandersi, question of systematic position of, **504.**

Tropidophryne melvillei, sp. n., parasite of Pseudococcus in Kenya, 503.

Tropidophryne natalensis, sp. n., presumably parasitic on Pseudo-

coccus citri in Natal, 503.

truculenta, Bačtra.

truncatulus, Lixus (Hypolixus). tryoni, Biosteres; Dacus (Chaetodacus, Strumeta) ferrugineus.

trypetes, Chilo.

Trypetids (Fruit-flies), legislation dealing with control of, in W. Australia, 130; key to intercepted, in Russian Union, 439.

Tryphon utilis, parasite of Pristiphora wesmaeli in Germany, 217. Trypodendron spp., Rhizophagus spp. predacious on, on Picea in Sakhalin and Japan, 339.

Trypopitys carpini, in building tim-

ber in Russia, 322.

Trypterygium wilfordi, tests of insecticide prepared from **450**.

Tsuga spp., restrictions on importation of, into Britain, 356; pests of, in U.S.A., 378, 499.

tsugae, Chermes (Adelges). Tuberolachnus (see Lachnus).

tuberosula, Platygaster. tubulorum, Lepidosaphes.

Tulip, Aphids and breaking disease of, in Oregon, 369.

tulipae, Anuraphis.

tulipaella, Rhopalosiphoninus (see R. staphyleae).

tumulosus, Ligyrus.

Tung-nut Tree (see Aleuvites).

Tunisia, locusts in, 478; Phylloxera on vines in, 10; summary of plant quarantines in, 344.

turanicum, Lecanium coryli; Macrosiphum (Acyrthosiphon) gossypii.

turca, Otiorrhynchus.

turcicus, Entedon auronitens; Sympiesis.

Turf, Tipulid larvae and their control in, in Britain, 95; damaged by Blissus hirtus in Canada, 601; insects damaging, in Holland, 269; treated with lead arsenate against Popillia japonica in U.S.A., 533; damaged by B. leucopterus in Trinidad, 25.

turionana, Rhyacionia (Evetria). turkestani, Tetranychus.

turkestanicus, Phloeosinus.

Turkey, miscellaneous pests in, 218, 452, 522; beneficial insects in, 219, 522; summary of plant quarantines in, 274.

Turkeys, destroying Cleonus punctiventris in Czechoslovakia, 470.

turneri, Nysius.

Turnip, pests of, in Denmark, 117; pests of, in Louisiana, 250, 371, 373; Hylemyia radicum on, in Morocco, 605; Aphids and mosaic disease of, in New York, 273, 274; Nysius turneri on, in Tasmania, 226; in baits for Listroderes obliquus, 249.

Turpentine, wood impregnated with mixture containing, against Coleoptera, 182; oil palms treated with, against Oryctes,

347.

Tychius spp., on lucerne in Russia, 108.

Tyloderma brassicae, sp. n., on cabbage in Brazil, 132, 183; new parasite of, 183.

Tyloderma foveolata, on strawberry

in Arkansas, 178.

Tyloderma fragariae, bionomics of, on strawberry in Arkansas, 178. typhiaeformis, Aegeria (Synanthedon).

Typhlocyba froggatti (australis, Frogg.), measures against, on fruit trees and hawthorn in Australia, 21, 292; in New Zealand, 292.

Typhlodromus thripsi, sp. n. (predacious on Thrips tabaci), bionomics of, in Britain, 672.

Typhlodromus tineivorus, predacious on Ephestia kuehniella in Germany, 400.

typographus, Ips.

Typophorus viridicyaneus, bionomics and control of, on sweet potato in U.S.A., 285, 286; in Mexico, 285.

Tyria jacobaeae (attacking ragwort), failure to establish, in Australia, 282; local establishment of, in

New Zealand, 449; insects and birds destroying, 282, 449.

Tyroglyphus farinae (in stored grain and flour in Russian Union), not surviving under field conditions in Siberia, 312; predacious enemies of, 319, 320; measures against, 685.

Tyrophagus putrescentiae, rare in stored grain in Siberia, 312; not surviving under field conditions,

312.

U.

Uganda, new Encyrtid parasite of Coccids in, 503; locusts in, 482, 666.

ugandum, Apion (Piezotrachelus).

uhleri, Horistonotus.

ullyetti, Telenomus.

ulmi, Eriosoma; Lepidosaphes; Tetraneura (see Byrsocrypta gallarum).

ulmi, auct., Oligonychus (see Paratetranychus pilosus).

ulmicola, Myzocallis (Sappocallis).

Ulmus (see Elm).

Ulmus japonica, Saperda octomaculata on, in Russian Far East, 42. Ulmus propinqua (and var. suberosa), bark characters affecting susceptibility of, to pests in

Japan, 443. Ulobaris loricata, bionomics of, on

beet in Kazakstan, 152.

Ultra-sulphur, 308.

Ultrawet, as a wetting and spreading agent for sprays, 71, 72; increasing efficiency of cubé dust, 177.

umbra, Pyrrhia.

umbrinus, Microterys.

Uncaria gambir (Gambier), Oreta carnea on, in Sumatra, 385.

undalis, Hellula.

undulata, Phyllotreta.

unicolor, Byturus.

unicostata, Monosteira.

unifasciatum, Lecanium (Sphaerolecanium).

unipuncta, Cirphis.

United States of America, cereal pests in, 80, 138, 229, 230, 280, 295, 327, 378, 379, 381, 542, 599, 600, 645; Citrus pests in, 343, 377, 642, 653; cotton pests in, 242, 243, 244, 245, 246, 247, 380, 381, 599; pests of forest and shade trees in, 53, 65, 67, 68, 77, 132,

138, 170, 324, 342, 368, 369, 377, 378, 382, 392, 434, 498, 499, 528, 540, 546, 599, 600, 644, 645, 647; pests of greenhouse and ornamental plants in, 141, 142, 419, 592, 600, 652, 654, 673; pests of leguminous plants in, 82, 83, 370, 371, 381, 415, 599, 600, 645; miscellaneous pests in, 377, 380, 599, 600; orchard pests in, 272, 273, 343, 377, 382, 419, 426, **434**, **643**, **648**; pests of small fruits in, **139**, **377**, **380**, **395**, **596**; ecology and control of pasture insects in, 527; pests of solanaceous plants in, 251, 252, 253, 254, ·378, 379, 380, 418, 422, 563, 599; pests of sweet potato in, 285, 373, 379, 539, 643; storedproduct and household pests in, 341, 378, 380, 382, 600, 640, 648; official testing of moth-proofing preparations in, 140; pests of timber in, 498, 499, 600; new Aphids in, 336, 596; distribution of Empoasca fabae in, 83; grasshoppers in, 76, 138, 139, 240, 569, 538, 646, 654, 655; Popillia japonica in, 104, 138, 139, 165, 180, 274, 377, 532, 533, 591; wireworms in, 28, 41, 71, 77, 418; insects and plant diseases in, 65, 68, 77, 370, 378, 395, 563, 645; Septobasidium associated with Coccids in, 242; beneficial insects and biological control in, 80, 81, 132, 170, 251, 253, 324, 377, 378, 379, 381, 382, 410, 422, 434, 499, 540, 596, 599, 615; new fungus infesting Heliothis armigera in, 392; annual insect population records in, organisation of economic entomology in, 261, 407, 408; plant pest legislation in, 176, 274, 275, **344, 591**; list of pests intercepted in quarantine in, 462; patents relating to pest control in, 224, 394: introduction of beneficial insects into other countries from, 382, 544, 546, 640; pests imported into other countries from, 328, 616; risk of introduction of Popillia japonica into Bermuda from, 85, 86; legislation restricting importation of potatoes and apples into Britain from, 356. (See also under the various States.)

urichi, Liothrips. Urochela luteovaria, on pear in China, **450**.

Uromyces caryophyllinus, Tetranychus telarius disseminating, on carnation in Massachusetts, 79. Urosigalphus, parasite of Chalcodermus bondari in Brazil, 414. Urosigalphus chalcodermi, 414. urozonus, Eupelmus. urticae, Vanessa.

Utah, Crioceris asparagi on asparagus in, 426; forest pests in, 427, 567, 568; pests of small fruits in, 31, 596; pests of leguminous crops in, 29, 462; Anabrus simplex and grasshoppers in, 29, 48; beneficial insects and biological control in, 30, 31, 463, 596; lizards destroying insects in, 48.

utibilis, Lariophagus (see L. distinguendus) utilis, Tryphon.

V.

Vaccinium myrtillus, Erannis spp. on, in Poland, 438.

Vacuum Cleaner, for volatilising paradichlorobenzene, 644.

Vacuum Fumigation, 174, 181, 416, 541, 542, 550, 605; chemicobiological bases of improvements in, 201, 202.

validirostris, Pissodes. vanderwulpi, Cadurcia.

vaneeckei, Liothrips.

Vanessa spp., Pentatomids experimentally attacking, 354.

Vanessa cardui, parasites and diseases of, on ramie in Transcaucasia, 587.

Vanilla, Ecpantheria icasia on, in Porto Rico, 501.

Vanilla Essence, in bait for Dacus psidii, 131.

vaporariorum, Trialeurodes.

variabilis, Hypera (Phytonomus); Lacon.

variana, Peronea.

varians, Anthonomus; Aphis; Chrysomela (Chrysolina); Cyrtopeltis (see Engytatus geniculatus). varicornis, Monolepta.

variegata, Adonia; Pachyprotasis.

varinervis, Antholcus.

variolosum, Asterolecanium.

varius, Anthrenus (see A. verbasci).

varivestis, Epilachna.

vassilievi, Telenomus. vastatrix, Perkinsiella ; Phylloxera (see P. vitifoliae).

vastella, Tinea. vayssierei, Bemisia.

Vegetable Marrow, probable reservoir of virus infecting spinach in Britain, 672.

Vegetable Weevil (see *Listroderes* obliquus).

velox, Oxya.

Velvet Bean (see Stizolobium). Velvet Bean Caterpillar (see Anticarsia gemmatalis).

Venezuela, Ceratitis capitata reported in, 24; Larra americana in,

85.

venosata, Diatraea.

ventralis, Exochomus; Scolytus.

ventricosus, Pediculoides.

Venturia inaequalis, 664.

verbasci, Anthrenus.

Verbascum thapsiforme, Pentatomids on, in Germany, **353.**

Vermont, Malacosoma disstria in, 377.

vernalis, Tiphia; Tipula.

vernata, Paleacrita.

verrucivorus, Decticus.

versicolor, Ampeloeca; Meteorus;

Trogoderma. versutus, Adoretus.

verticalis, Xylocopa.

verticillata, Eupristina.

vestitus, Anthonomus; Hylesinus (Chaetoptelius).

vesuviana, Carpomyia.

Vetch Bruchid (see Bruchus brachia-

Vetches, pests of, in Russian Union, 322, 454, 552; Bruchus brachialis on, in U.S.A., 381, 599; mites in stored seed of, 318.

Vetiveria nigritana, relation of locusts to, in Fr. W. Africa. 12, 571.

vetusta, Agrotis (Abagrotis).

viatrix, Tortilia.

Viburnum opulus, Aphis fabae on, in Germany, 686.

Vicia (see Vetches).

Vicia faba (see Beans).

vicina, Musca domestica.

vicinus, Camponotus maculatus; Homorocoryphus; Scapteriscus.

victima, Crematogaster.

Victoria, Anthrenus verbasci in, 187; cutworms in, 389; Cydia spp. on fruit trees in, 20, 293, 390, 544, 639; Phthorimaea operculella on potato in, 90, 389; beneficial insects and biological control in, 293, 545, 639, 640.

vigintioctomaculata, Epilachna. vigintioctopunctata, Epilachna.

Vigna (see Cowpeas). vigorsi, Periclitena.

villica, Arctia. vindicius, Microphanurus.

Vine, Grape, pests of, in Australia, 89, 164, 293, 365, 603; regulations for spraying, against Trypetids in W. Australia, 131; Lygus spinolae on, in Austria, 45; pests of, in Brazil, 34; Paranthrene regalis on, in China, 450; Polychrosis botrana on, in Cyprus, 270; mealybugs on, in Egypt, 507; pests of, in France, 155; pests of, in Germany, 46, 102, 103, 111, 155, 156, 458, 465, 496, **582, 688, 689**: locusts damaging, in Greece, 478; Paratetranychus punicae on, in India, 128; Targionia vitis on, in Italy, 471; Rhynchota on, in Japan, 440, 441; Polychrosis botrana on, in Jugoslavia, 494; pests of, in Palestine, 191; Phylloxera on, in Poland, 403; pests of, in Russian Union, 42, 43, 403, 688; Polychrosis botrana on, in Sicily, 516; Phylloxera on, in Switzerland, 46; legislative and other measures against Phylloxera on, in Tunisia, 10, 11; pests of, in Turkey, 452; pests of, in U.S.A., 31, 165, 377; susceptibility of varieties of, to pests, 11, 46, 191; condition of European, not affecting infestation by Phylloxera, 112; insecticides causing injury to, 45, 103. (See Grapes.)

Vine Moths (see Clysia ambiguella, Phalaenoides glycinae and Poly-

chrosis botrana).

Vinegar, in baits for *Cydia* and vine moths, **89**, **293**, **494**, **496**, **582**.

vinitor, Nysius.

vinula, Dicranura.

violacea, Necrobia.

Virachola bimaculata, food-plants and control of, in Sierra Leone, 271.

Virachola isocrates, probable parasite of, on pomegranate in India, 385.

virens, Coscineuta.

virescens, Heliothis.

virgata, Ferrisiana (Pseudococcus); Nephrotoma.

Virginia, apple Aphids in, 141, 533; pests of lima beans in, 595; miscellaneous pests in, 380, 530, 599; Dendroctorus

frontalis on Pinus taeda in, 377, **599**; Ophiostoma ulmi on elms in, 378.

virginica, Diacrisia. viridana, Tortrix. viride, Piesma maculatum. viridescens, Eupteromalus. viridicata, Elaeochlora. viridicyaneus, Typophorus. viridigrisea, Empoasca. viridipes, Phymateus. viridis. Anomala (Euchlora);

Chermes; Cicadella (Tettigoniella); Coccus; Smynthurus.

viridissima, Palomena. viridula, Nezara.

Virus Diseases (in Plants*), relation of Aleurodids to, 94, 124, 125, **153, 186, 360, 493, 637**; relation of Aphids to, 83, 84, 87, 88, 125, 126, 157, 158, 186, 213, 226, 236, 267, 273, 274, 280, 289, 301, 302, 303, 333, 334, 365, 369, 370, 395, 448, 472, 493, 563, 578, 590, 591, 633, 634, 652, 658, 667, 671, 672, 686, 687, 688; relation of Capsids to, 84, 157, 163, 303, 555. 555; Heliothis assulta transmitting, 444; relation of Jassids 84, 157, 286, 303, to, 412, 536, 688; relation Perkinsiella vastatrix to, 667; relation of Piesma to, 163, 351, 352, 683; relation of Thysanoptera to, 360, 472, 546; negative experiments with other insects and, 84, 95, 636, 688; bibliographies and general papers on 324 hosts and vectors of, 280, 334, **439, 456, 624**; (in beet), legislation against introduction of, into Britain, 356.

visci, Diaspis. viteana, Polychrosis.

Vitex, Bostrychoplites cornutus in planks of, in Mauritius, 611.

vitifoliae, Phylloxera.

Vitis vinifera (see Vine, Grape). vitis, Eriophyes; Targionia. vitripennis, Brachymeria; Dicero-

procta; Mormoniella. vittalis, Nymphula. vittata, Cassida; Phyllotreta.

vittatum, Phaulacridium. vittatus, Prepodes.

vittula, Phyllotreta. Voandzeia subterranea, Apion ugandum on, in Nyasaland, 126.

volucris, Eupeodes. vorax, Anthrenus.

Voria ruralis, parasite of Vanessa cardui in Transcaucasia, 587. vulgaris, Cyrtogaster; Gryllotalpa (see G. gryllotalpa); Melolontha

(see M. melolontha); Pterostichus (Omaseus); Zenillia (Phryxe).

vulnerator, Pristomerus.

W.

wahlbergi, Chilocorus.

Walnut, pests of, in California, 261, 649, 650; symbiosis of Tremex with fungi in, in Germany, 401; Cydia pomonella on, in Italy, 4, 197; slightly attacked by Caliroa limacina in Tasmania, Parlatoria oleae on, Uzbekistan, 474.

Walnut, Black (see Juglans nigra). Walnut-shell Flour, Bruchus obtectus not affected by, 531; as a carrier for dinitro-o-cyclohexylphenol, 261, 650.

Washington, pests of forest and shade trees in, 499, 599; miscellaneous pests in, 525, 541; orchard pests in, 343, 525; pests of small fruits in, 178, 280, 380; Bruchus pisorum on peas in, 325, 456; pests of timber in, 78, 498, 499; wireworms in, 378, 418; beneficial insects and biological control in, 379, 499; catalogue of Arthropods of, 525.

Washingtonia filifera (filamentosa), Dinapate wrighti on, in California

and Mexico, 569.

Water, treatment of plant material with, against weevils, 291, 539; timber stored under, against Coleoptera, 397, 612; bamboo rendered unattractive to Dinoderus by, 181; (under pressure), against Leptocoris trivittatus, 374; Citrus fruit injured by treatment with, against Aonidiella aurantii,

Water, Hot, treatment of plants and planting material with, 396, 419, 605, 657, 665; other uses of, against pests, 10, 131, 359, 374.

Water Lily (see Nymphaea alba). Watermelon, not attacked by Diaphania hyalinata in Porto Rico, 501; Etiella zinckenella on, in Russia, 152.

Wax Moth (see Galleria mellonella).

^{*} These references are also indexed under the individual plants infected.

Wax Moth, Lesser (see Achroia grisella).

Pristiphora (Lygaeonewesmaeli, matus)

Wesmaëlius concinnus, predacious on Chermes spp. in Britain, 262, 263.

Wheat, wireworms on, in Britain, 96, 97; pests of, in Canada, 66, 176, 368, 600; locusts damaging, in China, 450, 451; pests of, in Czechoslovakia, 468; pests of, in Denmark, 117; pests of, in Germany, 252, 253; pests of, in Holland, 268, 269; Locusta migratoria damaging, in India, 560; Contarinia tritici on, in Italy, 3; pests of, in Japan, 208, 556: pests of, in Manchuria, 361; Dipterous pests of, in Morocco, 124, 691; pests of, in New Zealand, 547, 548; Nomadacris septemfasciata damaging, Nyasaland, 126; Syringopais temperatella on, in Palestine, 127; slightly damaged by grasshoppers in Queensland, 602; pests of, in Russian Union, 311, 312, 315, 453, 554, 677, 680; Thysanoptera on, in Sweden, 121; pests of, in U.S.A., 83, 280, 284, 327, 374, 375, 376, 378, 542, 599; type of injury to, by Rhynchota, 352, 353, 548, 554; insects and white ear in, 680; varieties and time of sowing of, in relation to pests, 4, 66, 83, 269, 312, 327, 374, 375, 378; naphthalene and superphosphate reducing germination of, 97; intersown with other crops to protect them from wireworms, 97, 674.

Wheat (Stored), pests of, and their control, 119, 120, 137, 196, 220, 305, 318, 319, 551, 585; Lasioderma serricorne not feeding on,

234.

Wheat Gall-midges (see Contarinia tritici and Sitodiplosis mosellana). Wheat Strawworm (see Harmolita

grandis).

Wheat Thrips (see Haplothrips tritici).

Wheat-stem Sawfly (see Cephus cinctus).

Wheat-stem Sawfly, Black Trachelus tabidus).

White Ear, in cereals and grasses, Arthropods and other factors associated with, 121, 151, 680.

White-fringed Beetle (see Naupastus leucoloma).

White-lined Sphinx (see Celerio lineata).

Whiteflies, and plant diseases, 94, 124, 125, 153, 360, 493, 637; natural enemies of, 265, 333, 385, 437; new species of, 637.

Whitefly, Cabbage (see Aleurodes

brassicae).

Whitefly, Citrus (see Dialeurodes citri).

Whitefly, Greenhouse (see Trialeurodes vaporariorum).

Willow (Salix), Rhabdophaga salici-perda on, in Britain, 627; pests of, in Germany, 6, 398; pests of, in Japan, 207, 440, 615; pests of, in Morocco, 196, 359; pests of, in Russian Union, 43, 152, 474; Lachnosterna implicita on, in Wisconsin, 32; Ceresa bubalus reared on, 390; attacked by Pyrausta nubilalis, 152.

Willow, Cricket-bat (see Salix alba

var. coerulea).

Wilt Disease, in Lepidoptera, 499, 612: disease in Heliothis armigera

resembling, 423.

Wind, relation of locust flights to, 33, 109, 554, 560, 646; relation of, to flight and spread of other insects, 99, 167, 245, 330, 336, 351, 367, 409, 470, 471, 519, 614, 633, 647.

Wine, in baits for Cydia and vine moths, 20, 494, 496, 582.

Moth (see Operophtera Winter brumata).

Winthemia quadripustulata, parasite of Monima cruda in Poland, 438; parasite of Protoparce spp. in U.S.A., 253.

Wireworm, Northern Grain (see Corymbites aeripennis destructor). Wireworm, Sand (see Horistonotus

uhleri).

Wireworms (Elaterids), problem of, in Britain, 96; survey of, in Siberia, 678; importance of, on potatoes in U.S.A., 418; causing white ear in oats, 680; ecology of, 96, 97, 501, 656, 679, 683; movements of, in soil, 96, 500; soil sampling for, 656, 679; measures against, 97, 177, 378, 501, 656, 683; wheat planted among crops as a protection against, 97, 674; value of trapping adults of, 410; reactions of, to arsenicals, 28; predacious on other insects, 105, 239, 398, 660, 670.

Wisconsin, miscellaneous pests in, 31-33, 323, 378; Carabid predacious on *Lachnosterna* in, 423.

woodi, Acarapis.

Woodpeckers, destroying Scolytus sulcatus in U.S.A., 78; protection of Formica rufa against, 564. Wool Grease, uses of, in sprays, 426. Woollen Goods (see Textiles).

Woolly Apple Aphis (see Eriosoma

lanigerum).

wrighti, Dinapate.

X.

X-rays, for detecting pests in seeds, 147, 440.

Xanthium pungens, insects introduced into Australia against, 546.

xanthodes, Dacus (Notodacus).

xanthomelaena, Galerucella (see G. luteola).

xanthomelaenae, Tetrastichus. xanthophilum, Monalonion.

Xenosternum ornigis, parasite of Gelechia cercerisella in Kansas, 376.

Xeris (see Sirex).

Xerophilaphis scorzonerae, measures against, on rubber-producing plants in Russia, 44, 45; ant associated with, 44.

Xiphydria prolongata (dromedarius), on willow in Britain, 562.

xuthus, Papilio.

Xylamon Preparations, tests and action of, against *Hylotrupes bajulus* in timber, 110, 202, 520; against *Ips typographus*, 123; penetration of, into tracheae of insects, 580; question of toxicity of vapour of, 202, 520, 580.

Xyleborus, on teak in Burma, 693; not reducing yield of coffee in

Java, 522, 523.

Xyleborus badius, investigation of damage by, to puncheons exported from Br. Guiana, 328,524.

Xyleborus fornicatus, measures against, on tea in Formosa, 340.

Xyleborus morstatti (on avocado), in Fiji, 641; in Mauritius, 611.

Xyleborus myristicae, sp. n., on nutmeg in Sumatra, 512.

Xyleborus perforans, in Jamaica, 181, 239; on coconut, 239; measures against, in furniture, 181.

Xyleborus todo, Rhizophagus predacious on, on Abies in Japan, 339.

Xyleutes ceramicus, bionomics of, on teak and Gmelina arborea in Burma, 692.

Xyleutes persona, bionomics of, on Cassia in Burma, 693.

Xylia dolabriformis, pests of, in Burma, 693.

Xylion adustus, in timber in S.

Africa, 60.

Xylocopa spp., bionomics and control of, in timber, etc., in India, 337.

Xylodrepa quadripunctata (see

Thanatophilus).

Xylomyges eridania, spray against, on cotton in Barbados, 661; on vegetables in Florida, 287; disease in, 81; tests of baits for, 287; used in experiments with insecticides, 241, 421, 524.

Xylopsocus sellatus, in timber of Acacia melanoxylon in S. Africa,

60.

xylosteana, Tortrix (Cacoecia). xylostei, Hyadaphis (see H. sii).

Xylotrechus quadripes, bionomics and control of, on coffee in Indo-China, 49.

Xystrocera globosa, on Acacia mollissima in Formosa, 558.

Y.

Yeast, in food for *Dacus* and *Heliothis*, **345**, **346**, **427**; increasing populations of *Tribolium* confusum in maize meal, **80**.

Yellow-headed Spruce Sawfly (see Pikonema alaskensis).

ynca, Amerrhinus. ypsilon, Agrotis.

Z.

Zabrus tenebrioides, on rye in Czechoslovakia, 468.

Zamacra juglansiaria, bionomics of, on mulberry in Japan, 555.

Zanthoxylum, Papilio xuthus on, in China, 360; resistance of timber of, to termites in Porto Rico, 465.

Zanzibar, virus diseases of cassava in, **95**.

Zea mays (see Maize).

zehntneri, Elasmus.

Zele, parasite of Laphygma exempta in Kenya, 666; taxonomy and new species of, 263.

Zele calcarator, parasite of Erannis in Crete, 172.

Zelus cervicalis, predacious on Horistonotus uhleri in S. Carolina, 500.

Zenillia cosmophilae, sp. n., hosts of, in Queensland, 57.

Zenillia lucorum, parasite of Lymantria monacha in Germany, 581.

Zenillia noctuae, sp. n., hosts of, in Queensland, 57.

Zenillia picta, parasite of Plectoptera

reflexa in India, 612. Zenillia vulgaris, parasite of Sphinx

pinastri in Poland, 677.

Zeuxia cinerea, parasite of Lixus junci in Morocco, 283.

Zeuzera coffeae, on Xylia dolabriformis in Burma, 693; on tea in Formosa, 340; on coffee in India, 577; measures against, 340, 577.

Zeuzera pyrina, on fruit trees in Czechoslovakia, 469; on hazel in Greece, 452.

Zicrona coerulea, predacious on Leptinotarsa decemlineata in France, 107.

Zinc Arsenate, use of, against wireworms, 177.

Zinc Arsenite, tests of, in baits for grasshoppers, 134.

Zinc Chloride, unsatisfactory against Lyctus brunneus, 510.

Zinc Cyanamide, tests of, in baits for grasshoppers, 134.

Zinc Hydroxide, as a corrective for foliage injury by calcium arsenate, 373.

Zinc Phosphide, tests of, in baits for grasshoppers, 134. (See Phosphides.)

Zinc Sulphate, and lime, as adhesive for thiuram sulphide sprays, **532**. zinckenella, Etiella.

Zinckenia fascialis (see Hymenia recurvalis).

Zinnia, Myzus persicae on, in S. Rhodesia, 125; possible reservoir of tobacco rosette, 125.

Zizera (Zizina) labradus var. mangoensis, on beans in Fiji, 641.

Zizyphus, Carpomyia spp. intercepted in fruits of, in Russian Union, 439.

Zizyphus jujuba, pests of, in India, 129, 503.

zizyphus, Parlatoria.

zlocistii, Lepidosaphes (Coccomytilus).

zonatus, Dacus (Chaetodacus).

zonellus, Chilo.

Zootermopsis angusticollis, in rotten wood in U.S.A., 498; bacterial diseases of, 597.

Zygina parvula (see Erythroneura). Zyklon, as fumigant against cereal pests, 120, 400.

NOTICES.

Secretaries of Societies and Editors of Journals willing to exchange their publications with those of the Institute are requested to communicate with the Assistant Director.

The Annual Subscription, in advance, to the Review, Series A (Agricultural), is 30s. post free; Series B (Medical and Veterinary), 15s. post free. Subscriptions to the current Volume received after 30th June in each year will be charged at the published price, viz.: Series A, 36s.; Series B, 18s. Prices of Back Volumes on application.

Orders and Subscriptions should be sent direct to the Assistant Director, Imperial Institute of Entomology, 41, Queen's Gate, London, S.W.7, or through a bookseller.